

Random walks and diffusion on networks

Last updated: 2023/08/22, 11:48:21 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D
CSYS/MATH 6701, 6713, & a pretend number,
2023–2024 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
Santa Fe Institute | University of Vermont



Licensed under the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

Where is Barry?

- Consider simple undirected, ergodic (strongly connected) networks.
- As usual, represent network by **adjacency matrix** A where

$$a_{ij} = 1 \text{ if } i \text{ has an edge leading to } j,$$

$$a_{ij} = 0 \text{ otherwise.}$$

- Barry is at node j at time t with probability $p_j(t)$.
- In the next time step, he **randomly lurches** toward one of j 's neighbors.
- Barry arrives at node i from node j with probability $\frac{1}{k_j}$ if an edge connects j to i .
- Equation-wise:

$$p_i(t+1) = \sum_{j=1}^n \frac{1}{k_j} a_{ji} p_j(t).$$

where k_j is j 's degree. Note: $k_i = \sum_{j=1}^n a_{ij}$.

Where is Barry?

- By inspection, we see that

$$\vec{p}(\infty) = \frac{1}{\sum_{i=1}^n k_i} \vec{k}$$

- satisfies $\vec{p}(\infty) = A^T K^{-1} \vec{p}(\infty)$ with eigenvalue 1.
- We will find Barry at node i with probability proportional to its degree k_i .
- Beautiful implication: probability of finding Barry travelling along any edge is **uniform**.
- Diffusion in real space smooths things out.
- On networks, uniformity occurs on edges.
- So in fact, diffusion in real space is **about the edges too** but we just don't see that.

Outline

Random walks on networks

Inebriation and diffusion:

- Excellent observation:** The same equation applies for stuff moving around a network, such that at each time step all material at node i is sent to its neighbors.
- $x_i(t)$ = amount of stuff at node i at time t .

$$x_i(t+1) = \sum_{j=1}^n \frac{1}{k_j} a_{ji} x_j(t).$$

- Random walking is equivalent to [diffusion](#) ↗.

Other pieces:

- Goodness: $A^T K^{-1}$ is similar to a real symmetric matrix if $A = A^T$.
- Consider the transformation $M = K^{-1/2}$:

$$K^{-1/2} A^T K^{-1} K^{1/2} = K^{-1/2} A^T K^{-1/2}.$$

- Since $A^T = A$, we have

$$(K^{-1/2} A K^{-1/2})^T = K^{-1/2} A K^{-1/2}.$$

- Upshot: $A^T K^{-1} = A K^{-1}$ has real eigenvalues and a complete set of orthogonal eigenvectors.
- Can also show that maximum eigenvalue magnitude is indeed 1.

Random walks on networks—basics:

- Imagine a single random walker moving around on a network.
- At $t = 0$, start walker at node j and take time to be discrete.
- Q:** What's the long term probability distribution for where the walker will be?
- Define $p_i(t)$ as the probability that at time step t , our walker is at node i .
- We want to characterize the evolution of $\vec{p}(t)$.
- First task: connect $\vec{p}(t+1)$ to $\vec{p}(t)$.
- Let's call our walker **Barry**.
- Unfortunately for Barry, he lives on a high dimensional graph and is far from home.
- Worse still: Barry is **texting**.

Where is Barry?

- Linear algebra-based excitement:

$$p_i(t+1) = \sum_{j=1}^n a_{ji} \frac{1}{k_j} p_j(t)$$
 is more usefully viewed as

$$\vec{p}(t+1) = A^T K^{-1} \vec{p}(t)$$
 where $[K_{ij}] = [\delta_{ij} k_i]$ has node degrees on the main diagonal and zeros everywhere else.
- So... we need to find the **dominant eigenvalue** of $A^T K^{-1}$.
- Expect this eigenvalue will be 1 (doesn't make sense for total probability to change).
- The corresponding eigenvector will be the limiting probability distribution (or invariant measure).
- Extra concerns: multiplicity of eigenvalue = 1, and network connectedness.