

# Generating Functions and Networks

Last updated: 2023/08/22, 11:48:21 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D  
CSYS/MATH 6701, 6713, & a pretend number,  
2023-2024 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center  
Santa Fe Institute | University of Vermont



Licensed under the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

The PoCSverse  
Generating  
Functions and  
Networks  
1 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Simple examples:

Rolling dice and flipping coins:

$p_k^{(\square)} = \Pr(\text{throwing a } k) = 1/6 \text{ where } k = 1, 2, \dots, 6.$

$$F^{(\square)}(x) = \sum_{k=1}^6 p_k^{(\square)} x^k = \frac{1}{6}(x + x^2 + x^3 + x^4 + x^5 + x^6).$$

$p_0^{(\text{coin})} = \Pr(\text{head}) = 1/2, p_1^{(\text{coin})} = \Pr(\text{tail}) = 1/2.$

$$F^{(\text{coin})}(x) = p_0^{(\text{coin})}x^0 + p_1^{(\text{coin})}x^1 = \frac{1}{2}(1 + x).$$

A generating function for a probability distribution is called a **Probability Generating Function (p.g.f.)**.

We'll come back to these simple examples as we derive various delicious properties of generating functions.

The PoCSverse  
Generating  
Functions and  
Networks  
7 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

Useful pieces for probability distributions:

Normalization:  $F(1) = 1$

First moment:  $\langle k \rangle = F'(1)$

Higher moments:  $\langle k^n \rangle = \left( x \frac{d}{dx} \right)^n F(x) \Big|_{x=1}$

kth element of sequence (general):  $P_k = \frac{1}{k!} \frac{d^k}{dx^k} F(x) \Big|_{x=0}$

The PoCSverse  
Generating  
Functions and  
Networks  
11 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Outline

### Generating Functions

- Definitions
- Basic Properties
- Giant Component Condition
- Component sizes
- Useful results
- Size of the Giant Component
- A few examples
- Average Component Size

### References

The PoCSverse  
Generating  
Functions and  
Networks  
2 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Example

Take a degree distribution with exponential decay:

$$P_k = ce^{-\lambda k}$$

where geometrically, we have  $c = 1 - e^{-\lambda}$

The generating function for this distribution is

$$F(x) = \sum_{k=0}^{\infty} P_k x^k = \sum_{k=0}^{\infty} ce^{-\lambda k} x^k = \frac{c}{1 - xe^{-\lambda}}.$$

Notice that  $F(1) = c/(1 - e^{-\lambda}) = 1.$

For probability distributions, we must always have  $F(1) = 1$  since

$$F(1) = \sum_{k=0}^{\infty} P_k 1^k = \sum_{k=0}^{\infty} P_k = 1.$$

Check die and coin p.g.f.'s.

The PoCSverse  
Generating  
Functions and  
Networks  
8 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

A beautiful, fundamental thing:

The generating function for the sum of two random variables

$$W = U + V$$

is

$$F_W(x) = F_U(x)F_V(x).$$

Convolv yourself with Convolutions: [Insert assignment question](#)

- Try with die and coin p.g.f.'s.
- Add two coins (tail=0, head=1).
  - Add two dice.
  - Add a coin flip to one die roll.

The PoCSverse  
Generating  
Functions and  
Networks  
12 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Generatingfunctionology<sup>[1]</sup>

**Idea:** Given a sequence  $a_0, a_1, a_2, \dots$ , associate each element with a distinct function or other mathematical object.

Well-chosen functions allow us to manipulate sequences and retrieve sequence elements.

### Definition:

The **generating function (g.f.)** for a sequence  $\{a_n\}$  is

$$F(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Roughly: transforms a vector in  $R^\infty$  into a function defined on  $R^1$ .

Related to Fourier, Laplace, Mellin, ...

The PoCSverse  
Generating  
Functions and  
Networks  
6 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Properties:

Average degree:

$$\begin{aligned} \langle k \rangle &= \sum_{k=0}^{\infty} k P_k = \sum_{k=0}^{\infty} k P_k x^{k-1} \Big|_{x=1} \\ &= \frac{d}{dx} F(x) \Big|_{x=1} = F'(1) \end{aligned}$$

In general, many calculations become simple, if a little abstract.

For our exponential example:

$$F'(x) = \frac{(1 - e^{-\lambda})e^{-\lambda}}{(1 - xe^{-\lambda})^2}.$$

$$\text{So: } \langle k \rangle = F'(1) = \frac{e^{-\lambda}}{(1 - e^{-\lambda})}.$$

Check for die and coin p.g.f.'s.

The PoCSverse  
Generating  
Functions and  
Networks  
10 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Edge-degree distribution

Recall our condition for a giant component:

$$\langle k \rangle_R = \frac{\langle k^2 \rangle - \langle k \rangle}{\langle k \rangle} > 1.$$

Let's re-express our condition in terms of generating functions.

We first need the g.f. for  $R_k$ .

We'll now use this notation:

$$F_P(x) \text{ is the g.f. for } P_k.$$

$$F_R(x) \text{ is the g.f. for } R_k.$$

Giant component condition in terms of g.f. is:

$$\langle k \rangle_R = F'_R(1) > 1.$$

Now find how  $F_R$  is related to  $F_P$  ...

The PoCSverse  
Generating  
Functions and  
Networks  
14 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
References

## Edge-degree distribution

☞ We have

$$F_R(x) = \sum_{k=0}^{\infty} R_k x^k = \sum_{k=0}^{\infty} \frac{(k+1)P_{k+1}}{\langle k \rangle} x^k.$$

Shift index to  $j = k + 1$  and pull out  $\frac{1}{\langle k \rangle}$ :

$$F_R(x) = \frac{1}{\langle k \rangle} \sum_{j=1}^{\infty} j P_j x^{j-1} = \frac{1}{\langle k \rangle} \sum_{j=1}^{\infty} P_j \frac{d}{dx} x^j$$


$$= \frac{1}{\langle k \rangle} \frac{d}{dx} \sum_{j=1}^{\infty} P_j x^j = \frac{1}{\langle k \rangle} \frac{d}{dx} (F_P(x) - P_0) = \frac{1}{\langle k \rangle} F'_P(x).$$

Finally, since  $\langle k \rangle = F'_P(1)$ ,

$$F_R(x) = \frac{F'_P(x)}{F'_P(1)}$$

The PoCVerse  
Generating Functions and Networks  
15 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting probabilities:



☞ Markov property of random networks connects  $\pi_n, \rho_n$ , and  $P_k$ .

The PoCVerse  
Generating Functions and Networks  
19 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Useful results we'll need for g.f.'s

### Sneaky Result 1:

☞ Consider two random variables  $U$  and  $V$  whose values may be  $0, 1, 2, \dots$

☞ Write probability distributions as  $U_k$  and  $V_k$  and g.f.'s as  $F_U$  and  $F_V$ .

☞ SR1: If a third random variable is defined as

$$W = \sum_{i=1}^U V^{(i)} \text{ with each } V^{(i)} \stackrel{d}{=} V$$

then

$$F_W(x) = F_U(F_V(x))$$

The PoCVerse  
Generating Functions and Networks  
23 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Edge-degree distribution

☞ Recall giant component condition is  $\langle k \rangle_R = F'_R(1) > 1$ .

☞ Since we have  $F_R(x) = F'_P(x)/F'_P(1)$ ,


$$F'_R(x) = \frac{F''_P(x)}{F'_P(1)}$$

☞ Setting  $x = 1$ , our condition becomes

$$\frac{F''_P(1)}{F'_P(1)} > 1$$

The PoCVerse  
Generating Functions and Networks  
16 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting probabilities:



☞ Markov property of random networks connects  $\rho_n$  and  $R_k$ .

The PoCVerse  
Generating Functions and Networks  
20 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Proof of SR1:

Write probability that variable  $W$  has value  $k$  as  $W_k$ .

$$W_k = \sum_{j=0}^{\infty} U_j \times \Pr(\text{sum of } j \text{ draws of variable } V = k)$$

$$= \sum_{j=0}^{\infty} U_j \sum_{\{i_1, i_2, \dots, i_j\}} V_{i_1} V_{i_2} \dots V_{i_j}$$

$i_1 + i_2 + \dots + i_j = k$

$$\therefore F_{W'}(x) = \sum_{k=0}^{\infty} W_k x^k = \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} U_j \sum_{\{i_1, i_2, \dots, i_j\}} V_{i_1} V_{i_2} \dots V_{i_j} x^k$$

$i_1 + i_2 + \dots + i_j = k$

$$= \sum_{j=0}^{\infty} U_j \sum_{k=0}^{\infty} \sum_{\{i_1, i_2, \dots, i_j\}} V_{i_1} x^{i_1} V_{i_2} x^{i_2} \dots V_{i_j} x^{i_j}$$

$i_1 + i_2 + \dots + i_j = k$

The PoCVerse  
Generating Functions and Networks  
25 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Size distributions

To figure out the **size of the largest component** ( $S_1$ ), we need more resolution on component sizes.

Definitions:

☞  $\pi_n$  = probability that a random node belongs to a finite component of size  $n < \infty$ .

☞  $\rho_n$  = probability that a random end of a random link leads to a finite subcomponent of size  $n < \infty$ .

Local-global connection:

$$P_k, R_k \Leftrightarrow \pi_n, \rho_n$$

neighbors  $\Leftrightarrow$  components

The PoCVerse  
Generating Functions and Networks  
18 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## G.f.'s for component size distributions:

☞

$$F_{\pi}(x) = \sum_{n=0}^{\infty} \pi_n x^n \text{ and } F_{\rho}(x) = \sum_{n=0}^{\infty} \rho_n x^n$$

## The largest component:

☞ Subtle key:  $F_{\pi}(1)$  is the probability that a node belongs to a **finite** component.

☞ Therefore:  $S_1 = 1 - F_{\pi}(1)$ .

## Our mission, which we accept:

☞ Determine and connect the four generating functions

$$F_P, F_R, F_{\pi}, \text{ and } F_{\rho}.$$

The PoCVerse  
Generating Functions and Networks  
21 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Proof of SR1:

With some concentration, observe:

$$F_W(x) = \sum_{j=0}^{\infty} U_j \sum_{k=0}^{\infty} \sum_{\{i_1, i_2, \dots, i_j\}} V_{i_1} x^{i_1} V_{i_2} x^{i_2} \dots V_{i_j} x^{i_j}$$

$i_1 + i_2 + \dots + i_j = k$

$$\underbrace{x^k \text{ piece of } \left( \sum_{i'=0}^{\infty} V_{i'} x^{i'} \right)^j}_{\left( \sum_{i'=0}^{\infty} V_{i'} x^{i'} \right)^j} = (F_V(x))^j$$

$$= \sum_{j=0}^{\infty} U_j (F_V(x))^j$$

$$= F_U(F_V(x))$$

The PoCVerse  
Generating Functions and Networks  
26 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

☞ Alternate, groovier proof in the accompanying assignment.

## Useful results we'll need for g.f.'s

### Sneaky Result 2:

Start with a random variable  $U$  with distribution  $U_k$  ( $k = 0, 1, 2, \dots$ )

SR2: If a second random variable is defined as

$$V = U + 1 \text{ then } \boxed{F_V(x) = xF_U(x)}$$

Reason:  $V_k = U_{k-1}$  for  $k \geq 1$  and  $V_0 = 0$ .

$$\therefore F_V(x) = \sum_{k=0}^{\infty} V_k x^k = \sum_{k=1}^{\infty} U_{k-1} x^k$$

$$= x \sum_{j=0}^{\infty} U_j x^j = xF_U(x).$$

The PoCVerse  
Generating Functions and Networks  
27 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting generating functions:

$\pi_n$  = probability that a random node belongs to a finite component of size  $n$

$$= \sum_{k=0}^{\infty} P_k \times \Pr \left( \begin{array}{l} \text{sum of sizes of subcomponents} \\ \text{at end of } k \text{ random links} = n - 1 \end{array} \right)$$



Therefore: 
$$F_{\pi}(x) = \boxed{x \underbrace{F_P(F_{\rho}(x))}_{\text{SR1}}}_{\text{SR2}}$$

Extra factor of  $x$  accounts for random node itself.

The PoCVerse  
Generating Functions and Networks  
31 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting generating functions:

We now have two functional equations connecting our generating functions:

$$F_{\pi}(x) = xF_P(F_{\rho}(x)) \text{ and } F_{\rho}(x) = xF_R(F_{\rho}(x))$$

Taking stock: We know  $F_P(x)$  and  $F_R(x) = F'_P(x)/F'_P(1)$ .

We first untangle the second equation to find  $F_{\rho}$

We can do this because it **only involves**  $F_{\rho}$  and  $F_R$ .

The first equation then immediately gives us  $F_{\pi}$  in terms of  $F_{\rho}$  and  $F_R$ .

The PoCVerse  
Generating Functions and Networks  
34 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Useful results we'll need for g.f.'s

### Generalization of SR2:

(1) If  $V = U + i$  then

$$F_V(x) = x^i F_U(x).$$


(2) If  $V = U - i$  then

$$F_V(x) = x^{-i} F_U(x)$$

$$= x^{-i} \sum_{k=0}^{\infty} U_k x^k$$

The PoCVerse  
Generating Functions and Networks  
28 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting generating functions:



Relate  $\rho_n$  to  $R_k$  and  $\rho_n$  through one step of recursion.

The PoCVerse  
Generating Functions and Networks  
32 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Component sizes

Remembering vaguely what we are doing:

Finding  $F_{\pi}$  to obtain the **fractional size of the largest component**  $S_1 = 1 - F_{\pi}(1)$ .

Set  $x = 1$  in our two equations:

$$F_{\pi}(1) = F_P(F_{\rho}(1)) \text{ and } F_{\rho}(1) = F_R(F_{\rho}(1))$$


Solve second equation numerically for  $F_{\rho}(1)$ .

Plug  $F_{\rho}(1)$  into first equation to obtain  $F_{\pi}(1)$ .

The PoCVerse  
Generating Functions and Networks  
35 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting generating functions:

Goal: figure out forms of the component generating functions,  $F_{\pi}$  and  $F_{\rho}$ .



Relate  $\pi_n$  to  $P_k$  and  $\rho_n$  through one step of recursion.

The PoCVerse  
Generating Functions and Networks  
30 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Connecting generating functions:

$\rho_n$  = probability that a random link leads to a finite subcomponent of size  $n$ .

Invoke one step of recursion:

$\rho_n$  = probability that in following a random edge, the outgoing edges of the node reached lead to finite subcomponents of combined size  $n - 1$ ,

$$= \sum_{k=0}^{\infty} R_k \times \Pr \left( \begin{array}{l} \text{sum of sizes of subcomponents} \\ \text{at end of } k \text{ random links} = n - 1 \end{array} \right)$$



Therefore: 
$$F_{\rho}(x) = \boxed{x \underbrace{F_R(F_{\rho}(x))}_{\text{SR1}}}_{\text{SR2}}$$

Again, extra factor of  $x$  accounts for random node itself.

The PoCVerse  
Generating Functions and Networks  
33 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Component sizes

Example: Standard random graphs.

We can show  $F_P(x) = e^{-\langle k \rangle (1-x)}$

$$\Rightarrow F_R(x) = F'_P(x)/F'_P(1)$$

$$= \langle k \rangle e^{-\langle k \rangle (1-x)} / \langle k \rangle e^{-\langle k \rangle (1-x')} \Big|_{x'=1}$$

$$= e^{-\langle k \rangle (1-x)} = F_P(x) \quad \dots \text{aha!}$$

RHS's of our two equations are the same.

So  $F_{\pi}(x) = F_{\rho}(x) = xF_R(F_{\rho}(x)) = xF_R(F_{\pi}(x))$

Consistent with how our dirty (but wrong) trick worked earlier ...

$\pi_n = \rho_n$  just as  $P_k = R_k$ .

The PoCVerse  
Generating Functions and Networks  
36 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

## Component sizes

We are down to


$$F_{\pi}(x) = xF_R(F_{\pi}(x)) \text{ and } F_R(x) = e^{-(k)(1-x)}.$$

$$\therefore F_{\pi}(x) = xe^{-(k)(1-F_{\pi}(x))}$$

We're first after  $S_1 = 1 - F_{\pi}(1)$  so set  $x = 1$  and replace  $F_{\pi}(1)$  by  $1 - S_1$ :

$$1 - S_1 = e^{-(k)S_1}$$

$$\text{Or: } \langle k \rangle = \frac{1}{S_1} \ln \frac{1}{1 - S_1}$$



Just as we found with our dirty trick ...

Again, we (usually) have to resort to numerics ...

A few simple random networks to contemplate and play around with:

Notation: The Kronecker delta function  $\delta_{ij} = 1$  if  $i = j$  and 0 otherwise.

$$P_k = \delta_{k1}.$$

$$P_k = \delta_{k2}.$$

$$P_k = \delta_{k3}.$$

$$P_k = \delta_{kk'} \text{ for some fixed } k' \geq 0.$$

$$P_k = \frac{1}{2}\delta_{k1} + \frac{1}{2}\delta_{k3}.$$

$$P_k = a\delta_{k1} + (1-a)\delta_{k3}, \text{ with } 0 \leq a \leq 1.$$

$$P_k = \frac{1}{2}\delta_{k1} + \frac{1}{2}\delta_{kk'} \text{ for some fixed } k' \geq 2.$$

$$P_k = a\delta_{k1} + (1-a)\delta_{kk'} \text{ for some fixed } k' \geq 2 \text{ with } 0 \leq a \leq 1.$$

A joyful example  $\square$ :

$$P_k = \frac{1}{2}\delta_{k1} + \frac{1}{2}\delta_{k3}.$$

We find (two ways):  $R_k = \frac{1}{4}\delta_{k0} + \frac{3}{4}\delta_{k2}$ .

A giant component exists because:

$$\langle k \rangle_R = 0 \times 1/4 + 2 \times 3/4 = 3/2 > 1.$$

Generating functions for  $P_k$  and  $R_k$ :

$$F_P(x) = \frac{1}{2}x + \frac{1}{2}x^3 \text{ and } F_R(x) = \frac{1}{4}x^0 + \frac{3}{4}x^2$$

Check for goodness:

$$F_R(x) = F'_P(x)/F'_P(1) \text{ and } F_P(1) = F_R(1) = 1.$$

$$F'_P(1) = \langle k \rangle_P = 2 \text{ and } F'_R(1) = \langle k \rangle_R = \frac{3}{2}.$$


Things to figure out: Component size generating functions for  $\pi_n$  and  $\rho_n$ , and the size of the giant component.

The PoCSVerse  
Generating Functions and Networks  
37 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

Find  $F_{\rho}(x)$  first:

We know:

$$F_{\rho}(x) = xF_R(F_{\rho}(x)).$$



The PoCSVerse  
Generating Functions and Networks  
41 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

Let's define a binomial for arbitrary  $\theta$  and  $k = 0, 1, 2, \dots$ :

$$\binom{\theta}{k} = \frac{\Gamma(\theta + 1)}{\Gamma(k + 1)\Gamma(\theta - k + 1)}$$

For  $\theta = \frac{1}{2}$ , we have:

$$(1 + z)^{\frac{1}{2}} = \binom{\frac{1}{2}}{0} z^0 + \binom{\frac{1}{2}}{1} z^1 + \binom{\frac{1}{2}}{2} z^2 + \dots$$

$$= \frac{\Gamma(\frac{3}{2})}{\Gamma(1)\Gamma(\frac{3}{2})} z^0 + \frac{\Gamma(\frac{3}{2})}{\Gamma(2)\Gamma(\frac{1}{2})} z^1 + \frac{\Gamma(\frac{3}{2})}{\Gamma(3)\Gamma(-\frac{1}{2})} z^2 + \dots$$

$$= 1 + \frac{1}{2}z - \frac{1}{8}z^2 + \frac{1}{16}z^3 - \dots$$

where we've used  $\Gamma(x + 1) = x\Gamma(x)$  and noted that

$$\Gamma(\frac{1}{2}) = \frac{\sqrt{\pi}}{2}.$$

Note:  $(1 + z)^{\theta} \sim 1 + \theta z$  always.

Totally psyched, we go back to here:

$$F_{\rho}(x) = \frac{2}{3x} \left( 1 - \sqrt{1 - \frac{3}{4}x^2} \right).$$

Setting  $z = -\frac{3}{4}x^2$  and expanding, we have:

$$F_{\rho}(x) =$$

$$\frac{2}{3x} \left( 1 - \left[ 1 + \frac{1}{2} \left( -\frac{3}{4}x^2 \right)^1 - \frac{1}{8} \left( -\frac{3}{4}x^2 \right)^2 + \frac{1}{16} \left( -\frac{3}{4}x^2 \right)^3 + \dots \right] \right)$$

Giving:

$$F_{\rho}(x) = \sum_{n=0}^{\infty} \rho_n x^n =$$

$$\frac{1}{4}x + \frac{3}{64}x^3 + \frac{9}{512}x^5 + \dots + \frac{2}{3} \left( \frac{3}{4} \right)^k \frac{(-1)^{k+1} \Gamma(\frac{3}{2})}{\Gamma(k+1)\Gamma(\frac{3}{2}-k)} x^{2k-1} + \dots$$

Do odd powers make sense?

We can now find  $F_{\pi}(x)$  with:

$$F_{\pi}(x) = xF_P(F_{\rho}(x))$$

$$= x \frac{1}{2} \left( (F_{\rho}(x))^1 + (F_{\rho}(x))^3 \right)$$

$$= x \frac{1}{2} \left[ \frac{2}{3x} \left( 1 - \sqrt{1 - \frac{3}{4}x^2} \right) + \frac{2^3}{(3x)^3} \left( 1 - \sqrt{1 - \frac{3}{4}x^2} \right)^3 \right].$$

Delicious.

In principle, we can now extract all the  $\pi_n$ .

But let's just find the size of the giant component.

The PoCSVerse  
Generating Functions and Networks  
39 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

Sticking things in things, we have:

$$F_{\rho}(x) = x \left( \frac{1}{4} + \frac{3}{4} [F_{\rho}(x)]^2 \right).$$

Rearranging:

$$3x [F_{\rho}(x)]^2 - 4F_{\rho}(x) + x = 0.$$

Please and thank you:

$$F_{\rho}(x) = \frac{2}{3x} \left( 1 \pm \sqrt{1 - \frac{3}{4}x^2} \right)$$

Time for a Taylor series expansion.

The promise: non-negative powers of  $x$  with non-negative coefficients.

First: which sign do we take?

The PoCSVerse  
Generating Functions and Networks  
42 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

Because  $\rho_n$  is a probability distribution, we know

$$F_{\rho}(1) \leq 1 \text{ and } F_{\rho}(x) \leq 1 \text{ for } 0 \leq x \leq 1.$$

Thinking about the limit  $x \rightarrow 0$  in

$$F_{\rho}(x) = \frac{2}{3x} \left( 1 \pm \sqrt{1 - \frac{3}{4}x^2} \right),$$

we see that the positive sign solution blows to smithereens, and the negative one is okay.

So we must have:

$$F_{\rho}(x) = \frac{2}{3x} \left( 1 - \sqrt{1 - \frac{3}{4}x^2} \right),$$

We can now deploy the Taylor expansion:

$$(1 + z)^{\theta} = \binom{\theta}{0} z^0 + \binom{\theta}{1} z^1 + \binom{\theta}{2} z^2 + \binom{\theta}{3} z^3 + \dots$$

The PoCSVerse  
Generating Functions and Networks  
44 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

The PoCSVerse  
Generating Functions and Networks  
46 of 58  
Generating Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant Component  
A few examples  
Average Component Size  
References

First, we need  $F_\rho(1)$ :

$$F_\rho(x)|_{x=1} = \frac{2}{3 \cdot 1} \left( 1 - \sqrt{1 - \frac{3}{4}1^2} \right) = \frac{1}{3}.$$

This is the probability that a random edge leads to a sub-component of finite size.



Next:

$$F_\pi(1) = 1 - F_P(F_\rho(1)) = F_P\left(\frac{1}{3}\right) = \frac{1}{2} \cdot \frac{1}{3} + \frac{1}{2} \left(\frac{1}{3}\right)^3 = \frac{5}{27}.$$

This is the probability that a random chosen node belongs to a finite component.

Finally, we have

$$S_1 = 1 - F_\pi(1) = 1 - \frac{5}{27} = \frac{22}{27}.$$



## Average component size

Next: find **average size** of **finite** components  $\langle n \rangle$ .

Using standard G.F. result:  $\langle n \rangle = F'_\pi(1)$ .

Try to avoid finding  $F_\pi(x)$  ...

Starting from  $F_\pi(x) = xF_P(F_\rho(x))$ , we differentiate:

$$F'_\pi(x) = F_P(F_\rho(x)) + xF'_P(x)F'_P(F_\rho(x))$$

While  $F_\rho(x) = xF_R(F_\rho(x))$  gives

$$F'_\rho(x) = F_R(F_\rho(x)) + xF'_R(x)F'_R(F_\rho(x))$$

Now set  $x = 1$  in both equations.

We solve the second equation for  $F'_\rho(1)$  (we must already have  $F_\rho(1)$ ).

Plug  $F'_\rho(1)$  and  $F_\rho(1)$  into first equation to find  $F'_\pi(1)$ .

## Average component size

**Example:** Standard random graphs.

Use fact that  $F_P = F_R$  and  $F_\pi = F_\rho$ .

Two differentiated equations reduce to only one:

$$F'_\pi(x) = F_P(F_\pi(x)) + xF'_P(x)F'_P(F_\pi(x))$$

$$\text{Rearrange: } F'_\pi(x) = \frac{F_P(F_\pi(x))}{1 - xF'_P(F_\pi(x))}$$

Simplify denominator using  $F'_P(x) = \langle k \rangle F_P(x)$

Replace  $F_P(F_\pi(x))$  using  $F_\pi(x) = xF_P(F_\pi(x))$ .

Set  $x = 1$  and replace  $F_\pi(1)$  with  $1 - S_1$ .

$$\text{End result: } \langle n \rangle = F'_\pi(1) = \frac{(1 - S_1)}{1 - \langle k \rangle(1 - S_1)}$$

## Average component size

Our result for standard random networks:

$$\langle n \rangle = F'_\pi(1) = \frac{(1 - S_1)}{1 - \langle k \rangle(1 - S_1)}$$

Recall that  $\langle k \rangle = 1$  is the critical value of average degree for standard random networks.

Look at what happens when we increase  $\langle k \rangle$  to 1 from below.

We have  $S_1 = 0$  for all  $\langle k \rangle < 1$  so

$$\langle n \rangle = \frac{1}{1 - \langle k \rangle}$$

This blows up as  $\langle k \rangle \rightarrow 1$ .

**Reason:** we have a power law distribution of component sizes at  $\langle k \rangle = 1$ .

Typical critical point behavior ...

Let's return to our example:  $P_k = \frac{1}{2}\delta_{k1} + \frac{1}{2}\delta_{k3}$ .

We're after:

$$\langle n \rangle = F'_\pi(1) = F_P(F_\rho(1)) + F'_P(1)F'_P(F_\rho(1))$$

where we first need to compute

$$F'_\rho(1) = F_R(F_\rho(1)) + F'_R(1)F'_R(F_\rho(1)).$$

Place stick between teeth, and recall that we have:

$$F_P(x) = \frac{1}{2}x + \frac{1}{2}x^3 \text{ and } F_R(x) = \frac{1}{4}x^0 + \frac{3}{4}x^2.$$

Differentiation gives us:

$$F'_P(x) = \frac{1}{2} + \frac{3}{2}x^2 \text{ and } F'_R(x) = \frac{3}{2}x.$$

We bite harder and use  $F_\rho(1) = \frac{1}{3}$  to find:

$$\begin{aligned} F'_\rho(1) &= F_R(F_\rho(1)) + F'_R(1)F'_R(F_\rho(1)) \\ &= F_R\left(\frac{1}{3}\right) + F'_R(1)F'_R\left(\frac{1}{3}\right) \\ &= \frac{1}{4} + \frac{3}{4} \frac{1}{3^2} + F'_R(1) \frac{3}{2} \frac{1}{3}. \end{aligned}$$

After some reallocation of objects, we have  $F'_\rho(1) = \frac{13}{2}$ .

Finally:

$$\begin{aligned} \langle n \rangle &= F'_\pi(1) = F_P\left(\frac{1}{3}\right) + \frac{13}{2}F'_P\left(\frac{1}{3}\right) \\ &= \frac{1}{2} \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{1}{3^3} + \frac{13}{2} \left( \frac{1}{2} + \frac{3}{2} \frac{1}{3^2} \right) = \frac{5}{27} + \frac{13}{3} = \frac{122}{27}. \end{aligned}$$

So, kinda small.

## Average component size

Limits of  $\langle k \rangle = 0$  and  $\infty$  make sense for

$$\langle n \rangle = F'_\pi(1) = \frac{(1 - S_1)}{1 - \langle k \rangle(1 - S_1)}$$

As  $\langle k \rangle \rightarrow 0$ ,  $S_1 = 0$ , and  $\langle n \rangle \rightarrow 1$ .

All nodes are isolated.

As  $\langle k \rangle \rightarrow \infty$ ,  $S_1 \rightarrow 1$  and  $\langle n \rangle \rightarrow 0$ .

No nodes are outside of the giant component.

Extra on largest component size:

For  $\langle k \rangle = 1$ ,  $S_1 \sim N^{2/3}/N$ .

For  $\langle k \rangle < 1$ ,  $S_1 \sim (\log N)/N$ .

## Nutshell

Generating functions allow us to strangely calculate features of random networks.

They're a bit scary and magical.


Generating functions can be useful for contagion.

But: For the big results, more direct, physics-bearing calculations are possible.

# References I

[1] H. S. Wilf.

Generatingfunctionology.

A K Peters, Natick, MA, 3rd edition, 2006. pdf 

The PoCVerse  
**Generating  
Functions and  
Networks**  
58 of 58  
Generating  
Functions  
Definitions  
Basic Properties  
Giant Component  
Condition  
Component sizes  
Useful results  
Size of the Giant  
Component  
A few examples  
Average Component Size  
**References**