Lognormals and friends

Last updated: 2023/08/22, 11:48:25 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D CSYS/MATH 6701, 6713, & a pretend number, 2023–2024| @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
Santa Fe Institute | University of Vermont

These slides are brought to you by:

These slides are also brought to you by:

Special Guest Executive Producer

On Instagram at pratchett_the_cat

Outline


THE SUN
THE GOLDEN AGE OF REDUCTIONISM
THE MANIFESTO
THE WELL OF SCALES
THE LAWS OF DECAYING POWER
THE UNEXPECTED TRANSFIGURATION
MINISTRY OF RANDOM WALKS
THE RICH GET RICHER
THE DREAM OF EFFICIENT LANGUAGE
THE LAW OF DIGITS FIRST
THE SURPRISE OF BEING ROBUST YET FRAGILE
THE MISREADING OF THE BOOK OF SAND
THE MEASURING
THE DATA POOR
THE POURING DATA
THE CHARMS OF THE MANY WRONG PATHS
THE DOORS THAT SHOULD NOT BE OPENED
THE KNOWLEDGE OF DEAD MONKS
THE EMERGENCE OF STRUCTURE
THE EMERGENCE OF DESTRUCTION
THE EMERGENCE OF THINKING
It was the best of times,
THE EMERGENCE OF STORIES
THE MECHANICS STATISTICAL
THE THRESHOLDS OF PERCOLATION
THE NETWORKS COMPLEX
THE ELEMENTS OF WEBS
THE BIRDS OF A FEATHER
THE BIRDS OF NOT A FEATHER
THE ARK OF THE RANDOM NETWORK
THE STRANGENESS OF FRIENDS
THE SMALLNESS OF WORLDS
THE THEORY OF SIX DEGREES
THE NETWORKS FREE OF SCALE
CLASSIFIED
PHOENIX
THE CONTAGION
THE CONFUSION OF CONTAGIONS
THE DISEASE OF RANDOM MIXING
THE UNENDING PANDEMICS
THE THRESHOLDS OF THE MIND
THE SOCIAL WILD
THE SOCIAL CONTAGION
THE RATIO OF GAIN
THE MECHANICS OF FAME
THE TALK OF FAME
THE STORIES CONTAGIOUS
THE JOHN DORY
THE ADJACENT NARRATIVES
THE STORYTELLERS
THE FATED ASTRAL PATH
THE PATHS OF UNIVERSALITY
THE DEPENDENT PATHS
THE TROUSERS OF BIFURCATING REALITIES
THE UNKNOWN MECHANISM
THE LANDSCAPES OF FORKING PATHS
THE NETWORKS OF BLOOD
THE TREES OF REALITY
THE ORDERS OF STREAMS
THE LAWS OF BRANCHING
THE SCALING OF BROKEN HIERARCHIES
THE BRAID OR THE BRANCH
THE LAW OF OPTIMAL FORKS
THE CHURCH OF QUARTEROLOGY
THE TRUTHICIDE
THE FAMILY OF PANCAKE COWS
THE FAMILY OF SPHERICAL COWS
THE IMPROBABLE PACHYDERM
THE ILL-SCALING OF ALLOMETRIC BODIES
THE SUPPLY MOST FAIR
THE ORDERS SUPPLY TO THE MOST
THE DARK ART OF GENERATING FUNCTIONS
THE RANDOM SUM OF RANDOMNESS
THE TWO-SIDED WORLDS
THE STORIES AND THE TROPES
THE TRUTH OF DIFFUSION
THE UNVEILING OF HIDDEN POWERS
THE SCRYING FOR THE SHAPES OF THINGS
THE LIMITS OF ORGANIZATIONS
THE OVERCOMING OF CATASTROPHE
THE TROUBLES ILL-FORMED
THE NETWORKS OF MANY ILKS
THE CHAOS OF FETTERED CONTAGION
THE END


THE MANY TALES OF MANY TAILS

Outline

Alternative distributions

There are other 'heavy-tailed' distributions:

1. The Log-normal distribution

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

Alternative distributions

There are other 'heavy-tailed' distributions:

1. The Log-normal distribution

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

2. Weibull distributions

$$P(x)\mathrm{d}x = \frac{k}{\lambda}\left(\frac{x}{\lambda}\right)^{\mu-1} e^{-(x/\lambda)^{\mu}} \mathrm{d}x$$

CCDF = stretched exponential.

Alternative distributions

There are other 'heavy-tailed' distributions:

1. The Log-normal distribution

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

2. Weibull distributions

$$P(x)\mathrm{d}x = \frac{k}{\lambda}\left(\frac{x}{\lambda}\right)^{\mu-1} e^{-(x/\lambda)^{\mu}}\mathrm{d}x$$

   CCDF = stretched exponential.

3. Also: Gamma distribution, Erlang distribution, and more.

lognormals

The lognormal distribution:

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma}\text{exp}\left(-\frac{(\text{ln}x-\mu)^2}{2\sigma^2}\right)$$

- $\ln x$ is distributed according to a normal distribution with mean μ and variance σ.
- Appears in economics and biology where growth increments are distributed normally.

lognormals

- Standard form reveals the mean μ and variance σ^2 of the underlying normal distribution:

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

lognormals

- Standard form reveals the mean μ and variance σ^2 of the underlying normal distribution:

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma}\text{exp}\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

- For lognormals:

$$\mu_{\text{lognormal}} = e^{\mu+\frac{1}{2}\sigma^2}, \qquad \text{median}_{\text{lognormal}} = e^{\mu},$$

$$\sigma_{\text{lognormal}} = (e^{\sigma^2}-1)e^{2\mu+\sigma^2}, \qquad \text{mode}_{\text{lognormal}} = e^{\mu-\sigma^2}.$$

lognormals

- Standard form reveals the mean μ and variance σ^2 of the underlying normal distribution:

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

- For lognormals:

$$\mu_{\text{lognormal}} = e^{\mu + \frac{1}{2}\sigma^2}, \qquad \text{median}_{\text{lognormal}} = e^{\mu},$$

$$\sigma_{\text{lognormal}} = (e^{\sigma^2} - 1)e^{2\mu + \sigma^2}, \qquad \text{mode}_{\text{lognormal}} = e^{\mu - \sigma^2}.$$

- All moments of lognormals are finite.

Derivation from a normal distribution

Take Y as distributed normally:

Derivation from a normal distribution

Take Y as distributed normally:

$$P(y)\mathrm{d}y = \frac{1}{\sqrt{2\pi}\sigma}\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)\mathrm{d}y$$

Derivation from a normal distribution

Take Y as distributed normally:

$$P(y)\mathrm{d}y = \frac{1}{\sqrt{2\pi}\sigma}\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)\mathrm{d}y$$

Set $Y = \ln X$:

Derivation from a normal distribution

Take Y as distributed normally:

$$P(y)\mathrm{d}y = \frac{1}{\sqrt{2\pi}\sigma}\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)\mathrm{d}y$$

Set $Y = \ln X$:

- Transform according to $P(x)\mathrm{d}x = P(y)\mathrm{d}y$:

Derivation from a normal distribution

Take Y as distributed normally:

$$P(y)\mathrm{d}y = \frac{1}{\sqrt{2\pi}\sigma}\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)\mathrm{d}y$$

Set $Y = \ln X$:

- Transform according to $P(x)\mathrm{d}x = P(y)\mathrm{d}y$:
- $$\frac{\mathrm{d}y}{\mathrm{d}x} = 1/x \Rightarrow \mathrm{d}y = \mathrm{d}x/x$$

Derivation from a normal distribution

Take Y as distributed normally:

- $$P(y)\mathrm{d}y = \frac{1}{\sqrt{2\pi}\sigma}\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)\mathrm{d}y$$

Set $Y = \ln X$:

- Transform according to $P(x)\mathrm{d}x = P(y)\mathrm{d}y$:
- $$\frac{\mathrm{d}y}{\mathrm{d}x} = 1/x \Rightarrow \mathrm{d}y = \mathrm{d}x/x$$
- $$\Rightarrow P(x)\mathrm{d}x = \frac{1}{x\sqrt{2\pi}\sigma}\exp\left(-\frac{(\ln x-\mu)^2}{2\sigma^2}\right)\mathrm{d}x$$

Confusion between lognormals and pure power laws



Near agreement over four orders of magnitude!

- For lognormal (blue), $\mu = 0$ and $\sigma = 10$.
- For power law (red), $\gamma = 1$ and $c = 0.03$.

Confusion

What's happening:

$$\ln P(x) = \ln \left\{ \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2} \right) \right\}$$

Confusion

What's happening:

$$\ln P(x) = \ln \left\{ \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2} \right) \right\}$$

$$= -\ln x - \ln\sqrt{2\pi}\sigma - \frac{(\ln x - \mu)^2}{2\sigma^2}$$

Confusion

What's happening:

$$\ln P(x) = \ln \left\{ \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2} \right) \right\}$$

$$= -\ln x - \ln\sqrt{2\pi}\sigma - \frac{(\ln x - \mu)^2}{2\sigma^2}$$

$$= -\frac{1}{2\sigma^2}(\ln x)^2 + \left(\frac{\mu}{\sigma^2} - 1 \right) \ln x - \ln\sqrt{2\pi}\sigma - \frac{\mu^2}{2\sigma^2}.$$

Confusion

What's happening:

$$\ln P(x) = \ln\left\{\frac{1}{x\sqrt{2\pi}\sigma}\exp\left(-\frac{(\ln x-\mu)^2}{2\sigma^2}\right)\right\}$$

$$= -\ln x - \ln\sqrt{2\pi}\sigma - \frac{(\ln x-\mu)^2}{2\sigma^2}$$

$$= -\frac{1}{2\sigma^2}(\ln x)^2 + \left(\frac{\mu}{\sigma^2}-1\right)\ln x - \ln\sqrt{2\pi}\sigma - \frac{\mu^2}{2\sigma^2}.$$

If the first term is relatively small,

Confusion

What's happening:

$$\ln P(x) = \ln\left\{\frac{1}{x\sqrt{2\pi}\sigma}\exp\left(-\frac{(\ln x-\mu)^2}{2\sigma^2}\right)\right\}$$

$$= -\ln x - \ln\sqrt{2\pi}\sigma - \frac{(\ln x-\mu)^2}{2\sigma^2}$$

$$= -\frac{1}{2\sigma^2}(\ln x)^2 + \left(\frac{\mu}{\sigma^2}-1\right)\ln x - \ln\sqrt{2\pi}\sigma - \frac{\mu^2}{2\sigma^2}.$$

If the first term is relatively small,

$$\boxed{\ln P(x) \sim -\left(1-\frac{\mu}{\sigma^2}\right)\ln x + \text{const.}}$$

Confusion

What's happening:

$$\ln P(x) = \ln\left\{\frac{1}{x\sqrt{2\pi}\sigma}\exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)\right\}$$

$$= -\ln x - \ln\sqrt{2\pi}\sigma - \frac{(\ln x - \mu)^2}{2\sigma^2}$$

$$= -\frac{1}{2\sigma^2}(\ln x)^2 + \left(\frac{\mu}{\sigma^2} - 1\right)\ln x - \ln\sqrt{2\pi}\sigma - \frac{\mu^2}{2\sigma^2}.$$

If the first term is relatively small,

$$\boxed{\ln P(x) \sim -\left(1 - \frac{\mu}{\sigma^2}\right)\ln x + \text{const.}} \Rightarrow \boxed{\gamma = 1 - \frac{\mu}{\sigma^2}}$$

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.
- If $\sigma^2 \gg 1$ and μ,

$$\boxed{\ln P(x) \sim -\ln x + \text{const.}}$$

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.
- If $\sigma^2 \gg 1$ and μ,

$$\boxed{\ln P(x) \sim -\ln x + \text{const.}}$$

- Expect -1 scaling to hold until $(\ln x)^2$ term becomes significant compared to $(\ln x)$:

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.
- If $\sigma^2 \gg 1$ and μ,

$$\boxed{\ln P(x) \sim -\ln x + \text{const.}}$$

- Expect -1 scaling to hold until $(\ln x)^2$ term becomes significant compared to $(\ln x)$:

$$-\frac{1}{2\sigma^2}(\ln x)^2 \simeq 0.05\left(\frac{\mu}{\sigma^2} - 1\right)\ln x$$

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.
- If $\sigma^2 \gg 1$ and μ,

$$\boxed{\ln P(x) \sim -\ln x + \text{const.}}$$

- Expect -1 scaling to hold until $(\ln x)^2$ term becomes significant compared to $(\ln x)$:

$$-\frac{1}{2\sigma^2}(\ln x)^2 \simeq 0.05\left(\frac{\mu}{\sigma^2} - 1\right)\ln x$$

$$\Rightarrow \log_{10} x \lesssim 0.05 \times 2(\sigma^2 - \mu)\log_{10} e$$

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.
- If $\sigma^2 \gg 1$ and μ,

$$\boxed{\ln P(x) \sim -\ln x + \text{const.}}$$

- Expect -1 scaling to hold until $(\ln x)^2$ term becomes significant compared to $(\ln x)$:

$$-\frac{1}{2\sigma^2}(\ln x)^2 \simeq 0.05\left(\frac{\mu}{\sigma^2} - 1\right)\ln x$$

$$\Rightarrow \log_{10} x \lesssim 0.05 \times 2(\sigma^2 - \mu)\log_{10} e \simeq 0.05(\sigma^2 - \mu)$$

Confusion

- If $\mu < 0$, $\gamma > 1$ which is totally cool.
- If $\mu > 0$, $\gamma < 1$, not so much.
- If $\sigma^2 \gg 1$ and μ,

$$\boxed{\ln P(x) \sim -\ln x + \text{const.}}$$

- Expect -1 scaling to hold until $(\ln x)^2$ term becomes significant compared to $(\ln x)$:

$$-\frac{1}{2\sigma^2}(\ln x)^2 \simeq 0.05\left(\frac{\mu}{\sigma^2} - 1\right)\ln x$$

$$\Rightarrow \log_{10} x \lesssim 0.05 \times 2(\sigma^2 - \mu)\log_{10} e \simeq 0.05(\sigma^2 - \mu)$$

- $\Rightarrow$ If you find a -1 exponent,
  you may have a lognormal distribution...

Outline

Generating lognormals:

Random multiplicative growth:

$$x_{n+1} = r x_n$$

where $r > 0$ is a random growth variable

Generating lognormals:

Random multiplicative growth:

- $$x_{n+1} = r x_n$$

  where $r > 0$ is a random growth variable
- (Shrinkage is allowed)

Generating lognormals:

Random multiplicative growth:

- $$x_{n+1} = r x_n$$

  where $r > 0$ is a random growth variable
- (Shrinkage is allowed)
- In log space, growth is by addition:

$$\ln x_{n+1} = \ln r + \ln x_n$$

Generating lognormals:

Random multiplicative growth:

- $$x_{n+1} = r x_n$$

  where $r > 0$ is a random growth variable
- (Shrinkage is allowed)
- In log space, growth is by addition:

  $$\ln x_{n+1} = \ln r + \ln x_n$$

- $\Rightarrow \ln x_n$ is normally distributed

Generating lognormals:

Random multiplicative growth:

- $$x_{n+1} = r x_n$$
  where $r > 0$ is a random growth variable
- (Shrinkage is allowed)
- In log space, growth is by addition:
  $$\ln x_{n+1} = \ln r + \ln x_n$$
- $\Rightarrow \ln x_n$ is normally distributed
- $\Rightarrow x_n$ is lognormally distributed

Lognormals or power laws?

- Gibrat [2] (1931) uses preceding argument to explain lognormal distribution of firm sizes ($\gamma \simeq 1$).

Lognormals or power laws?

- Gibrat [2] (1931) uses preceding argument to explain lognormal distribution of firm sizes ($\gamma \simeq 1$).
- But Robert Axtell [1] (2001) shows a power law fits the data very well with $\gamma = 2$, not $\gamma = 1$ (!)

Lognormals or power laws?

- Gibrat [2] (1931) uses preceding argument to explain lognormal distribution of firm sizes ($\gamma \simeq 1$).
- But Robert Axtell [1] (2001) shows a power law fits the data very well with $\gamma = 2$, not $\gamma = 1$ (!)
- Problem of data censusing (missing small firms).

Lognormals or power laws?

- Gibrat [2] (1931) uses preceding argument to explain lognormal distribution of firm sizes ($\gamma \simeq 1$).
- But Robert Axtell [1] (2001) shows a power law fits the data very well with $\gamma = 2$, not $\gamma = 1$ (!)
- Problem of data censusing (missing small firms).



Freq $\propto$ (size)$^{-\gamma}$

$\gamma \simeq 2$

Lognormals or power laws?

- Gibrat [2] (1931) uses preceding argument to explain lognormal distribution of firm sizes ($\gamma \simeq 1$).
- But Robert Axtell [1] (2001) shows a power law fits the data very well with $\gamma = 2$, not $\gamma = 1$ (!)
- Problem of data censusing (missing small firms).



Freq $\propto$ (size)$^{-\gamma}$
$\gamma \simeq 2$

- One piece in Gibrat's model seems okay empirically: Growth rate r appears to be independent of firm size. [1].

An explanation

- Axtel cites Malcai et al.'s (1999) argument [5] for why power laws appear with exponent $\gamma \simeq 2$

An explanation

- Axtel cites Malcai et al.'s (1999) argument [5] for why power laws appear with exponent $\gamma \simeq 2$
- The set up: N entities with size $x_i(t)$

An explanation

- Axtel cites Malcai et al.'s (1999) argument [5] for why power laws appear with exponent $\gamma \simeq 2$
- The set up: N entities with size $x_i(t)$
- Generally:

$$x_i(t+1) = r x_i(t)$$

where r is drawn from some happy distribution

An explanation

- Axtel cites Malcai et al.'s (1999) argument [5] for why power laws appear with exponent $\gamma \simeq 2$
- The set up: N entities with size $x_i(t)$
- Generally:

$$x_i(t+1) = r x_i(t)$$

where r is drawn from some happy distribution
- Same as for lognormal but one extra piece.

An explanation

- Axtel cites Malcai et al.'s (1999) argument [5] for why power laws appear with exponent $\gamma \simeq 2$
- The set up: N entities with size $x_i(t)$
- Generally:

$$x_i(t+1) = r x_i(t)$$

  where r is drawn from some happy distribution
- Same as for lognormal but one extra piece.
- Each x_i cannot drop too low with respect to the other sizes:

$$x_i(t+1) = \max(r x_i(t), c \langle x_i \rangle)$$

Some math later...

Insert assignment question

Some math later...

Insert assignment question

Find $P(x) \sim x^{-\gamma}$

Some math later...

Insert assignment question

$$\text{Find} \quad P(x) \sim x^{-\gamma}$$

- where γ is implicitly given by

$$N = \frac{(\gamma - 2)}{(\gamma - 1)} \left[\frac{(c/N)^{\gamma-1} - 1}{(c/N)^{\gamma-1} - (c/N)} \right]$$

N = total number of firms.

Some math later...

Insert assignment question

$$\text{Find} \quad P(x) \sim x^{-\gamma}$$

- where γ is implicitly given by

$$N = \frac{(\gamma-2)}{(\gamma-1)}\left[\frac{(c/N)^{\gamma-1}-1}{(c/N)^{\gamma-1}-(c/N)}\right]$$

N = total number of firms.

- Now, if $c/N \ll 1$ and $\gamma > 2$ $\quad N = \frac{(\gamma-2)}{(\gamma-1)}\left[\frac{-1}{-(c/N)}\right]$

Some math later...

Insert assignment question

Find $P(x) \sim x^{-\gamma}$

where γ is implicitly given by

$$N = \frac{(\gamma-2)}{(\gamma-1)} \left[\frac{(c/N)^{\gamma-1} - 1}{(c/N)^{\gamma-1} - (c/N)} \right]$$

N = total number of firms.

Now, if $c/N \ll 1$ and $\gamma > 2$ $N = \frac{(\gamma-2)}{(\gamma-1)} \left[\frac{-1}{-(c/N)} \right]$

Which gives $\gamma \sim 1 + \frac{1}{1-c}$

Some math later...

Insert assignment question

- Find $P(x) \sim x^{-\gamma}$

- where γ is implicitly given by

$$N = \frac{(\gamma-2)}{(\gamma-1)}\left[\frac{(c/N)^{\gamma-1}-1}{(c/N)^{\gamma-1}-(c/N)}\right]$$

N = total number of firms.

- Now, if $c/N \ll 1$ and $\gamma > 2$ $\quad N = \frac{(\gamma-2)}{(\gamma-1)}\left[\frac{-1}{-(c/N)}\right]$

- Which gives $\gamma \sim 1 + \frac{1}{1-c}$

- Groovy... c small $\Rightarrow \gamma \simeq 2$

Outline

The second tweak

Ages of firms/people/... may not be the same

The second tweak

Ages of firms/people/... may not be the same

- Allow the number of updates for each size x_i to vary

The second tweak

Ages of firms/people/... may not be the same

- Allow the number of updates for each size x_i to vary
- Example: $P(t)\mathrm{d}t = ae^{-at}\mathrm{d}t$ where t = age.

The second tweak

Ages of firms/people/... may not be the same

- Allow the number of updates for each size x_i to vary
- Example: $P(t)\mathrm{d}t = ae^{-at}\mathrm{d}t$ where t = age.
- Back to no bottom limit: each x_i follows a lognormal

The second tweak

Ages of firms/people/... may not be the same

- Allow the number of updates for each size x_i to vary
- Example: $P(t)\mathrm{d}t = ae^{-at}\mathrm{d}t$ where t = age.
- Back to no bottom limit: each x_i follows a lognormal
- Sizes are distributed as [6]

$$P(x) = \int_{t=0}^{\infty} ae^{-at} \frac{1}{x\sqrt{2\pi t}} \exp\left(-\frac{(\ln x - \mu)^2}{2t}\right) \mathrm{d}t$$

(Assume for this example that $\sigma \sim t$ and $\mu = \ln m$)

The second tweak

Ages of firms/people/... may not be the same

- Allow the number of updates for each size x_i to vary
- Example: $P(t)\mathrm{d}t = ae^{-at}\mathrm{d}t$ where t = age.
- Back to no bottom limit: each x_i follows a lognormal
- Sizes are distributed as [6]

$$P(x) = \int_{t=0}^{\infty} ae^{-at} \frac{1}{x\sqrt{2\pi t}} \exp\left(-\frac{(\ln x - \mu)^2}{2t}\right) \mathrm{d}t$$

(Assume for this example that $\sigma \sim t$ and $\mu = \ln m$)

- Now averaging different lognormal distributions.

Averaging lognormals

$$P(x) = \int_{t=0}^{\infty} a e^{-at} \frac{1}{x\sqrt{2\pi t}} \exp\left(-\frac{(\ln\frac{x}{m})^2}{2t}\right) \mathrm{d}t$$

Averaging lognormals

- $$P(x) = \int_{t=0}^{\infty} a e^{-at} \frac{1}{x\sqrt{2\pi t}} \mathrm{exp}\left(- \frac{(\ln \frac{x}{m})^2}{2t} \right) \mathrm{d}t$$

- Insert fabulous calculation (team is spared).

Averaging lognormals

$$P(x) = \int_{t=0}^{\infty} a e^{-at} \frac{1}{x\sqrt{2\pi t}} \exp\left(-\frac{(\ln \frac{x}{m})^2}{2t} \right) \mathrm{d}t$$

- Insert fabulous calculation (team is spared).
- Some enjoyable suffering leads to:

$$P(x) \propto x^{-1} e^{-\sqrt{2\lambda (\ln \frac{x}{m})^2}}$$

The second tweak

$$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln\frac{x}{m})^2}}$$

The second tweak

$$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln\frac{x}{m})^2}}$$

- Depends on sign of $\ln\frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.

The second tweak

- $$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln\frac{x}{m})^2}}$$
- Depends on sign of $\ln\frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.
- $$P(x) \propto \begin{cases} x^{-1+\sqrt{2\lambda}} & \text{if } \frac{x}{m} < 1 \\ x^{-1-\sqrt{2\lambda}} & \text{if } \frac{x}{m} > 1 \end{cases}$$

The second tweak

- $$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln\frac{x}{m})^2}}$$
- Depends on sign of $\ln\frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.
- $$P(x) \propto \begin{cases} x^{-1+\sqrt{2\lambda}} & \text{if } \frac{x}{m} < 1 \\ x^{-1-\sqrt{2\lambda}} & \text{if } \frac{x}{m} > 1 \end{cases}$$
- 'Break' in scaling (not uncommon)

The second tweak

- $$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln\frac{x}{m})^2}}$$
- Depends on sign of $\ln\frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.
- $$P(x) \propto \begin{cases} x^{-1+\sqrt{2\lambda}} & \text{if } \frac{x}{m} < 1 \\ x^{-1-\sqrt{2\lambda}} & \text{if } \frac{x}{m} > 1 \end{cases}$$
- 'Break' in scaling (not uncommon)
- Double-Pareto distribution

The second tweak

- $$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln \frac{x}{m})^2}}$$
- Depends on sign of $\ln \frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.
- $$P(x) \propto \begin{cases} x^{-1+\sqrt{2\lambda}} & \text{if } \frac{x}{m} < 1 \\ x^{-1-\sqrt{2\lambda}} & \text{if } \frac{x}{m} > 1 \end{cases}$$
- 'Break' in scaling (not uncommon)
- Double-Pareto distribution
- First noticed by Montroll and Shlesinger [7, 8]

The second tweak

- $$P(x) \propto x^{-1} e^{-\sqrt{2\lambda (\ln \frac{x}{m})^2}}$$
- Depends on sign of $\ln \frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.
- $$P(x) \propto \begin{cases} x^{-1+\sqrt{2\lambda}} & \text{if } \frac{x}{m} < 1 \\ x^{-1-\sqrt{2\lambda}} & \text{if } \frac{x}{m} > 1 \end{cases}$$
- 'Break' in scaling (not uncommon)
- Double-Pareto distribution
- First noticed by Montroll and Shlesinger [7, 8]
- Later: Huberman and Adamic [3, 4]: Number of pages per website

Summary of these exciting developments:

- Lognormals and power laws can be awfully similar

Summary of these exciting developments:

- Lognormals and power laws can be awfully similar
- Random Multiplicative Growth leads to lognormal distributions

Summary of these exciting developments:

- Lognormals and power laws can be awfully similar
- Random Multiplicative Growth leads to lognormal distributions
- Enforcing a minimum size leads to a power law tail

Summary of these exciting developments:

- Lognormals and power laws can be awfully similar
- Random Multiplicative Growth leads to lognormal distributions
- Enforcing a minimum size leads to a power law tail
- With no minimum size but a distribution of lifetimes, the double Pareto distribution appears

Summary of these exciting developments:

- Lognormals and power laws can be awfully similar
- Random Multiplicative Growth leads to lognormal distributions
- Enforcing a minimum size leads to a power law tail
- With no minimum size but a distribution of lifetimes, the double Pareto distribution appears
- Take-home message: Be careful out there...

References I

[1] R. Axtell.
Zipf distribution of U.S. firm sizes.
Science, 293(5536):1818–1820, 2001. pdf

[2] R. Gibrat.
Les inégalités économiques.
Librairie du Recueil Sirey, Paris, France, 1931.

[3] B. A. Huberman and L. A. Adamic.
Evolutionary dynamics of the World Wide Web.
Technical report, Xerox Palo Alto Research Center, 1999.

[4] B. A. Huberman and L. A. Adamic.
The nature of markets in the World Wide Web.
Quarterly Journal of Economic Commerce, 1:5–12, 2000.

References II

[5] O. Malcai, O. Biham, and S. Solomon.
Power-law distributions and lévy-stable intermittent fluctuations in stochastic systems of many autocatalytic elements.
Phys. Rev. E, 60(2):1299–1303, 1999. pdf

[6] M. Mitzenmacher.
A brief history of generative models for power law and lognormal distributions.
Internet Mathematics, 1:226–251, 2003. pdf

[7] E. W. Montroll and M. W. Shlesinger.
On $1/f$ noise and other distributions with long tails.
Proc. Natl. Acad. Sci., 79:3380–3383, 1982. pdf

References III

[8] E. W. Montroll and M. W. Shlesinger.
Maximum entropy formalism, fractals, scaling phenomena, and $1/f$ noise: a tale of tails.
J. Stat. Phys., 32:209–230, 1983.