

Mixed, correlated random networks

Last updated: 2023/08/22, 11:48:25 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D
CSYS/MATH 6701, 6713, & a pretend number,
2023–2024 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
Santa Fe Institute | University of Vermont



These slides are brought to you by:

Sealie & Lambie
Productions

The PoCverse
Mixed, correlated
random networks
2 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



These slides are also brought to you by:

Special Guest Executive Producer



 On Instagram at [pratchett_the_cat](https://www.instagram.com/pratchett_the_cat) 

The PoCverse
Mixed, correlated
random networks
3 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Outline

Directed random networks

Mixed random networks

Definition

Correlations

Mixed Random Network Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

The PoCverse
Mixed, correlated
random networks
4 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Random directed networks:



So far, we've largely studied networks with undirected, unweighted edges.

The PoCverse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Random directed networks:



So far, we've largely studied networks with undirected, unweighted edges.



Now consider directed, unweighted edges.



The PoCVerse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Random directed networks:



So far, we've largely studied networks with undirected, unweighted edges.



Now consider directed, unweighted edges.



Nodes have k_i and k_o incoming and outgoing edges, otherwise random.

The PoCVerse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Random directed networks:



So far, we've largely studied networks with undirected, unweighted edges.



Now consider directed, unweighted edges.



Nodes have k_i and k_o incoming and outgoing edges, otherwise random.



Network defined by joint in- and out-degree distribution: P_{k_i, k_o}

The PoCVerse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Random directed networks:



So far, we've largely studied networks with undirected, unweighted edges.

Now consider directed, unweighted edges.



Nodes have k_i and k_o incoming and outgoing edges, otherwise random.

Network defined by joint in- and out-degree distribution: P_{k_i, k_o}

Normalization: $\sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} P_{k_i, k_o} = 1$

The PoCVerse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell


References




Random directed networks:





 So far, we've largely studied networks with undirected, unweighted edges.


 Now consider directed, unweighted edges.



 Nodes have k_i and k_o incoming and outgoing edges, otherwise random.

 Network defined by joint in- and out-degree distribution: P_{k_i, k_o}

 Normalization: $\sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} P_{k_i, k_o} = 1$

 Marginal in-degree and out-degree distributions:

$$P_{k_i} = \sum_{k_o=0}^{\infty} P_{k_i, k_o} \quad \text{and} \quad P_{k_o} = \sum_{k_i=0}^{\infty} P_{k_i, k_o}$$

The PoCVerse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Random directed networks:



So far, we've largely studied networks with undirected, unweighted edges.

Now consider directed, unweighted edges.



Nodes have k_i and k_o incoming and outgoing edges, otherwise random.

Network defined by joint in- and out-degree distribution: P_{k_i, k_o}

Normalization: $\sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} P_{k_i, k_o} = 1$

Marginal in-degree and out-degree distributions:

$$P_{k_i} = \sum_{k_o=0}^{\infty} P_{k_i, k_o} \quad \text{and} \quad P_{k_o} = \sum_{k_i=0}^{\infty} P_{k_i, k_o}$$

Required balance:

$$\langle k_i \rangle = \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_i P_{k_i, k_o} = \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_o P_{k_i, k_o} = \langle k_o \rangle$$

The PoCVerse
Mixed, correlated
random networks
7 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion


Spreading condition
Full generalization
Triggering probabilities

Nutshell


References





Directed network structure:





From Boguñá and Serano. [1]

 GWCC = Giant Weakly Connected Component (directions removed);

 GIN = Giant In-Component;

 GOUT = Giant Out-Component;

 GSCC = Giant Strongly Connected Component;

 DC = Disconnected Components (finite).

The PoCverse
Mixed, correlated
random networks
8 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion


Spreading condition
Full generalization
Triggering probabilities

Nutshell


References





Directed network structure:





From Boguñá and Serano. [1]


 GWCC = Giant Weakly Connected Component (directions removed);

 GIN = Giant In-Component;

 GOUT = Giant Out-Component;

 GSCC = Giant Strongly Connected Component;

 DC = Disconnected Components (finite).

 When moving through a family of increasingly connected directed random networks, GWCC usually appears before GIN, GOUT, and GSCC which tend to appear together. [4, 1]

The PoCverse
Mixed, correlated
random networks
8 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Outline

Directed random networks

Mixed random networks

Definition

Correlations

Mixed Random Network Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

The PoCverse
Mixed, correlated
random networks
9 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Observation:



Directed and undirected random networks are separate families ...

The PoCverse
Mixed, correlated
random networks
10 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Observation:



Directed and undirected random networks are separate families ...



...and analyses are also disjoint.

The PoCverse
Mixed, correlated
random networks
10 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Observation:

- Directed and undirected random networks are separate families ...
- ...and analyses are also disjoint.
- Need to examine a larger family of random networks with mixed directed and undirected edges.

The PoCverse
Mixed, correlated
random networks
10 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References




Observation:

- Directed and undirected random networks are separate families ...
- ...and analyses are also disjoint.
- Need to examine a larger family of random networks with mixed directed and undirected edges.

Consider nodes with three types of edges:

- k_u undirected edges,
- k_i incoming directed edges,
- k_o outgoing directed edges.



The PoCverse
Mixed, correlated
random networks
10 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Observation:


- Directed and undirected random networks are separate families ...
- ...and analyses are also disjoint.
- Need to examine a larger family of random networks with mixed directed and undirected edges.

Consider nodes with three types of edges:

- k_u undirected edges,
- k_i incoming directed edges,
- k_o outgoing directed edges.

Define a node by generalized degree:

$$\vec{k} = [k_u \ k_i \ k_o]^T.$$



The PoCverse
Mixed, correlated
random networks
10 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Joint degree distribution:

$$P_{\vec{k}} \text{ where } \vec{k} = [k_u \ k_i \ k_o]^T.$$

The PoCverse
Mixed, correlated
random networks
11 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell

References




Joint degree distribution:

$$P_{\vec{k}} \text{ where } \vec{k} = [k_u \ k_i \ k_o]^T.$$


 As for directed networks, require in- and out-degree averages to match up:

$$\langle k_i \rangle = \sum_{k_u=0}^{\infty} \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_i P_{\vec{k}} = \sum_{k_u=0}^{\infty} \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_o P_{\vec{k}} = \langle k_o \rangle$$




 Joint degree distribution:

$$P_{\vec{k}} \text{ where } \vec{k} = [k_u \ k_i \ k_o]^T.$$

 As for directed networks, require in- and out-degree averages to match up:

$$\langle k_i \rangle = \sum_{k_u=0}^{\infty} \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_i P_{\vec{k}} = \sum_{k_u=0}^{\infty} \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_o P_{\vec{k}} = \langle k_o \rangle$$

 Otherwise, no other restrictions and connections are random.

The PoCverse
Mixed, correlated
random networks
11 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network


Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell

References





 Joint degree distribution:

$$P_{\vec{k}} \text{ where } \vec{k} = [k_u \ k_i \ k_o]^T.$$

 As for directed networks, require in- and out-degree averages to match up:

$$\langle k_i \rangle = \sum_{k_u=0}^{\infty} \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_i P_{\vec{k}} = \sum_{k_u=0}^{\infty} \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} k_o P_{\vec{k}} = \langle k_o \rangle$$

 Otherwise, no other restrictions and connections are random.

 Directed and undirected random networks are disjoint subfamilies:

$$\text{Undirected: } P_{\vec{k}} = P_{k_u} \delta_{k_i,0} \delta_{k_o,0},$$

$$\text{Directed: } P_{\vec{k}} = \delta_{k_u,0} P_{k_i, k_o}.$$



Outline

Directed random networks

Mixed random networks

Definition

Correlations

Mixed Random Network Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

The PoCverse
Mixed, correlated
random networks
12 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References



Correlations:



Now add correlations (two point or Markovian)

The PoCverse
Mixed, correlated
random networks
13 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network
Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}')$ = probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.

The PoCSverse
Mixed, correlated
random networks
13 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}')$ = probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.
2. $P^{(i)}(\vec{k} | \vec{k}')$ = probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an in-directed edge relative to the destination node.

The PoCverse
Mixed, correlated
random networks
13 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network
Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}')$ = probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.
2. $P^{(i)}(\vec{k} | \vec{k}')$ = probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an in-directed edge relative to the destination node.
3. $P^{(o)}(\vec{k} | \vec{k}')$ = probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an out-directed edge relative to the destination node.



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}') =$ probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.
2. $P^{(i)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an in-directed edge relative to the destination node.
3. $P^{(o)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an out-directed edge relative to the destination node.



Now require more refined (detailed) balance.



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}') =$ probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.
2. $P^{(i)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an in-directed edge relative to the destination node.
3. $P^{(o)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an out-directed edge relative to the destination node.



Now require more refined (detailed) balance.



Conditional probabilities cannot be arbitrary.



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}') =$ probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.
2. $P^{(i)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an in-directed edge relative to the destination node.
3. $P^{(o)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an out-directed edge relative to the destination node.



Now require more refined (detailed) balance.



Conditional probabilities cannot be arbitrary.

1. $P^{(u)}(\vec{k} | \vec{k}')$ must be related to $P^{(u)}(\vec{k}' | \vec{k})$.



Correlations:



Now add correlations (two point or Markovian) □:

1. $P^{(u)}(\vec{k} | \vec{k}') =$ probability that an undirected edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node.
2. $P^{(i)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an in-directed edge relative to the destination node.
3. $P^{(o)}(\vec{k} | \vec{k}') =$ probability that an edge leaving a degree \vec{k}' nodes arrives at a degree \vec{k} node is an out-directed edge relative to the destination node.



Now require more refined (detailed) balance.





Conditional probabilities cannot be arbitrary.

1. $P^{(u)}(\vec{k} | \vec{k}')$ must be related to $P^{(u)}(\vec{k}' | \vec{k})$.
2. $P^{(o)}(\vec{k} | \vec{k}')$ and $P^{(i)}(\vec{k} | \vec{k}')$ must be connected.



Correlations—Undirected edge balance:

 Randomly choose an edge, and randomly choose one end.



The PoCverse
Mixed, correlated
random networks
14 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network
Contagion

Spreading condition

Full generalization

Triggering probabilities


Nutshell

References



Correlations—Undirected edge balance:

- ☄ Randomly choose an edge, and randomly choose one end.
- ☄ Say we find a degree \vec{k} node at this end, and a degree \vec{k}' node at the other end.



The PoCverse
Mixed, correlated
random networks
14 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization

Triggering probabilities


Nutshell

References



Correlations—Undirected edge balance:

- ☰ Randomly choose an edge, and randomly choose one end.
- ☰ Say we find a degree \vec{k} node at this end, and a degree \vec{k}' node at the other end.
- ☰ Define probability this happens as $P^{(u)}(\vec{k}, \vec{k}')$.



The PoCverse
Mixed, correlated
random networks
14 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization

Triggering probabilities


Nutshell

References




Correlations—Undirected edge balance:

- ☰ Randomly choose an edge, and randomly choose one end.
- ☰ Say we find a degree \vec{k} node at this end, and a degree \vec{k}' node at the other end.
- ☰ Define probability this happens as $P^{(u)}(\vec{k}, \vec{k}')$.
- ☰ Observe we must have $P^{(u)}(\vec{k}, \vec{k}') = P^{(u)}(\vec{k}', \vec{k})$.



Correlations—Undirected edge balance:

- Randomly choose an edge, and randomly choose one end.
- Say we find a degree \vec{k} node at this end, and a degree \vec{k}' node at the other end.
- Define probability this happens as $P^{(u)}(\vec{k}, \vec{k}')$.
- Observe we must have $P^{(u)}(\vec{k}, \vec{k}') = P^{(u)}(\vec{k}', \vec{k})$.



- Conditional probability connection:

$$P^{(u)}(\vec{k}, \vec{k}') = P^{(u)}(\vec{k} | \vec{k}') \frac{k'_u P(\vec{k}')}{\langle k'_u \rangle}$$

$$P^{(u)}(\vec{k}', \vec{k}) = P^{(u)}(\vec{k}' | \vec{k}) \frac{k_u P(\vec{k})}{\langle k_u \rangle}$$



Correlations—Directed edge balance:




The quantities

$$\frac{k_o P(\vec{k})}{\langle k_o \rangle} \text{ and } \frac{k_i P(\vec{k})}{\langle k_i \rangle}$$

give the probabilities that in starting at a random end of a randomly selected edge, we begin at a degree \vec{k} node and then find ourselves travelling:

1. along an outgoing edge, or
2. against the direction of an incoming edge.



The PoCverse
Mixed, correlated
random networks
15 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion


Spreading condition
Full generalization
Triggering probabilities

Nutshell

References




Correlations—Directed edge balance:

 The quantities


$$\frac{k_o P(\vec{k})}{\langle k_o \rangle} \text{ and } \frac{k_i P(\vec{k})}{\langle k_i \rangle}$$

give the probabilities that in starting at a random end of a randomly selected edge, we begin at a degree \vec{k} node and then find ourselves travelling:

1. along an outgoing edge, or
2. against the direction of an incoming edge.

 We therefore have

$$P^{(\text{dir})}(\vec{k}, \vec{k}') = P^{(i)}(\vec{k} | \vec{k}') \frac{k'_o P(\vec{k}')}{\langle k'_o \rangle} = P^{(o)}(\vec{k}' | \vec{k}) \frac{k_i P(\vec{k})}{\langle k_i \rangle}.$$



Correlations—Directed edge balance:



The quantities

$$\frac{k_o P(\vec{k})}{\langle k_o \rangle} \text{ and } \frac{k_i P(\vec{k})}{\langle k_i \rangle}$$

give the probabilities that in starting at a random end of a randomly selected edge, we begin at a degree \vec{k} node and then find ourselves travelling:

1. along an outgoing edge, or
2. against the direction of an incoming edge.




We therefore have

$$P^{(\text{dir})}(\vec{k}, \vec{k}') = P^{(i)}(\vec{k} | \vec{k}') \frac{k'_o P(\vec{k}')}{\langle k'_o \rangle} = P^{(o)}(\vec{k}' | \vec{k}) \frac{k_i P(\vec{k})}{\langle k_i \rangle}.$$



Note that $P^{(\text{dir})}(\vec{k}, \vec{k}')$ and $P^{(\text{dir})}(\vec{k}', \vec{k})$ are in general not related if $\vec{k} \neq \vec{k}'$.



The PoCverse
Mixed, correlated
random networks
15 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Outline

Directed random networks

Mixed random networks

Definition

Correlations

Mixed Random Network Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

The PoCverse
Mixed, correlated
random networks
16 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References



Global spreading condition: [2]

When are cascades possible?:

The PoCverse
Mixed, correlated
random networks
17 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell

References



Global spreading condition: [2]

When are cascades possible?:

 Consider uncorrelated mixed networks first.

The PoCverse
Mixed, correlated
random networks
17 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition: [2]

When are cascades possible?:

- Consider uncorrelated mixed networks first.
- Recall our first result for undirected random networks, that edge gain ratio must exceed 1:

$$\mathbf{R} = \sum_{k_u=0}^{\infty} \frac{k_u P_{k_u}}{\langle k_u \rangle} \bullet (k_u - 1) \bullet B_{k_u,1} > 1.$$

The PoCverse
Mixed, correlated
random networks
17 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition: [2]

When are cascades possible?:

- Consider uncorrelated mixed networks first.
- Recall our first result for undirected random networks, that edge gain ratio must exceed 1:

$$R = \sum_{k_u=0}^{\infty} \frac{k_u P_{k_u}}{\langle k_u \rangle} \cdot (k_u - 1) \cdot B_{k_u, 1} > 1.$$

- Similar form for purely directed networks:

$$R = \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} \frac{k_i P_{k_i, k_o}}{\langle k_i \rangle} \cdot k_o \cdot B_{k_i, 1} > 1.$$

The PoCverse
Mixed, correlated
random networks
17 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition: [2]

When are cascades possible?:

- Consider uncorrelated mixed networks first.
- Recall our first result for undirected random networks, that edge gain ratio must exceed 1:

$$\mathbf{R} = \sum_{k_u=0}^{\infty} \frac{k_u P_{k_u}}{\langle k_u \rangle} \cdot (k_u - 1) \cdot B_{k_u, 1} > 1.$$

- Similar form for purely directed networks:

$$\mathbf{R} = \sum_{k_i=0}^{\infty} \sum_{k_o=0}^{\infty} \frac{k_i P_{k_i, k_o}}{\langle k_i \rangle} \cdot k_o \cdot B_{k_i, 1} > 1.$$

- Both are composed of (1) probability of connection to a node of a given type; (2) number of newly infected edges if successful; and (3) probability of infection.

The PoCverse
Mixed, correlated
random networks
17 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell

References



Global spreading condition:

Local growth equation:

 Define number of infected edges leading to nodes a distance d away from the original seed as $f(d)$.

The PoCverse
Mixed, correlated
random networks
18 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition:

Local growth equation:

- Define number of infected edges leading to nodes a distance d away from the original seed as $f(d)$.
- Infected edge growth equation:

$$f(d + 1) = \mathbf{R}f(d).$$

The PoCVerse
Mixed, correlated
random networks
18 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell


References




Global spreading condition:

Local growth equation:

 Define number of infected edges leading to nodes a distance d away from the original seed as $f(d)$.

 Infected edge growth equation:

$$f(d + 1) = \mathbf{R}f(d).$$

 Applies for discrete time and continuous time contagion processes.

The PoCverse
Mixed, correlated
random networks
18 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition:

Local growth equation:

- Define number of infected edges leading to nodes a distance d away from the original seed as $f(d)$.
- Infected edge growth equation:

$$f(d + 1) = \mathbf{R}f(d).$$

- Applies for discrete time and continuous time contagion processes.
- Now see $B_{k_u, 1}$ is the probability that an infected edge eventually infects a node.

The PoCverse
Mixed, correlated
random networks
18 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition:

Local growth equation:

- Define number of infected edges leading to nodes a distance d away from the original seed as $f(d)$.
- Infected edge growth equation:

$$f(d + 1) = \mathbf{R}f(d).$$

- Applies for discrete time and continuous time contagion processes.
- Now see $B_{k_u, 1}$ is the probability that an infected edge eventually infects a node.
- Also allows for recovery of nodes (SIR).

The PoCVerse
Mixed, correlated
random networks
18 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell

References



Global spreading condition:

Mixed, uncorrelated random networks:

 Now have two types of edges spreading infection: directed and undirected.

The PoCverse
Mixed, correlated
random networks
19 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell


References



Global spreading condition:

Mixed, uncorrelated random networks:

 Now have two types of edges spreading infection: directed and undirected.

 Gain ratio now more complicated:

1. Infected directed edges can lead to infected directed or undirected edges.
2. Infected undirected edges can lead to infected directed or undirected edges.

The PoCverse
Mixed, correlated
random networks
19 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network

Contagion

Spreading condition
Full generalization
Triggering probabilities


Nutshell


References




Global spreading condition:

Mixed, uncorrelated random networks:

 Now have two types of edges spreading infection: directed and undirected.

 Gain ratio now more complicated:

1. Infected directed edges can lead to infected directed or undirected edges.
2. Infected undirected edges can lead to infected directed or undirected edges.

 Define $f^{(u)}(d)$ and $f^{(o)}(d)$ as the expected number of infected undirected and directed edges leading to nodes a distance d from seed.

The PoCverse
Mixed, correlated
random networks
19 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References





Gain ratio now has a matrix form:

$$\begin{bmatrix} f^{(u)}(d+1) \\ f^{(o)}(d+1) \end{bmatrix} = \mathbf{R} \begin{bmatrix} f^{(u)}(d) \\ f^{(o)}(d) \end{bmatrix}$$

Gain ratio now has a matrix form:

$$\begin{bmatrix} f^{(u)}(d+1) \\ f^{(o)}(d+1) \end{bmatrix} = \mathbf{R} \begin{bmatrix} f^{(u)}(d) \\ f^{(o)}(d) \end{bmatrix}$$

Two separate gain equations:

$$f^{(u)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet (k_u - 1) \bullet B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_u \bullet B_{k_u+k_i,1} f^{(o)}(d) \right]$$

🧩 Gain ratio now has a matrix form:

$$\begin{bmatrix} f^{(u)}(d+1) \\ f^{(o)}(d+1) \end{bmatrix} = \mathbf{R} \begin{bmatrix} f^{(u)}(d) \\ f^{(o)}(d) \end{bmatrix}$$

🧩 Two separate gain equations:

$$f^{(u)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \cdot (k_u - 1) \cdot B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \cdot k_u \cdot B_{k_u+k_i,1} f^{(o)}(d) \right]$$

$$f^{(o)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \cdot k_o B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \cdot k_o \cdot B_{k_u+k_i,1} f^{(o)}(d) \right]$$

Gain ratio now has a matrix form:

$$\begin{bmatrix} f^{(u)}(d+1) \\ f^{(o)}(d+1) \end{bmatrix} = \mathbf{R} \begin{bmatrix} f^{(u)}(d) \\ f^{(o)}(d) \end{bmatrix}$$


Two separate gain equations:

$$f^{(u)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet (k_u - 1) \bullet B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_u \bullet B_{k_u+k_i,1} f^{(o)}(d) \right]$$


$$f^{(o)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet k_o B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_o \bullet B_{k_u+k_i,1} f^{(o)}(d) \right]$$

Gain ratio matrix:

$$\mathbf{R} = \sum_{\bar{k}} \begin{bmatrix} \frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet (k_u - 1) & \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_u \\ \frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet k_o & \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_o \end{bmatrix} \bullet B_{k_u+k_i,1}$$


 Gain ratio now has a matrix form:

$$\begin{bmatrix} f^{(u)}(d+1) \\ f^{(o)}(d+1) \end{bmatrix} = \mathbf{R} \begin{bmatrix} f^{(u)}(d) \\ f^{(o)}(d) \end{bmatrix}$$


 Two separate gain equations:

$$f^{(u)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet (k_u - 1) \bullet B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_u \bullet B_{k_u+k_i,1} f^{(o)}(d) \right]$$


$$f^{(o)}(d+1) = \sum_{\bar{k}} \left[\frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet k_o B_{k_u+k_i,1} f^{(u)}(d) + \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_o \bullet B_{k_u+k_i,1} f^{(o)}(d) \right]$$

 Gain ratio matrix:

$$\mathbf{R} = \sum_{\bar{k}} \begin{bmatrix} \frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet (k_u - 1) & \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_u \\ \frac{k_u P_{\bar{k}}}{\langle k_u \rangle} \bullet k_o & \frac{k_i P_{\bar{k}}}{\langle k_i \rangle} \bullet k_o \end{bmatrix} \bullet B_{k_u+k_i,1}$$

 Spreading condition: max eigenvalue of $\mathbf{R} > 1$.

Global spreading condition:

 Useful change of notation for making results more general: write $P^{(u)}(\vec{k} | *) = \frac{k_u P_{\vec{k}}}{\langle k_u \rangle}$ and $P^{(i)}(\vec{k} | *) = \frac{k_i P_{\vec{k}}}{\langle k_i \rangle}$ where * indicates the starting node's degree is irrelevant (no correlations).

The PoCverse
Mixed, correlated
random networks
21 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition:

- Useful change of notation for making results more general: write $P^{(u)}(\vec{k} | *) = \frac{k_u P_{\vec{k}}}{\langle k_u \rangle}$ and $P^{(i)}(\vec{k} | *) = \frac{k_i P_{\vec{k}}}{\langle k_i \rangle}$ where $*$ indicates the starting node's degree is irrelevant (no correlations).
- Also write $B_{k_u k_i, *}$ to indicate a more general infection probability, but one that does not depend on the edge's origin.

The PoCVerse
Mixed, correlated
random networks
21 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Global spreading condition:

Useful change of notation for making results more general: write $P^{(u)}(\vec{k} | *) = \frac{k_u P_{\vec{k}}}{\langle k_u \rangle}$ and

$P^{(i)}(\vec{k} | *) = \frac{k_i P_{\vec{k}}}{\langle k_i \rangle}$ where $*$ indicates the starting node's degree is irrelevant (no correlations).


Also write $B_{k_u k_i, *}$ to indicate a more general infection probability, but one that does not depend on the edge's origin.

Now have, for the example of mixed, uncorrelated random networks:

$$\mathbf{R} = \sum_{\vec{k}} \begin{bmatrix} P^{(u)}(\vec{k} | *) \bullet (k_u - 1) & P^{(i)}(\vec{k} | *) \bullet k_u \\ P^{(u)}(\vec{k} | *) \bullet k_o & P^{(i)}(\vec{k} | *) \bullet k_o \end{bmatrix} \bullet B_{k_u k_i, *}$$



Summary of contagion conditions for uncorrelated networks:

 I. Undirected, Uncorrelated— $f(d + 1) = \mathbf{f}(d)$:

$$\mathbf{R} = \sum_{k_U} P^{(u)}(k_U | *) \bullet (k_U - 1) \bullet B_{k_U, *}$$

The PoCverse
Mixed, correlated
random networks
22 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition

Full generalization
Triggering probabilities

Nutshell

References



Summary of contagion conditions for uncorrelated networks:

I. Undirected, Uncorrelated— $f(d + 1) = \mathbf{f}(d)$:

$$\mathbf{R} = \sum_{k_u} P^{(u)}(k_u | *) \bullet (k_u - 1) \bullet B_{k_u, *}$$

II. Directed, Uncorrelated— $f(d + 1) = \mathbf{f}(d)$:

$$\mathbf{R} = \sum_{k_i, k_o} P^{(i)}(k_i, k_o | *) \bullet k_o \bullet B_{k_i, *}$$

The PoCverse
Mixed, correlated
random networks
22 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion


Spreading condition
Full generalization
Triggering probabilities

Nutshell


References




Summary of contagion conditions for uncorrelated networks:

 I. Undirected, Uncorrelated— $f(d + 1) = \mathbf{f}(d)$:

$$\mathbf{R} = \sum_{k_u} P^{(u)}(k_u | *) \bullet (k_u - 1) \bullet B_{k_u, *}$$

 II. Directed, Uncorrelated— $f(d + 1) = \mathbf{f}(d)$:

$$\mathbf{R} = \sum_{k_i, k_o} P^{(i)}(k_i, k_o | *) \bullet k_o \bullet B_{k_i, *}$$

 III. Mixed Directed and Undirected, Uncorrelated—

$$\begin{bmatrix} f^{(u)}(d + 1) \\ f^{(o)}(d + 1) \end{bmatrix} = \mathbf{R} \begin{bmatrix} f^{(u)}(d) \\ f^{(o)}(d) \end{bmatrix}$$

$$\mathbf{R} = \sum_{\vec{k}} \begin{bmatrix} P^{(u)}(\vec{k} | *) \bullet (k_u - 1) & P^{(i)}(\vec{k} | *) \bullet k_u \\ P^{(u)}(\vec{k} | *) \bullet k_o & P^{(i)}(\vec{k} | *) \bullet k_o \end{bmatrix} \bullet B_{k_u, k_i, *}$$



Correlated version:



Now have to think of transfer of infection from edges emanating from degree \vec{k}' nodes to edges emanating from degree \vec{k} nodes.

The PoCverse
Mixed, correlated
random networks
23 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion


Spreading condition
Full generalization
Triggering probabilities


Nutshell

References



Correlated version:

 Now have to think of transfer of infection from edges emanating from degree \vec{k}' nodes to edges emanating from degree \vec{k} nodes.

 Replace $P^{(i)}(\vec{k} | *)$ with $P^{(i)}(\vec{k} | \vec{k}')$ and so on.



Correlated version:

- Now have to think of transfer of infection from edges emanating from degree \vec{k}' nodes to edges emanating from degree \vec{k} nodes.
- Replace $P^{(i)}(\vec{k} | *)$ with $P^{(i)}(\vec{k} | \vec{k}')$ and so on.
- Edge types are now more diverse beyond directed and undirected as originating node type matters.



Correlated version:

- Now have to think of transfer of infection from edges emanating from degree \vec{k}' nodes to edges emanating from degree \vec{k} nodes.
- Replace $P^{(i)}(\vec{k} | *)$ with $P^{(i)}(\vec{k} | \vec{k}')$ and so on.
- Edge types are now more diverse beyond directed and undirected as originating node type matters.
- Sums are now over \vec{k}' .



Summary of contagion conditions for correlated networks:



IV. Undirected,

Correlated— $f_{k_u}(d+1) = \sum_{k'_u} R_{k_u k'_u} f_{k'_u}(d)$

$$R_{k_u k'_u} = P^{(u)}(k_u | k'_u) \cdot (k_u - 1) \cdot B_{k_u k'_u}$$

The PoCverse
Mixed, correlated
random networks
24 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Summary of contagion conditions for correlated networks:



IV. Undirected,

Correlated— $f_{k_u}(d+1) = \sum_{k'_u} R_{k_u k'_u} f_{k'_u}(d)$

$$R_{k_u k'_u} = P^{(u)}(k_u | k'_u) \cdot (k_u - 1) \cdot B_{k_u k'_u}$$



V. Directed,

Correlated— $f_{k_i k_o}(d+1) = \sum_{k'_i, k'_o} R_{k_i k_o k'_i k'_o} f_{k'_i k'_o}(d)$

$$R_{k_i k_o k'_i k'_o} = P^{(i)}(k_i, k_o | k'_i, k'_o) \cdot k_o \cdot B_{k_i k_o k'_i k'_o}$$

The PoCverse
Mixed, correlated
random networks
24 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Summary of contagion conditions for correlated networks:



IV. Undirected,

Correlated— $f_{k_u}(d+1) = \sum_{k'_u} R_{k_u k'_u} f_{k'_u}(d)$

$$R_{k_u k'_u} = P^{(u)}(k_u | k'_u) \cdot (k_u - 1) \cdot B_{k_u k'_u}$$



V. Directed,

Correlated— $f_{k_i k_o}(d+1) = \sum_{k'_i, k'_o} R_{k_i k_o k'_i k'_o} f_{k'_i k'_o}(d)$

$$R_{k_i k_o k'_i k'_o} = P^{(i)}(k_i, k_o | k'_i, k'_o) \cdot k_o \cdot B_{k_i k_o k'_i k'_o}$$



VI. Mixed Directed and Undirected, Correlated—

$$\begin{bmatrix} f_{\vec{k}}^{(u)}(d+1) \\ f_{\vec{k}}^{(o)}(d+1) \end{bmatrix} = \sum_{\vec{k}'} \mathbf{R}_{\vec{k} \vec{k}'} \begin{bmatrix} f_{\vec{k}'}^{(u)}(d) \\ f_{\vec{k}'}^{(o)}(d) \end{bmatrix}$$

$$\mathbf{R}_{\vec{k} \vec{k}'} = \begin{bmatrix} P^{(u)}(\vec{k} | \vec{k}') \cdot (k_u - 1) & P^{(i)}(\vec{k} | \vec{k}') \cdot k_u \\ P^{(u)}(\vec{k} | \vec{k}') \cdot k_o & P^{(i)}(\vec{k} | \vec{k}') \cdot k_o \end{bmatrix} \cdot B_{\vec{k} \vec{k}'}$$



Outline

Directed random networks

Mixed random networks

Definition

Correlations

Mixed Random Network Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

The PoCverse
Mixed, correlated
random networks
25 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization


Triggering probabilities

Nutshell

References



Full generalization:




$$f_{\vec{\alpha}}(d+1) = \sum_{\vec{\alpha}'} R_{\vec{\alpha}\vec{\alpha}'} f_{\vec{\alpha}'}(d)$$

$R_{\vec{\alpha}\vec{\alpha}'}$ is the gain ratio matrix and has the form:

$$R_{\vec{\alpha}\vec{\alpha}'} = P_{\vec{\alpha}\vec{\alpha}'} \bullet k_{\vec{\alpha}\vec{\alpha}'} \bullet B_{\vec{\alpha}\vec{\alpha}'}$$




Full generalization:



$$f_{\vec{\alpha}}(d+1) = \sum_{\vec{\alpha}'} R_{\vec{\alpha}\vec{\alpha}'} f_{\vec{\alpha}'}(d)$$


$R_{\vec{\alpha}\vec{\alpha}'}$ is the gain ratio matrix and has the form:

$$R_{\vec{\alpha}\vec{\alpha}'} = P_{\vec{\alpha}\vec{\alpha}'} \bullet k_{\vec{\alpha}\vec{\alpha}'} \bullet B_{\vec{\alpha}\vec{\alpha}'}$$

 $P_{\vec{\alpha}\vec{\alpha}'}$ = conditional probability that a type λ' edge emanating from a type ν' node leads to a type ν node.




Full generalization:




$$f_{\vec{\alpha}}(d+1) = \sum_{\vec{\alpha}'} R_{\vec{\alpha}\vec{\alpha}'} f_{\vec{\alpha}'}(d)$$

$R_{\vec{\alpha}\vec{\alpha}'}$ is the gain ratio matrix and has the form:


$$R_{\vec{\alpha}\vec{\alpha}'} = P_{\vec{\alpha}\vec{\alpha}'} \bullet k_{\vec{\alpha}\vec{\alpha}'} \bullet B_{\vec{\alpha}\vec{\alpha}'}$$

 $P_{\vec{\alpha}\vec{\alpha}'}$ = conditional probability that a type λ' edge emanating from a type ν' node leads to a type ν node.

 $k_{\vec{\alpha}\vec{\alpha}'}$ = potential number of newly infected edges of type λ emanating from nodes of type ν .




Full generalization:





$$f_{\vec{\alpha}}(d+1) = \sum_{\vec{\alpha}'} R_{\vec{\alpha}\vec{\alpha}'} f_{\vec{\alpha}'}(d)$$

$R_{\vec{\alpha}\vec{\alpha}'}$ is the gain ratio matrix and has the form:

$$R_{\vec{\alpha}\vec{\alpha}'} = P_{\vec{\alpha}\vec{\alpha}'} \cdot k_{\vec{\alpha}\vec{\alpha}'} \cdot B_{\vec{\alpha}\vec{\alpha}'}$$


 $P_{\vec{\alpha}\vec{\alpha}'}$ = conditional probability that a type λ' edge emanating from a type ν' node leads to a type ν node.

 $k_{\vec{\alpha}\vec{\alpha}'}$ = potential number of newly infected edges of type λ emanating from nodes of type ν .

 $B_{\vec{\alpha}\vec{\alpha}'}$ = probability that a type ν node is eventually infected by a single infected type λ' link arriving from a neighboring node of type ν' .




Full generalization:





$$f_{\vec{\alpha}}(d+1) = \sum_{\vec{\alpha}'} R_{\vec{\alpha}\vec{\alpha}'} f_{\vec{\alpha}'}(d)$$


$R_{\vec{\alpha}\vec{\alpha}'}$ is the gain ratio matrix and has the form:

$$R_{\vec{\alpha}\vec{\alpha}'} = P_{\vec{\alpha}\vec{\alpha}'} \bullet k_{\vec{\alpha}\vec{\alpha}'} \bullet B_{\vec{\alpha}\vec{\alpha}'}$$

 $P_{\vec{\alpha}\vec{\alpha}'}$ = conditional probability that a type λ' edge emanating from a type ν' node leads to a type ν node.

 $k_{\vec{\alpha}\vec{\alpha}'}$ = potential number of newly infected edges of type λ emanating from nodes of type ν .

 $B_{\vec{\alpha}\vec{\alpha}'}$ = probability that a type ν node is eventually infected by a single infected type λ' link arriving from a neighboring node of type ν' .

 Generalized contagion condition:

$$\max|\mu| : \mu \in \sigma(\mathbf{R}) > 1$$



Outline

Directed random networks

Mixed random networks

Definition

Correlations

Mixed Random Network Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

The PoCverse
Mixed, correlated
random networks
27 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References





As we saw earlier, the triggering probability for simple contagion on random networks can be determined with a straightforward physical argument.

The PoCverse
Mixed, correlated
random networks
28 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization

Triggering probabilities

Nutshell

References



As we saw earlier, the triggering probability for simple contagion on random networks can be determined with a straightforward physical argument.

Two good things:

$$Q_{\text{trig}} = \sum_{k=0}^{\infty} \frac{kP_k}{\langle k \rangle} \cdot B_{k1} \cdot \left[1 - (1 - Q_{\text{trig}})^{k-1} \right],$$

$$P_{\text{trig}} = S_{\text{trig}} = \sum_k P_k \cdot \left[1 - (1 - Q_{\text{trig}})^k \right].$$

The PoCverse
Mixed, correlated
random networks
28 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization

Triggering probabilities

Nutshell

References



As we saw earlier, the triggering probability for simple contagion on random networks can be determined with a straightforward physical argument.

Two good things:

$$Q_{\text{trig}} = \sum_{k=0}^{\infty} \frac{k P_k}{\langle k \rangle} \cdot B_{k1} \cdot \left[1 - (1 - Q_{\text{trig}})^{k-1} \right],$$

$$P_{\text{trig}} = S_{\text{trig}} = \sum_k P_k \cdot \left[1 - (1 - Q_{\text{trig}})^k \right].$$

Equivalent to result found via the eldritch route of generating functions.



As we saw earlier, the triggering probability for simple contagion on random networks can be determined with a straightforward physical argument.

Two good things:

$$Q_{\text{trig}} = \sum_{k=0}^{\infty} \frac{k P_k}{\langle k \rangle} \cdot B_{k1} \cdot \left[1 - (1 - Q_{\text{trig}})^{k-1} \right],$$

$$P_{\text{trig}} = S_{\text{trig}} = \sum_k P_k \cdot \left[1 - (1 - Q_{\text{trig}})^k \right].$$

Equivalent to result found via the eldritch route of generating functions.

Generating functions arguably make some kinds of calculations easier (but perhaps we don't care about component sizes that much).



As we saw earlier, the triggering probability for simple contagion on random networks can be determined with a straightforward physical argument.

Two good things:

$$Q_{\text{trig}} = \sum_{k=0}^{\infty} \frac{k P_k}{\langle k \rangle} \cdot B_{k1} \cdot \left[1 - (1 - Q_{\text{trig}})^{k-1} \right],$$

$$P_{\text{trig}} = S_{\text{trig}} = \sum_k P_k \cdot \left[1 - (1 - Q_{\text{trig}})^k \right].$$

Equivalent to result found via the eldritch route of generating functions.

Generating functions arguably make some kinds of calculations easier (but perhaps we don't care about component sizes that much).

On the other hand, a plainspoken physical argument helps us generalize to correlated networks more easily.



Summary of triggering probabilities for uncorrelated networks: ^[3] □

I. Undirected, Uncorrelated—

$$Q_{\text{trig}} = \sum_{k'_u} P^{(u)}(k'_u | \cdot) B_{k'_u 1} \left[1 - (1 - Q_{\text{trig}})^{k'_u - 1} \right]$$

$$P_{\text{trig}} = S_{\text{trig}} = \sum_{k'_u} P(k'_u) \left[1 - (1 - Q_{\text{trig}})^{k'_u} \right]$$

Summary of triggering probabilities for uncorrelated networks: ^[3] □

I. Undirected, Uncorrelated—

$$Q_{\text{trig}} = \sum_{k'_u} P^{(u)}(k'_u | \cdot) B_{k'_u 1} \left[1 - (1 - Q_{\text{trig}})^{k'_u - 1} \right]$$

$$P_{\text{trig}} = S_{\text{trig}} = \sum_{k'_u} P(k'_u) \left[1 - (1 - Q_{\text{trig}})^{k'_u} \right]$$

II. Directed, Uncorrelated—

$$Q_{\text{trig}} = \sum_{k'_i, k'_o} P^{(u)}(k'_i, k'_o | \cdot) B_{k'_i 1} \left[1 - (1 - Q_{\text{trig}})^{k'_o} \right]$$

$$S_{\text{trig}} = \sum_{k'_i, k'_o} P(k'_i, k'_o) \left[1 - (1 - Q_{\text{trig}})^{k'_o} \right]$$

Summary of triggering probabilities for uncorrelated networks:

III. Mixed Directed and Undirected, Uncorrelated—

$$Q_{\text{trig}}^{(u)} = \sum_{\vec{k}'} P^{(u)}(\vec{k}' | \cdot) B_{\vec{k}'1} \left[1 - (1 - Q_{\text{trig}}^{(u)})^{k'_u - 1} (1 - Q_{\text{trig}}^{(o)})^{k'_o} \right]$$

$$Q_{\text{trig}}^{(o)} = \sum_{\vec{k}'} P^{(i)}(\vec{k}' | \cdot) B_{\vec{k}'1} \left[1 - (1 - Q_{\text{trig}}^{(u)})^{k'_u} (1 - Q_{\text{trig}}^{(o)})^{k'_o} \right]$$

$$S_{\text{trig}} = \sum_{\vec{k}'} P(\vec{k}') \left[1 - (1 - Q_{\text{trig}}^{(u)})^{k'_u} (1 - Q_{\text{trig}}^{(o)})^{k'_o} \right]$$



Summary of triggering probabilities for correlated networks:



IV. Undirected, Correlated— $Q_{\text{trig}}(k_u) =$

$$\sum_{k'_u} P^{(u)}(k'_u | k_u) B_{k'_u} \left[1 - (1 - Q_{\text{trig}}(k'_u))^{k'_u - 1} \right]$$

$$S_{\text{trig}} = \sum_{k'_u} P(k'_u) \left[1 - (1 - Q_{\text{trig}}(k'_u))^{k'_u} \right]$$



Summary of triggering probabilities for correlated networks:



IV. Undirected, Correlated— $Q_{\text{trig}}(k_u) =$

$$\sum_{k'_u} P^{(u)}(k'_u | k_u) B_{k'_u-1} \left[1 - (1 - Q_{\text{trig}}(k'_u))^{k'_u-1} \right]$$

$$S_{\text{trig}} = \sum_{k'_u} P(k'_u) \left[1 - (1 - Q_{\text{trig}}(k'_u))^{k'_u} \right]$$



V. Directed, Correlated— $Q_{\text{trig}}(k_i, k_o) =$

$$\sum_{k'_i, k'_o} P^{(u)}(k'_i, k'_o | k_i, k_o) B_{k'_i-1} \left[1 - (1 - Q_{\text{trig}}(k'_i, k'_o))^{k'_o} \right]$$

$$S_{\text{trig}} = \sum_{k'_i, k'_o} P(k'_i, k'_o) \left[1 - (1 - Q_{\text{trig}}(k'_i, k'_o))^{k'_o} \right]$$



Summary of triggering probabilities for correlated networks:


VI. Mixed Directed and Undirected, Correlated—

$$Q_{\text{trig}}^{(u)}(\vec{k}) = \sum_{\vec{k}'} P^{(u)}(\vec{k}' | \vec{k}) B_{\vec{k}'1} \left[1 - (1 - Q_{\text{trig}}^{(u)}(\vec{k}'))^{k'_u - 1} (1 - Q_{\text{trig}}^{(o)}(\vec{k}'))^{k'_o} \right]$$

$$Q_{\text{trig}}^{(o)}(\vec{k}) = \sum_{\vec{k}'} P^{(i)}(\vec{k}' | \vec{k}) B_{\vec{k}'1} \left[1 - (1 - Q_{\text{trig}}^{(u)}(\vec{k}'))^{k'_u} (1 - Q_{\text{trig}}^{(o)}(\vec{k}'))^{k'_o} \right]$$

$$S_{\text{trig}} = \sum_{\vec{k}'} P(\vec{k}') \left[1 - (1 - Q_{\text{trig}}^{(u)}(\vec{k}'))^{k'_u} (1 - Q_{\text{trig}}^{(o)}(\vec{k}'))^{k'_o} \right]$$

Nutshell:

 Mixed, correlated random networks with undirected and directed edges form natural inclusive generalization of purely undirected and purely directed random networks.

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network
Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Nutshell:

- ⊞ Mixed, correlated random networks with undirected and directed edges form natural inclusive generalization of purely undirected and purely directed random networks.
- ⊞ Spreading conditions and triggering probabilities of contagion processes can be determined using a direct, physical approach.

Directed random networks

Mixed random networks

Definition
Correlations

Mixed Random Network

Contagion




Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Nutshell:

-  Mixed, correlated random networks with undirected and directed edges form natural inclusive generalization of purely undirected and purely directed random networks.
-  Spreading conditions and triggering probabilities of contagion processes can be determined using a direct, physical approach.
-  These conditions can be generalized to arbitrary random networks with arbitrary node and edge types.

Directed random networks

Mixed random networks

Definition
Correlations

Mixed Random Network

Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



Nutshell:

- Mixed, correlated random networks with undirected and directed edges form natural inclusive generalization of purely undirected and purely directed random networks.
- Spreading conditions and triggering probabilities of contagion processes can be determined using a direct, physical approach.
- These conditions can be generalized to arbitrary random networks with arbitrary node and edge types.
- More generalizations: bipartite affiliation graphs and multilayer networks.

Directed random networks

Mixed random networks

Definition
Correlations

Mixed Random Network

Contagion

Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



References I

- [1] M. Boguñá and M. Ángeles Serrano.
Generalized percolation in random directed networks.
[Phys. Rev. E, 72:016106, 2005. pdf](#)
- [2] P. S. Dodds, K. D. Harris, and J. L. Payne.
Direct, physically motivated derivation of the contagion condition for spreading processes on generalized random networks.
[Phys. Rev. E, 83:056122, 2011. pdf](#)
- [3] K. D. Harris, J. L. Payne, and P. S. Dodds.
Direct, physically-motivated derivation of triggering probabilities for contagion processes acting on correlated random networks.
<https://arxiv.org/abs/1108.5398>, 2014.

The PoCverse
Mixed, correlated
random networks
34 of 35

Directed random
networks

Mixed random
networks

Definition
Correlations

Mixed Random
Network

Contagion


Spreading condition
Full generalization
Triggering probabilities

Nutshell

References



References II

- [4] M. E. J. Newman, S. H. Strogatz, and D. J. Watts.
Random graphs with arbitrary degree distributions
and their applications.
[Phys. Rev. E, 64:026118, 2001. pdf](#) 

The PoCverse
Mixed, correlated
random networks
35 of 35

Directed random
networks

Mixed random
networks

Definition

Correlations

Mixed Random
Network

Contagion

Spreading condition

Full generalization

Triggering probabilities

Nutshell

References

