Chapter 2: Lecture 1

Linear Algebra, Course 124C, Spring, 2009

Prof. Peter Dodds

Department of Mathematics & Statistics
University of Vermont

Basics:

- Instructor: Prof. Peter Dodds
- Lecture room and meeting times:
  367 Votey, Tuesday and Thursday, 1:00 pm to 2:15 pm
- Office: 203 Lord House, 16 Colchester Avenue
- E-mail: pdodds@uvm.edu
- Course website:
  `http://www.uvm.edu/~pdodds/teaching/courses/2009-01UVM-124/`
- Textbook: "Introduction to Linear Algebra" (3rd ed.) by Gilbert Strang; Wellesley-Cambridge Press.

Admin:

Paper products:

1. Outline
2. "The Fundamental Theorem of Linear Algebra" [1]
3. "Too Much Calculus" [2]

Office hours:

- Tuesday: 2:30 pm to 4:30 pm
  Thursday: 11:30 am to 12:30 pm
  Rm 203, Math Building

Admin:

Paper products:

1. Outline
2. “The Fundamental Theorem of Linear Algebra” [1]
3. “Too Much Calculus” [2]

Office hours:

- Tuesday: 2:30 pm to 4:30 pm
  Thursday: 11:30 am to 12:30 pm
  Rm 203, Math Building

Grading breakdown:

1. Assignments (40%)
   - Ten one-week assignments.
   - Lowest assignment score will be dropped.
   - The last assignment cannot be dropped!
   - Each assignment will have a random bonus point question which has nothing to do with linear algebra.
2. Midterm exams (35%)
   - Three 75 minutes tests distributed throughout the course, all of equal weighting.
3. Final exam (24%)
   - Three hours of pure happiness.
   - May 4, 8:00 am to 11:00 am; held in normal lecture room.

Grading breakdown:

1. Assignments (40%)
   - Ten one-week assignments.
   - Lowest assignment score will be dropped.
   - The last assignment cannot be dropped!
   - Each assignment will have a random bonus point question which has nothing to do with linear algebra.
2. Midterm exams (35%)
   - Three 75 minutes tests distributed throughout the course, all of equal weighting.
3. Final exam (24%)
   - Three hours of pure happiness.
   - May 4, 8:00 am to 11:00 am; held in normal lecture room.

Grading breakdown:

1. Assignments (40%)
    - Ten one-week assignments.
    - Lowest assignment score will be dropped.
    - The last assignment cannot be dropped!
    - Each assignment will have a random bonus point question which has nothing to do with linear algebra.
2. Midterm exams (35%)
    - Three 75 minutes tests distributed throughout the course, all of equal weighting.
3. Final exam (24%)
    - Three hours of pure happiness.
    - May 4, 8:00 am to 11:00 am; held in normal lecture room.

Grading breakdown:

1. Homework (0%)—Problems assigned online from the textbook. Doing these exercises will be most beneficial and will increase happiness.
2. General attendance (1%)—it is extremely desirable that students attend class, and class presence will be taken into account if a grade is borderline.

How grading works:

Questions are worth 3 points according to the following scale:

- 3 = correct or very nearly so.
- 2 = acceptable but needs some revisions.
- 1 = needs major revisions.
- 0 = way off.

Schedule:

The course will mainly cover chapters 2 through 6 of the textbook. (You should know all about Chapter 1.)

Week # (dates)	Tuesday	Thursday
1 (1/13 and 1/15)	Lecture	Lecture + A 1
2 (1/20 and 1/22)	Lecture	Lecture + A 2
3 (1/27 and 1/29)	Lecture	Lecture + A 3
4 (2/3 and 2/5)	Lecture	*Test 1*
5 (2/10 and 2/12)	Lecture	Lecture + A 4
6 (2/17 and 2/19)	Lecture	Lecture + A 5
7 (2/24 and 2/26)	Lecture	Lecture + A 6
8 (3/3 and 3/5)	Town Recess	Lecture
9 (3/10 and 3/12)	Spring recess	Spring recess
10 (3/17 and 3/19)	*Test 2*	Lecture + A 7
11 (3/24 and 3/26)	Lecture	Lecture + A 8
12 (3/31 and 4/2)	Lecture	Lecture + A 9
13 (4/7 and 4/9)	Lecture	*Test 3*
13 (4/14 and 4/16)	Lecture	Lecture + A 10
14 (4/21 and 4/23)	Lecture	Lecture
15 (4/28)	Lecture	—

Important dates:

1. Classes run from Monday, January 12 to Wednesday, April 29.
2. Add/Drop, Audit, Pass/No Pass deadline—Monday, January 26.
3. Last day to withdraw—Friday, March 30.
4. Reading and exam period—Thursday, April 30th to Friday, May 8.

More stuff:

Do check your zoo account for updates regarding the course.

Academic assistance: Anyone who requires assistance in any way (as per the ACCESS program or due to athletic endeavors), please see or contact me as soon as possible.

More stuff:

Being good people:

1. In class there will be no electronic gadgetry, no cell phones, no beeping, no text messaging, etc. You really just need your brain, some paper, and a writing implement here (okay, and Matlab or similar).
2. Second, I encourage you to email me questions, ideas, comments, etc., about the class but request that you please do so in a respectful fashion.
3. Finally, as in all UVM classes, Academic honesty will be expected and departures will be dealt with appropriately. See `http://www.uvm.edu/cses/` for guidelines.

More stuff:

Being good people:

1. In class there will be no electronic gadgetry, no cell phones, no beeping, no text messaging, etc. You really just need your brain, some paper, and a writing implement here (okay, and Matlab or similar).
2. Second, I encourage you to email me questions, ideas, comments, etc., about the class but request that you please do so in a respectful fashion.
3. Finally, as in all UVM classes, Academic honesty will be expected and departures will be dealt with appropriately. See `http://www.uvm.edu/cses/` for guidelines.

More stuff:

Being good people:

1. In class there will be no electronic gadgetry, no cell phones, no beeping, no text messaging, etc. You really just need your brain, some paper, and a writing implement here (okay, and Matlab or similar).
2. Second, I encourage you to email me questions, ideas, comments, etc., about the class but request that you please do so in a respectful fashion.
3. Finally, as in all UVM classes, Academic honesty will be expected and departures will be dealt with appropriately. See `http://www.uvm.edu/cses/` for guidelines.

More stuff:

Late policy: Unless in the case of an emergency (a real one) or if an absence has been predeclared and a make-up version sorted out, assignments that are not turned in on time or tests that are not attended will be given 0%.

Computing: Students are encouraged to use Matlab or something similar to check their work.

Note: for assignment problems, written details of calculations will be required.

Grading:

A+	97–100	B+	87–89	C+	77–79	D+	67–69
A	93–96	B	83–86	C	73–76	D	63–66
A-	90–92	B-	80–82	C-	70–72	D-	60–62

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:
If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:

If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:

If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:

If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:

If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:

If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is

a body of mathematics
that deals with discrete problems.

Many things are discrete:

- Information (0's & 1's, letters, words)
- People (sociology)
- Networks (the Web, people again, food webs, ...)
- Sounds (musical notes)

Even more:

If real data is continuous, we almost always discretize it (0's and 1's)

Why are we doing this?

Linear Algebra is used in many fields to solve problems:

- Engineering
- Computer Science (Google's Pagerank)
- Physics
- Economics
- Biology
- Ecology
- ...

Why are we doing this?

Linear Algebra is used in many fields to solve problems:

- Engineering
- Computer Science (Google's Pagerank)
- Physics
- Economics
- Biology
- Ecology
- ...

Linear Algebra is as important as calculus.

Matrices as gadgets:

A transforms $\vec{x}$ into $\vec{x}'$ through multiplication

$$\vec{x}' = A\vec{x}$$

Can use matrices to:

- Grow vectors
- Shrink vectors
- Rotate vectors
- Flip vectors
- Do all these things to different directions

Image approximation (80x60)

$$A = \sum_{i=1}^{1} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$

 

Image approximation (80x60)

$$A = \sum_{i=1}^{2} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$





Image approximation (80x60)

$$A = \sum_{i=1}^{3} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$





Image approximation (80x60)

$$A = \sum_{i=1}^{4} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{5} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{6} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{7} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{8} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{9} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{10} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{20} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{30} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{40} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{50} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Image approximation (80x60)

$$A = \sum_{i=1}^{60} \sigma_i \hat{u}_i \hat{v}_i^{\mathrm{T}}$$



Three key problems of Linear Algebra

1. Given a matrix A and a vector $\vec{b}$, find $\vec{x}$ such that

$$A\vec{x} = \vec{b}.$$

2. Eigenvalue problem: Given A, find λ and $\vec{v}$ such that

$$A\vec{v} = \lambda\vec{v}.$$

3. Coupled linear differential equations:

$$\frac{\mathrm{d}}{\mathrm{d}t}y(t) = A\,y(t)$$

- Our focus will be largely on #1, partly on #2.

Three key problems of Linear Algebra

1. Given a matrix A and a vector $\vec{b}$, find $\vec{x}$ such that

$$A\vec{x} = \vec{b}.$$

2. Eigenvalue problem: Given A, find λ and $\vec{v}$ such that

$$A\vec{v} = \lambda\vec{v}.$$

3. Coupled linear differential equations:

$$\frac{\mathrm{d}}{\mathrm{d}t}y(t) = A\,y(t)$$

- Our focus will be largely on #1, partly on #2.

Three key problems of Linear Algebra

1. Given a matrix A and a vector $\vec{b}$, find $\vec{x}$ such that
$$A\vec{x} = \vec{b}.$$
2. Eigenvalue problem: Given A, find λ and $\vec{v}$ such that
$$A\vec{v} = \lambda\vec{v}.$$
3. Coupled linear differential equations:
$$\frac{\mathrm{d}}{\mathrm{d}t} y(t) = A\, y(t)$$

- Our focus will be largely on #1, partly on #2.

Three key problems of Linear Algebra

1. Given a matrix A and a vector $\vec{b}$, find $\vec{x}$ such that
$$A\vec{x} = \vec{b}.$$
2. Eigenvalue problem: Given A, find λ and $\vec{v}$ such that
$$A\vec{v} = \lambda\vec{v}.$$
3. Coupled linear differential equations:
$$\frac{\mathrm{d}}{\mathrm{d}t}y(t) = A\,y(t)$$

- Our focus will be largely on #1, partly on #2.

Major course objective:

To deeply understand the equation $A\vec{x} = \vec{b}$, the Fundamental Theorem of Linear Algebra, and the following picture:



Major course objective:

To deeply understand the equation $A\vec{x} = \vec{b}$, the Fundamental Theorem of Linear Algebra, and the following picture:



What is going on here? We have 26 lectures to find out...

Our friend $A\vec{x} = \vec{b}$

Broadly speaking, $A\vec{x} = \vec{b}$ translates as follows:

- $\vec{b}$ represents reality (e.g., music, structure)
- A contains building blocks (e.g., notes, shapes)
- $\vec{x}$ specifies how we combine our building blocks to represent $\vec{b}$.

Our friend $A\vec{x} = \vec{b}$

Broadly speaking, $A\vec{x} = \vec{b}$ translates as follows:

- $\vec{b}$ represents reality (e.g., music, structure)
- A contains building blocks (e.g., notes, shapes)
- $\vec{x}$ specifies how we combine our building blocks to represent $\vec{b}$.

Our friend $A\vec{x} = \vec{b}$

Broadly speaking, $A\vec{x} = \vec{b}$ translates as follows:

- $\vec{b}$ represents reality (e.g., music, structure)
- A contains building blocks (e.g., notes, shapes)
- $\vec{x}$ specifies how we combine our building blocks to represent $\vec{b}$.

Our friend $A\vec{x} = \vec{b}$

Broadly speaking, $A\vec{x} = \vec{b}$ translates as follows:

- $\vec{b}$ represents reality (e.g., music, structure)
- A contains building blocks (e.g., notes, shapes)
- $\vec{x}$ specifies how we combine our building blocks to represent $\vec{b}$.

Our friend $A\vec{x} = \vec{b}$

Broadly speaking, $A\vec{x} = \vec{b}$ translates as follows:

- $\vec{b}$ represents reality (e.g., music, structure)
- A contains building blocks (e.g., notes, shapes)
- $\vec{x}$ specifies how we combine our building blocks to represent $\vec{b}$.

How can we disentangle an orchestra's sound?

Our friend $A\vec{x} = \vec{b}$

Broadly speaking, $A\vec{x} = \vec{b}$ translates as follows:

- $\vec{b}$ represents reality (e.g., music, structure)
- A contains building blocks (e.g., notes, shapes)
- $\vec{x}$ specifies how we combine our building blocks to represent $\vec{b}$.

How can we disentangle an orchestra's sound?

What about pictures, waves, signals, ...?

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

If we can represent reality as a superposition (or combination) of simple elements, we can do many things:

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

If we can represent reality as a superposition (or combination) of simple elements, we can do many things:

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

If we can represent reality as a superposition (or combination) of simple elements, we can do many things:

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

If we can represent reality as a superposition (or combination) of simple elements, we can do many things:

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

If we can represent reality as a superposition (or combination) of simple elements, we can do many things:

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Our friend $A\vec{x} = \vec{b}$

What does knowing $\vec{x}$ give us?

If we can represent reality as a superposition (or combination) of simple elements, we can do many things:

- Compress information
- See how we can alter information
- Find a system's simplest representation
- Find a system's most important elements
- See how to adjust a system in a principled defined way

Three ways to understand $A\vec{x} = \vec{b}$:

- Way 1: The Row Picture
- Way 2: The Column Picture
- Way 3: The Matrix Picture

Example:

- $$\begin{array}{rcrcr} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$
- Call this a 2 by 2 system of equations.
- 2 equations with 2 unknowns.
- Standard method of solving by adding and subtracting multiples of equations from each other

Three ways to understand $A\vec{x} = \vec{b}$:

- Way 1: The Row Picture
- Way 2: The Column Picture
- Way 3: The Matrix Picture

Example:

- $$\begin{array}{rcrcr} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$
- Call this a 2 by 2 system of equations.
- 2 equations with 2 unknowns.
- Standard method of solving by adding and subtracting multiples of equations from each other

Three ways to understand $A\vec{x} = \vec{b}$:

- Way 1: The Row Picture
- Way 2: The Column Picture
- Way 3: The Matrix Picture

Example:

- $$\begin{array}{rcrcr} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$
- Call this a 2 by 2 system of equations.
- 2 equations with 2 unknowns.
- Standard method of solving by adding and subtracting multiples of equations from each other

Three ways to understand $A\vec{x} = \vec{b}$:

- Way 1: The Row Picture
- Way 2: The Column Picture
- Way 3: The Matrix Picture

Example:

- $$\begin{array}{rcrcl} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$
- Call this a 2 by 2 system of equations.
- 2 equations with 2 unknowns.
- Standard method of solving by adding and subtracting multiples of equations from each other

Three ways to understand $A\vec{x} = \vec{b}$:

- Way 1: The Row Picture
- Way 2: The Column Picture
- Way 3: The Matrix Picture

Example:

- $$\begin{array}{rcrcr} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$
- Call this a 2 by 2 system of equations.
- 2 equations with 2 unknowns.
- Standard method of solving by adding and subtracting multiples of equations from each other = Row Picture

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection: (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
(a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point —One, unique solution
2. Lines are parallel and disjoint
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point —One, unique solution
2. Lines are parallel and disjoint —No solutions
3. Lines are the same

Three ways to understand $A\vec{x} = \vec{b}$:

Row Picture—what we are doing:

- (a) Finding intersection of two lines
- (b) Finding the values of x_1 and x_2 for which both equations are satisfied (true/happy)
- A splendid and deep connection:
  (a) Geometry $\rightleftharpoons$ (b) Algebra

Three possible kinds of solution:

1. Lines intersect at one point —One, unique solution
2. Lines are parallel and disjoint —No solutions
3. Lines are the same —Infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

The column picture:

- Column vectors are 'building blocks'
- Key idea: try to 'reach' $\vec{b}$ by combining multiples of column vectors $\vec{a}_1$ and $\vec{a}_2$.

Three ways to understand $A\vec{x} = \vec{b}$:

The column picture:

See

$$\begin{array}{rcccc} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$

- Column vectors are 'building blocks'
- Key idea: try to 'reach' $\vec{b}$ by combining multiples of column vectors $\vec{a}_1$ and $\vec{a}_2$.

Three ways to understand $A\vec{x} = \vec{b}$:

The column picture:

See

$$\begin{array}{rcrcl} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$

as

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

- Column vectors are ‘building blocks’
- Key idea: try to ‘reach’ $\vec{b}$ by combining multiples of column vectors $\vec{a}_1$ and $\vec{a}_2$.

Three ways to understand $A\vec{x} = \vec{b}$:

The column picture:

See

$$\begin{array}{rcrcl} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$

as

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

General problem

$$x_1 \vec{a}_1 + x_2 \vec{a}_2 = \vec{b}$$

- Column vectors are 'building blocks'
- Key idea: try to 'reach' $\vec{b}$ by combining multiples of column vectors $\vec{a}_1$ and $\vec{a}_2$.

Three ways to understand $A\vec{x} = \vec{b}$:

The column picture:

See

$$\begin{array}{rcrcr} -x_1 & + & x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 4 \end{array}$$

as

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

General problem

$$x_1 \vec{a}_1 + x_2 \vec{a}_2 = \vec{b}$$

- Column vectors are ‘building blocks’
- Key idea: try to ‘reach’ $\vec{b}$ by combining multiples of column vectors $\vec{a}_1$ and $\vec{a}_2$.

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Three ways to understand $A\vec{x} = \vec{b}$:

We love the column picture:

- Intuitive.
- Generalizes easily to many dimensions.

Three possible kinds of solution:

1. $\vec{a}_1 \not\parallel \vec{a}_2$: 1 solution
2. $\vec{a}_1 \parallel \vec{a}_2 \not\parallel \vec{b}$: No solutions
3. $\vec{a}_1 \parallel \vec{a}_2 \parallel \vec{b}$: infinitely many solutions

Assuming neither $\vec{a}_1$ or $\vec{a}_1$ are $\vec{0}$.

Three ways to understand $A\vec{x} = \vec{b}$:

Difficulties:

- Do we give up if $A\vec{x} = \vec{b}$ has no solution?
- No! We can still find the $\vec{x}$ that gets us as close to $\vec{b}$ as possible.
- Method of approximation—very important!
- We may not have the right building blocks but we can do our best.

Three ways to understand $A\vec{x} = \vec{b}$:

Difficulties:

- Do we give up if $A\vec{x} = \vec{b}$ has no solution?
- No! We can still find the $\vec{x}$ that gets us as close to $\vec{b}$ as possible.
- Method of approximation—very important!
- We may not have the right building blocks but we can do our best.

Three ways to understand $A\vec{x} = \vec{b}$:

Difficulties:

- Do we give up if $A\vec{x} = \vec{b}$ has no solution?
- No! We can still find the $\vec{x}$ that gets us as close to $\vec{b}$ as possible.
- Method of approximation—very important!
- We may not have the right building blocks but we can do our best.

Three ways to understand $A\vec{x} = \vec{b}$:

Difficulties:

- Do we give up if $A\vec{x} = \vec{b}$ has no solution?
- No! We can still find the $\vec{x}$ that gets us as close to $\vec{b}$ as possible.
- Method of approximation—very important!
- We may not have the right building blocks but we can do our best.

Three ways to understand $A\vec{x} = \vec{b}$:

The Matrix Picture:

A is now an operator:

- A transforms $\vec{x}$ into $\vec{b}$.
- In general, A does two things to $\vec{x}$:
  1. Rotation
  2. Dilation (stretching/contraction)

Three ways to understand $A\vec{x} = \vec{b}$:

The Matrix Picture:

Now see

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

A is now an operator:

- A transforms $\vec{x}$ into $\vec{b}$.
- In general, A does two things to $\vec{x}$:
  1. Rotation
  2. Dilation (stretching/contraction)

Three ways to understand $A\vec{x} = \vec{b}$:

The Matrix Picture:

Now see

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

as

$$A\vec{x} = \vec{b} : \begin{bmatrix} -1 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

A is now an operator:

- A transforms $\vec{x}$ into $\vec{b}$.
- In general, A does two things to $\vec{x}$:
  1. Rotation
  2. Dilation (stretching/contraction)

Three ways to understand $A\vec{x} = \vec{b}$:

The Matrix Picture:

Now see

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

as

$$A\vec{x} = \vec{b} : \begin{bmatrix} -1 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

A is now an operator:

- A transforms $\vec{x}$ into $\vec{b}$.
- In general, A does two things to $\vec{x}$:
  1. Rotation
  2. Dilation (stretching/contraction)

Three ways to understand $A\vec{x} = \vec{b}$:

The Matrix Picture:

Now see

$$x_1 \begin{bmatrix} -1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}.$$

as

$$A\vec{x} = \vec{b} : \begin{bmatrix} -1 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

A is now an operator:

- A transforms $\vec{x}$ into $\vec{b}$.
- In general, A does two things to $\vec{x}$:
  1. Rotation
  2. Dilation (stretching/contraction)

The Matrix Picture

Key idea in linear algebra:

- Decomposition (or factorization) of matrices.
- Matrices can often be written as products or sums of simpler matrices
- $A = LU$, $A = QR$, $A = U\Sigma V^{\mathrm{T}}$, $A = \sum_i \lambda_i \vec{v}\vec{v}^{\mathrm{T}}$, ...

The Matrix Picture

Key idea in linear algebra:

- Decomposition (or factorization) of matrices.
- Matrices can often be written as products or sums of simpler matrices
- $A = LU$, $A = QR$, $A = U\Sigma V^{\mathrm{T}}$, $A = \sum_i \lambda_i \vec{v}\vec{v}^{\mathrm{T}}$, ...

The Matrix Picture

Key idea in linear algebra:

- Decomposition (or factorization) of matrices.
- Matrices can often be written as products or sums of simpler matrices
- $A = LU$, $A = QR$, $A = U\Sigma V^{\mathrm{T}}$, $A = \sum_i \lambda_i \vec{v}\vec{v}^{\mathrm{T}}$, ...

The truth about mathematics

The Colbert Report on Math (February 7, 2006)

References I

G. Strang.
The fundamental theorem of linear algebra.
The American Mathematical Monthly, 100(9):848–855, 1993. pdf (⊞)

G. Strang.
Too much calculus, 2002.
SIAM Linear Algebra Activity Group Newsletter. pdf (⊞)