

# Lecture 3 (Chapter 2)

## Linear Algebra, Course 124A, Fall, 2009

Prof. Peter Dodds

Department of Mathematics & Statistics  
University of Vermont



Licensed under the *Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License*.

Solving  $A\vec{x} = \vec{b}$ :Solving  $A\vec{x} = \vec{b}$ 

- ▶ We (people + computers) solve systems of linear equations by a systematic method of **Elimination** followed by **Back substitution**
- ▶ Due to our man Gauss, hence Gaussian elimination.
- ▶ Our first example:

$$\begin{array}{rclcl} -x_1 & + & 3x_2 & = & 1 \\ 2x_1 & + & x_2 & = & 5 \end{array}$$

# Gaussian elimination:

## Basic elimination rules (roughly):

1. Strategically, mechanically remove unwanted entries by subtracting a multiple of a row from another.
  2. Swap rows if needed to create an 'upper triangular form'
- e.g.

$$\begin{array}{rcl} & x_2 & = & 3 \\ 2x_1 & - & x_2 & = & -1 \end{array} \Rightarrow \begin{array}{rcl} 2x_1 & - & x_2 & = & -1 \\ & & x_2 & = & 3 \end{array}$$

# Gaussian elimination:

Solving  $A\vec{x} = \vec{b}$ 

Solve:

$$2x_1 - 3x_2 = 3$$

$$4x_1 - 5x_2 + x_3 = 7$$

$$2x_1 - x_2 - 3x_3 = 5$$

# Gaussian elimination:

## Summary:

Using **row operations**, we turned this problem:

$$A\vec{x} = \vec{b} : \begin{bmatrix} 2 & -3 & 0 \\ 4 & -5 & 1 \\ 2 & -1 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 5 \end{bmatrix}$$

into this problem:

$$U\vec{x} = \vec{d} : \begin{bmatrix} 2 & -3 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix}$$

and the latter is **easy to solve** using **back substitution**.

## Gaussian elimination:

### Defn:

The entries along  $U$ 's main diagonal are the **pivots** of  $A$ .  
(The pivots are hidden—elimination finds them.)

### Defn:

A matrix with only zeros below the main diagonal is called **upper triangular**. A matrix with only zeros above the main diagonal is called **lower triangular**. We get from  $A$  to  $U$  and the latter is always upper triangular.

### Defn:

**Singular** means a system has no unique solution.

- ▶ It may have no solutions or infinitely many solutions.
- ▶ Singular = archaic way of saying 'messed up.'

### Truth:

If at least one pivot is zero, the matrix will be **singular**.  
(but the reverse is not necessarily true).

# Gaussian elimination:

## The one true method:

- ▶ We simplify  $A$  using elimination in **the same way every time**.
- ▶ Eliminate entries one column at a time, moving left to right, and down each column.

$$\begin{array}{cccccccc}
 X & + & X & + & X & + & X & = & X \\
 1 \downarrow & + & X & + & X & + & X & = & X \\
 2 \downarrow & + & 4 \downarrow & + & X & + & X & = & X \\
 3 \nearrow & + & 5 \rightarrow & + & 6 & + & X & = & X
 \end{array}$$

## Gaussian elimination:

- ▶ To eliminate entry in row  $i$  of  $j$ th column, subtract a multiple  $l_{ij}$  of the  $j$ th row from  $i$ .
- ▶ For example:

$$\begin{array}{rccccrcr}
 2x_1 & + & 3x_2 & + & -2x_3 & + & x_4 & = & 1 \\
 x_1 & - & 7x_2 & + & 3x_3 & + & x_4 & = & 1 \\
 -x_1 & - & 3x_2 & - & x_3 & + & 5x_4 & = & -2 \\
 2x_1 & + & x_2 & - & 2x_3 & + & 2x_4 & = & 0
 \end{array}$$

$$l_{21} = 1/2, l_{31} = -1/2, l_{41} = ?.$$

- ▶ **Note:** we cannot find  $l_{32}$  etc., until we are finished with row 1. Pivots are hidden!
- ▶ **Note:** the denominator of each  $l_{ij}$  multiplier is the pivot in the  $j$ th column.