

## Lecture 2/25—Chapter 2

Linear Algebra  
MATH 124, Fall, 2010

Prof. Peter Dodds

Department of Mathematics & Statistics  
Center for Complex Systems  
Vermont Advanced Computing Center  
University of Vermont



Licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License.

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



1 of 9

## Gaussian elimination:

### Basic elimination rules (roughly):

1. Strategically, mechanically remove unwanted entries by subtracting a multiple of a row from another.
2. Swap rows if needed to create an 'upper triangular form'

e.g.

$$\begin{array}{r} x_2 = 3 \\ 2x_1 - x_2 = -1 \end{array} \cdot \begin{array}{r} 2x_1 - x_2 = -1 \\ x_2 = 3 \end{array}$$

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



4 of 9

## Outline

Solving  $A\vec{x} = \vec{b}$

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



2 of 9

## Gaussian elimination:

Solve:

$$\begin{array}{r} 2x_1 - 3x_2 = 3 \\ 4x_1 - 5x_2 + x_3 = 7 \\ 2x_1 - x_2 - 3x_3 = 5 \end{array}$$

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



5 of 9

## Solving $A\vec{x} = \vec{b}$ :

- ▶ We (people + computers) solve systems of linear equations by a systematic method of **Elimination** followed by **Back substitution**
- ▶ Due to our man Gauss, hence Gaussian elimination.
- ▶ Our first example:

$$\begin{array}{r} -x_1 + 3x_2 = 1 \\ 2x_1 + x_2 = 5 \end{array}$$



3 of 9

## Gaussian elimination:

### Summary:

Using **row operations**, we turned this problem:

$$A\vec{x} = \vec{b} : \begin{bmatrix} 2 & -3 & 0 \\ 4 & -5 & 1 \\ 2 & -1 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 5 \end{bmatrix}$$

into this problem:

$$U\vec{x} = \vec{d} : \begin{bmatrix} 2 & -3 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix}$$

and the latter is **easy to solve** using **back substitution**.

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



6 of 9

## Gaussian elimination:

### Defn:

The entries along  $U$ 's main diagonal are the **pivots** of  $A$ .  
(The pivots are hidden—elimination finds them.)

### Defn:

A matrix with only zeros below the main diagonal is called **upper triangular**. A matrix with only zeros above the main diagonal is called **lower triangular**. We get from  $A$  to  $U$  and the latter is always upper triangular.

### Defn:

**Singular** means a system has no unique solution.

- ▶ It may have no solutions or infinitely many solutions.
- ▶ Singular = archaic way of saying 'messed up.'

### Truth:

If at least one pivot is zero, the matrix will be **singular**.  
(but the reverse is not necessarily true).

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



7 of 9

## Gaussian elimination:

### The one true method:

- ▶ We simplify  $A$  using elimination in **the same way every time**.
- ▶ Eliminate entries one column at a time, moving left to right, and down each column.

$$\begin{array}{cccccc} X & + & X & + & X & + & X & = & X \\ 1 \downarrow & + & X & + & X & + & X & = & X \\ 2 \downarrow & + & 4 \downarrow & + & X & + & X & = & X \\ 3 \nearrow & + & 5 \rightarrow & + & 6 & + & X & = & X \end{array}$$

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



8 of 9

## Gaussian elimination:

- ▶ To eliminate entry in row  $i$  of  $j$ th column, subtract a multiple  $\ell_{ij}$  of the  $j$ th row from  $i$ .
- ▶ For example:

$$\begin{array}{cccccc} 2x_1 & + & 3x_2 & + & -2x_3 & + & x_4 & = & 1 \\ x_1 & - & 7x_2 & + & 3x_3 & + & x_4 & = & 1 \\ -x_1 & - & 3x_2 & - & x_3 & + & 5x_4 & = & -2 \\ 2x_1 & + & x_2 & - & 2x_3 & + & 2x_4 & = & 0 \end{array}$$

$$\ell_{21} = 1/2, \ell_{31} = -1/2, \ell_{41} = ?.$$

- ▶ **Note:** we cannot find  $\ell_{32}$  etc., until we are finished with row 1. Pivots are hidden!
- ▶ **Note:** the denominator of each  $\ell_{ij}$  multiplier is the pivot in the  $j$ th column.

Ch. 2: Lec. 2

Solving  $A\vec{x} = \vec{b}$



9 of 9