

Complex Networks, CSYS/MATH 303—Assignment 2 University of Vermont, Spring 2011

Dispersed: Thursday, February 3 (the day before the Teletherm), 2011.

Due: By start of lecture, 2:30 pm, Thursday, February 17, 2011.

Some useful reminders:

Instructor: Peter Dodds

Office: Farrell Hall, second floor, Trinity Campus

E-mail: peter.dodds@uvm.edu

Office hours: 1:00 pm to 3:00 pm, Wednesday

Course website: <http://www.uvm.edu/~pdodds/teaching/courses/2011-01UVM-303>

All parts are worth 3 points unless marked otherwise. Please show all your working clearly and list the names of others with whom you collaborated.

Graduate students are requested to use \LaTeX (or related variant).

1. Tokunaga's law is statistical but we can consider a rigid version. Take $T_1 = 2$ and $R_T = 2$ and draw an example network of order $\Omega = 4$ with these parameters.
2. Show $R_s = R_\ell$. In other words show that Horton's law of stream segments matches that of main stream lengths.
3. Tokunaga's law implies Horton's laws:

In lectures, we established the following:

$$n_\omega = \underbrace{2n_{\omega+1}}_{\text{generation}} + \sum_{\omega'=\omega+1}^{\Omega} \underbrace{T_{\omega'-\omega}n_{\omega'}}_{\text{absorption}}$$

From here, derive Horton's law for stream numbers: $n_\omega/n_{\omega+1} = R_n$, where $R_n > 1$ and is independent of ω , and find R_n in terms of Tokunaga's two parameters T_1 and R_T .

4. Show $R_n = R_a$ by using Tokunaga's law to find the average area of an order ω basin, \bar{a}_ω , in terms of the average area of basins of order 1 to $\omega - 1$.
(In lectures, we use Horton's laws to roughly demonstrate this result.)
5. For river networks, basin areas are distributed according to $P(a) \propto a^{-\tau}$.
Determine the exponent τ in terms of the Horton ratios R_n and R_s .