

Complex Networks, CSYS/MATH 303—Assignment 3
University of Vermont, Spring 2011

Dispersed: Sunday, February 20, 2011.

Due: By start of lecture, 2:30 pm, Thursday, February 24, 2011.

Some useful reminders:

Instructor: Peter Dodds

Office: Farrell Hall, second floor, Trinity Campus

E-mail: peter.dodds@uvm.edu

Office hours: 11:45 am to 12:15 pm and 1:00 pm to 3:00 pm, Wednesday

Course website: <http://www.uvm.edu/~pdodds/teaching/courses/2011-01UVM-303>

All parts are worth 3 points unless marked otherwise. Please show all your working clearly and list the names of others with whom you collaborated.

Graduate students are requested to use L^AT_EX (or related variant).

Supply networks and allometry:

1. From lectures on Supply Networks:

Show that for large V and $0 < \epsilon < 1/2$

$$\min V_{\text{net}} \propto \int_{\Omega_{d,D}(V)} \rho \|\vec{x}\|^{1-2\epsilon} d\vec{x} \sim \rho V^{1+\gamma_{\max}(1-2\epsilon)}$$

Reminders: we defined $L_i = c_i^{-1} V^{\gamma_i}$ where $\gamma_1 + \gamma_2 + \dots + \gamma_d = 1$, $\gamma_1 = \gamma_{\max} \geq \gamma_2 \geq \dots \geq \gamma_d$, and $c = \prod_i c_i \leq 1$ is a shape factor.

Hints: assume the first k lengths scale in the same way with

$\gamma_1 = \dots = \gamma_k = \gamma_{\max}$, and write $\|\vec{x}\| = (x_1^2 + x_2^2 + \dots + x_d^2)^{1/2}$.

2. Consider a set of rectangular areas with side lengths L_1 and L_2 such that $L_1 \propto A^{\gamma_1}$ and $L_2 \propto A^{\gamma_2}$ where A is area and $\gamma_1 + \gamma_2 = 1$. Assume $\gamma_1 > \gamma_2$ and that $\epsilon = 0$.


Now imagine that material has to be distributed from a central source in each of these areas to sinks distributed with density $\rho(A)$, and that these sinks draw the same amount of material per unit time independent of L_1 and L_2 .

Find an exact form for how the volume of the most efficient distribution network scales with overall area $A = L_1 L_2$. (Hint: you will have to set up a double integration over the rectangle.)

If network volume must remain a constant fraction of overall area, determine the maximal scaling of sink density ρ with A .

Extra hints:

- Integrate over triangles as follows.
- You need to only perform calculations for one triangle.



3. (a) For a family of d -dimensional regions, with scaling as per Question 1, determine, to leading order, the scaling of hyper-surface area S with volume V . In other words, find the exponent β in $S \propto V^\beta$ as $V \rightarrow \infty$. Assume that nothing peculiar happens with the shapes (as we have always implicitly done), in that there is no fractal roughening. Hint: figure out how the circumference for the rectangles in the previous question scales with area A . For d dimensions, think about how the hyper-surface area of a hyperrectangle (or orthotope) would scale.
- (b) For general d , what is the minimum and maximum possible values of β and for what values of the γ_i does these extrema occur?