

Mechanisms for Generating Power-Law Size Distributions, Part 1

Principles of Complex Systems | @pocsvox
 CSYS/MATH 300, Fall, 2016 | #FallPoCS2016

Prof. Peter Dodds | @peterdodds


Dept. of Mathematics & Statistics | Vermont Complex Systems Center
 Vermont Advanced Computing Core | University of Vermont



Licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License.

PoCS | @pocsvox
 Power-Law Mechanisms, Pt. 1

Random Walks
 The First Return Problem
 Examples
 Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO
 References



1 of 60

PoCS | @pocsvox
 Power-Law Mechanisms, Pt. 1

Random Walks
 The First Return Problem
 Examples
 Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO
 References



4 of 60

These slides are brought to you by:



PoCS | @pocsvox
 Power-Law Mechanisms, Pt. 1

Random Walks
 The First Return Problem
 Examples
 Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO
 References



2 of 60

PoCS | @pocsvox
 Power-Law Mechanisms, Pt. 1

Random Walks
 The First Return Problem
 Examples
 Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO
 References



5 of 60

Outline


Random Walks
 The First Return Problem
 Examples

Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO

References

PoCS | @pocsvox
 Power-Law Mechanisms, Pt. 1

Random Walks
 The First Return Problem
 Examples
 Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO
 References



3 of 60

PoCS | @pocsvox
 Power-Law Mechanisms, Pt. 1

Random Walks
 The First Return Problem
 Examples
 Variable transformation
 Basics
 Holtsmark's Distribution
 PLIPLO
 References



6 of 60

Mechanisms:

A powerful story in the rise of complexity:

- 🔗 structure arises out of randomness.
- 🔗 Exhibit A: Random walks. [↗](#)

The essential random walk:

- 🔗 One spatial dimension.
- 🔗 Time and space are discrete
- 🔗 Random walker (e.g., a drunk) starts at origin $x = 0$.
- 🔗 Step at time t is ϵ_t :

$$\epsilon_t = \begin{cases} +1 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/2 \end{cases}$$

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
🔗 7 of 60

Variances sum: [↗*](#)

$$\begin{aligned} \text{Var}(x_t) &= \text{Var}\left(\sum_{i=1}^t \epsilon_i\right) \\ &= \sum_{i=1}^t \text{Var}(\epsilon_i) = \sum_{i=1}^t 1 = t \end{aligned}$$

* Sum rule = a good reason for using the variance to measure spread; only works for independent distributions.

So typical displacement from the origin scales as:

$$\sigma = t^{1/2}$$

- 🔗 A non-trivial scaling law arises out of additive aggregation or accumulation.


PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
🔗 10 of 60

A few random random walks:



PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
🔗 8 of 60

Random walk basics:

Counting random walks:

- 🔗 Each **specific** random walk of length t appears with a chance $1/2^t$.
- 🔗 We'll be more interested in how many random walks end up at the same place.
- 🔗 Define $N(i, j, t)$ as # distinct walks that start at $x = i$ and end at $x = j$ after t time steps.
- 🔗 Random walk must displace by $+(j - i)$ after t steps.
- 🔗 [Insert question from assignment 3 ↗](#)

$$N(i, j, t) = \binom{t}{(t+j-i)/2}$$

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
🔗 13 of 60

Random walks:

Displacement after t steps:

$$x_t = \sum_{i=1}^t \epsilon_i$$

Expected displacement:

$$\langle x_t \rangle = \left\langle \sum_{i=1}^t \epsilon_i \right\rangle = \sum_{i=1}^t \langle \epsilon_i \rangle = 0$$

- 🔗 At any time step, we 'expect' our drunkard to be back at the pub.
- 🔗 Obviously fails for odd number of steps...
- 🔗 But as time goes on, the chance of our drunkard lurching back to the pub must diminish, right?

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
🔗 9 of 60

How does $P(x_t)$ behave for large t ?

- 🔗 Take time $t = 2n$ to help ourselves.
- 🔗 $x_{2n} \in \{0, \pm 2, \pm 4, \dots, \pm 2n\}$
- 🔗 x_{2n} is even so set $x_{2n} = 2k$.
- 🔗 Using our expression $N(i, j, t)$ with $i = 0, j = 2k$, and $t = 2n$, we have

$$\Pr(x_{2n} = 2k) \propto \binom{2n}{n+k}$$

- 🔗 For large n , the binomial deliciously approaches the Normal Distribution of Snoredom:

$$\Pr(x_t = x) \simeq \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}}$$

[Insert question from assignment 3 ↗](#)

- 🔗 The whole is different from the parts. **#nutritious**
- 🔗 See also: [Stable Distributions ↗](#)


PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
🔗 14 of 60

Universality is also not left-handed:



- This is Diffusion: the most essential kind of spreading (more later).
- View as Random Additive Growth Mechanism.

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
15 of 60

Random walks are even weirder than you might think...

- $\xi_{r,t}$ = the probability that by time step t , a random walk has crossed the origin r times.
- Think of a coin flip game with ten thousand tosses.
- If you are behind early on, what are the chances you will make a comeback?
- The most likely number of lead changes is... 0.
- In fact: $\xi_{0,t} > \xi_{1,t} > \xi_{2,t} > \dots$
- Even crazier:
The expected time between tied scores = ∞

See Feller, Intro to Probability Theory, Volume I [3]

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
18 of 60



PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
16 of 60

Applied knot theory:



"Designing tie knots by random walks"
Fink and Mao,
Nature, 398, 31-32, 1999. [4]





Figure 1 All diagrams are drawn in the frame of reference of the mirror image of the actual tie. a. The two ways of beginning a knot, L and R. For knots beginning with L, the tie must begin inside-out. b. The four-in-hand, denoted by the sequence L₁R₁L₂C₁T. c. A knot may be represented by a persistent random walk on a triangular lattice. The example shown is the four-in-hand, indicated by the walk 1111a

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
19 of 60



PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
17 of 60

Applied knot theory:

h	γ	γ/h	K(h, γ)	s	b	Name	Sequence
3	1	0.33	1	0	0		L ₁ R ₁ C ₁ T
4	1	0.25	1	-1	1	Four-in-hand	L ₁ R ₁ L ₂ C ₁ T
5	2	0.40	2	-1	0	Pratt knot	L ₁ C ₁ R ₁ L ₂ C ₁ T
6	2	0.33	4	0	0	Half-Windsor	L ₁ R ₁ C ₁ L ₂ R ₁ C ₁ T
7	2	0.29	6	-1	1		L ₁ R ₁ L ₂ C ₁ R ₁ L ₂ C ₁ T
7	3	0.43	4	0	1		L ₁ C ₁ R ₁ C ₁ L ₂ R ₁ C ₁ T
8	2	0.25	8	0	2		L ₁ R ₁ L ₂ C ₁ R ₁ L ₂ R ₁ C ₁ T
8	3	0.38	12	-1	0	Windsor	L ₁ C ₁ R ₁ L ₂ C ₁ R ₁ L ₂ C ₁ T
9	4	0.33	24	0	0		L ₁ R ₁ C ₁ L ₂ R ₁ C ₁ L ₂ R ₁ C ₁ T
9	4	0.44	8	-1	2		L ₁ C ₁ R ₁ C ₁ L ₂ C ₁ R ₁ L ₂ C ₁ T

Knots are characterized by half-winding number h , centre number γ , centre fraction γ/h , knots per class $K(h, \gamma)$, symmetry s , balance b , name and sequence.

- h = number of moves
- γ = number of center moves
- $K(h, \gamma) = \frac{2^{\gamma-1}(h-\gamma-2)}{\gamma-1}$
- $s = \sum_{i=1}^h x_i$ where $x = -1$ for L and +1 for R.
- $b = \frac{1}{2} \sum_{i=2}^{h-1} |\omega_i + \omega_{i-1}|$ where $\omega = \pm 1$ represents winding direction.

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



UNIVERSITY OF VERMONT
20 of 60

Random walks #crazytownbananapants

The problem of first return:

- What is the probability that a random walker in one dimension returns to the origin for the first time after t steps?
- Will our drunkard always return to the origin?
- What about higher dimensions?

Reasons for caring:

- We will find a power-law size distribution with an interesting exponent.
- Some physical structures may result from random walks.
- We'll start to see how different scalings relate to each other.

PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



22 of 60

Counting first returns:

Approach:

- Move to counting numbers of walks.
- Return to probability at end.
- Again, $N(i, j, t)$ is the # of possible walks between $x = i$ and $x = j$ taking t steps.
- Consider all paths starting at $x = 1$ and ending at $x = 1$ after $t = 2n - 2$ steps.
- Idea: If we can compute the number of walks that hit $x = 0$ at least once, then we can subtract this from the total number to find the ones that maintain $x \geq 1$.
- Call walks that drop below $x = 1$ excluded walks.
- We'll use a method of images to identify these excluded walks.


PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



25 of 60

For random walks in 1-d:



- A return to origin can only happen when $t = 2n$.
- In example above, returns occur at $t = 8, 10,$ and 14 .
- Call $P_{fr(2n)}$ the probability of first return at $t = 2n$.
- Probability calculation \equiv Counting problem (combinatorics/statistical mechanics).
- Idea: Transform first return problem into an easier return problem.


PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



23 of 60

Examples of excluded walks:



Key observation for excluded walks:


- For any path starting at $x=1$ that hits 0 , there is a unique matching path starting at $x=-1$.
- Matching path first mirrors and then tracks after first reaching $x=0$.
- # of t -step paths starting and ending at $x=1$ and hitting $x=0$ at least once = # of t -step paths starting at $x=-1$ and ending at $x=1 = N(-1, 1, t)$
- So $N_{\text{first return}}(2n) = N(1, 1, 2n-2) - N(-1, 1, 2n-2)$

PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



26 of 60



- Can assume drunkard first lurches to $x = 1$.
- Observe walk first returning at $t = 16$ stays at or above $x = 1$ for $1 \leq t \leq 15$ (dashed red line).
- Now want walks that can return many times to $x = 1$.
- $P_{fr}(2n) = 2 \cdot \frac{1}{2} Pr(x_t \geq 1, 1 \leq t \leq 2n-1, \text{ and } x_1 = x_{2n-1} = 1)$
- The $\frac{1}{2}$ accounts for $x_{2n} = 2$ instead of 0 .
- The 2 accounts for drunkards that first lurch to $x = -1$.

PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



24 of 60

Probability of first return:

Insert question from assignment 3:

- Find

$$N_{fr}(2n) \sim \frac{2^{2n-3/2}}{\sqrt{2\pi n^{3/2}}}$$

- Normalized number of paths gives probability.
- Total number of possible paths = 2^{2n} .

$$P_{fr}(2n) = \frac{1}{2^{2n}} N_{fr}(2n) \approx \frac{1}{2^{2n}} \frac{2^{2n-3/2}}{\sqrt{2\pi n^{3/2}}} = \frac{1}{\sqrt{2\pi}} (2n)^{-3/2} \propto t^{-3/2}$$

PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLIO
References



27 of 60

- We have $P(t) \propto t^{-3/2}$, $\gamma = 3/2$.
- Same scaling holds for continuous space/time walks.
- $P(t)$ is normalizable.
- Recurrence:** Random walker always returns to origin
- But mean, variance, and all higher moments are infinite. #totalmadness
- Even though walker must return, expect a long wait...
- One moral:** Repeated gambling against an infinitely wealthy opponent must lead to ruin.

Higher dimensions

- Walker in $d = 2$ dimensions must also return
- Walker may not return in $d \geq 3$ dimensions

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtzmark's Distribution
PLIPLD
References



UNIVERSITY OF VERMONT
28 of 60

Scheidegger networks

- Creates basins with random walk boundaries.
- Observe** that subtracting one random walk from another gives random walk with increments:

$$\epsilon_t = \begin{cases} +1 & \text{with probability } 1/4 \\ 0 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/4 \end{cases}$$

- Random walk with probabilistic pauses.
- Basin termination = first return random walk problem.
- Basin length ℓ distribution: $P(\ell) \propto \ell^{-3/2}$
- For real river networks, generalize to $P(\ell) \propto \ell^{-\gamma}$.

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtzmark's Distribution
PLIPLD
References



UNIVERSITY OF VERMONT
32 of 60

Random walks

On finite spaces:

- In any finite homogeneous space, a random walker will visit every site with equal probability
- Call this probability the **Invariant Density** of a dynamical system
- Non-trivial Invariant Densities arise in chaotic systems.

On networks:

- On networks, a random walker visits each node with frequency \propto node degree #groovy
- Equal probability still present: walkers traverse **edges** with equal frequency. #totallygroovy

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtzmark's Distribution
PLIPLD
References



UNIVERSITY OF VERMONT
29 of 60

Connections between exponents:

- For a basin of length ℓ , width $\propto \ell^{1/2}$
- Basin area $a \propto \ell \cdot \ell^{1/2} = \ell^{3/2}$
- Invert: $\ell \propto a^{2/3}$
- $d\ell \propto d(a^{2/3}) = 2/3 a^{-1/3} da$
- $\Pr(\text{basin area} = a)da = \Pr(\text{basin length} = \ell)d\ell$
- $\propto \ell^{-3/2} d\ell$
- $\propto (a^{2/3})^{-3/2} a^{-1/3} da$
- $= a^{-4/3} da$
- $= a^{-\tau} da$


PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtzmark's Distribution
PLIPLD
References



UNIVERSITY OF VERMONT
33 of 60

Scheidegger Networks [9, 2]



- Random directed network on triangular lattice.
- Toy model of real networks.
- 'Flow' is southeast or southwest with equal probability.

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtzmark's Distribution
PLIPLD
References



UNIVERSITY OF VERMONT
31 of 60

Connections between exponents:

- Both basin area and length obey power law distributions
- Observed for real river networks
- Reportedly: $1.3 < \tau < 1.5$ and $1.5 < \gamma < 2$

Generalize relationship between area and length:

- Hack's law [5]: $\ell \propto a^h$.

- For real, large networks $h \simeq 0.5$
- Smaller basins possibly $h > 1/2$ (see: allometry).
- Models exist with interesting values of h .
- Plan:** Redo calc with γ , τ , and h .

PoCS | @pocsvox
Power-Law
Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtzmark's Distribution
PLIPLD
References



UNIVERSITY OF VERMONT
34 of 60

Connections between exponents:

Given

$$\ell \propto a^h, P(a) \propto a^{-\tau}, \text{ and } P(\ell) \propto \ell^{-\gamma}$$

$d\ell \propto d(a^h) = ha^{h-1}da$

Find τ in terms of γ and h .

$$\begin{aligned} \Pr(\text{basin area} = a)da &= \Pr(\text{basin length} = \ell)d\ell \\ &\propto \ell^{-\gamma}d\ell \\ &\propto (a^h)^{-\gamma}a^{h-1}da \\ &= a^{-(1+h(\gamma-1))}da \end{aligned}$$

$$\tau = 1 + h(\gamma - 1)$$

Excellent example of the **Scaling Relations** found between exponents describing power laws for many systems.



More than randomness

Can generalize to Fractional Random Walks [7, 8, 6]

Levy flights, Fractional Brownian Motion

See Montroll and Shlesinger for example: [6]
"On $1/f$ noise and other distributions with long tails."
Proc. Natl. Acad. Sci., 1982.

In 1-d, standard deviation σ scales as

$$\sigma \sim t^\alpha$$

$\alpha = 1/2$ — diffusive

$\alpha > 1/2$ — superdiffusive

$\alpha < 1/2$ — subdiffusive

Extensive memory of path now matters...



Connections between exponents:

With more detailed description of network structure, $\tau = 1 + h(\gamma - 1)$ simplifies to: [1]

$$\tau = 2 - h$$

and


$$\gamma = 1/h$$

Only one exponent is independent (take h).

Simplifies system description.

Expect Scaling Relations where power laws are found.

Need only characterize Universality class with independent exponents.



Other First Returns or First Passage Times:

Failure:

A very simple model of failure/death: [11]

x_t = entity's 'health' at time t



Start with $x_0 > 0$.



Entity fails when x hits 0.

Streams

Dispersion of suspended sediments in streams.

Long times for clearing.





Now make transformation:

$$P_y(y)dy = P_x(x)dx$$

$$P_y(y)dy = P_x\left(\left(\frac{y}{c}\right)^{-1/\alpha}\right) \frac{c^{1/\alpha}}{\alpha} y^{-1-1/\alpha} dy$$

If $P_x(x) \rightarrow$ non-zero constant as $x \rightarrow 0$ then

$$P_y(y) \propto y^{-1-1/\alpha} \text{ as } y \rightarrow \infty.$$

If $P_x(x) \rightarrow x^\beta$ as $x \rightarrow 0$ then

$$P_y(y) \propto y^{-1-1/\alpha-\beta/\alpha} \text{ as } y \rightarrow \infty.$$



Variable Transformation

Understand power laws as arising from

1. Elementary distributions (e.g., exponentials).
2. Variables connected by power relationships.

- Random variable X with known distribution P_x
- Second random variable Y with $y = f(x)$.

$$P_Y(y)dy = \sum_{x|f(x)=y} P_X(x)dx = \sum_{y|f(x)=y} P_X(f^{-1}(y)) |f'(f^{-1}(y))| dy$$

Often easier to do by hand...

General Example

- Assume relationship between x and y is 1-1.
- Power-law relationship between variables:
 $y = cx^{-\alpha}$, $\alpha > 0$
- Look at y large and x small

$$dy = d(cx^{-\alpha})$$

$$= c(-\alpha)x^{-\alpha-1}dx$$

$$\text{invert: } dx = \frac{-1}{c\alpha} x^{\alpha+1} dy$$

$$dx = \frac{-1}{c\alpha} \left(\frac{y}{c}\right)^{-(\alpha+1)/\alpha} dy$$

$$dx = \frac{-c^{1/\alpha}}{\alpha} y^{-1-1/\alpha} dy$$



Example

Exponential distribution

Given $P_x(x) = \frac{1}{\lambda} e^{-x/\lambda}$ and $y = cx^{-\alpha}$, then


$$P(y) \propto y^{-1-1/\alpha} + O(y^{-1-2/\alpha})$$

- Exponentials arise from randomness (easy)...
- More later when we cover robustness.



Gravity

- Select a random point in the universe \vec{x}
- Measure the force of gravity $F(\vec{x})$
- Observe that $P_F(F) \sim F^{-5/2}$.



Matter is concentrated in stars: [10]

- F is distributed unevenly
- Probability of being a distance r from a single star at $\vec{x} = \vec{0}$:

$$P_r(r)dr \propto r^2 dr$$

- Assume stars are distributed randomly in space (oops?)
- Assume only one star has significant effect at \vec{x} .
- Law of gravity:

$$F \propto r^{-2}$$

- invert:


$$r \propto F^{-1/2}$$

- Also invert:

$$dF \propto d(r^{-2}) \propto r^{-3} dr \rightarrow dr \propto r^3 dF \propto F^{-3/2} dF.$$



□ Todo: Build Dalek army.



Transformation:

Using $r \propto F^{-1/2}$, $dr \propto F^{-3/2} dF$, and $P_r(r) \propto r^2$

$$\begin{aligned} P_F(F)dF &= P_r(r)dr \\ &\propto P_r(\text{const} \times F^{-1/2})F^{-3/2}dF \\ &\propto (F^{-1/2})^2 F^{-3/2}dF \\ &= F^{-1-3/2}dF \\ &= F^{-5/2}dF. \end{aligned}$$



Extreme Caution!

- PLIPLD = Power law in, power law out
- Explain a power law as resulting from another unexplained power law.
- Yet another homunculus argument?
- Don't do this!!! (slap, slap)
- MIWO = Mild in, Wild out is fine.
- In general: We need mechanisms!



Gravity:

$$\begin{aligned} P_F(F) &= F^{-5/2} dF \\ \gamma &= 5/2 \end{aligned}$$

- Mean is finite.
- Variance = ∞ .
- A wild distribution.
- Upshot: Random sampling of space usually safe but can end badly...



References I

- P. S. Dodds and D. H. Rothman. Unified view of scaling laws for river networks. *Physical Review E*, 59(5):4865-4877, 1999. [pdf](#)
- P. S. Dodds and D. H. Rothman. Scaling, universality, and geomorphology. *Annu. Rev. Earth Planet. Sci.*, 28:571-610, 2000. [pdf](#)
- W. Feller. *An Introduction to Probability Theory and Its Applications*, volume I. John Wiley & Sons, New York, third edition, 1968.
- T. M. Fink and Y. Mao. Designing tie knots by random walks. *Nature*, 398:31-32, 1999. [pdf](#)



References II

- [5] J. T. Hack.
Studies of longitudinal stream profiles in Virginia and Maryland.
[United States Geological Survey Professional Paper, 294-B:45-97, 1957. pdf](#)
- [6] E. W. Montroll and M. F. Shlesinger.
On the wonderful world of random walks, volume XI of *Studies in statistical mechanics*, chapter 1, pages 1-121.
New-Holland, New York, 1984.
- [7] E. W. Montroll and M. W. Shlesinger.
On $1/f$ noise and other distributions with long tails.
[Proc. Natl. Acad. Sci., 79:3380-3383, 1982. pdf](#)

PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLI
[References](#)



59 of 60

References III

- [8] E. W. Montroll and M. W. Shlesinger.
Maximum entropy formalism, fractals, scaling phenomena, and $1/f$ noise: a tale of tails.
[J. Stat. Phys., 32:209-230, 1983.](#)
- [9] A. E. Scheidegger.
The algebra of stream-order numbers.
[United States Geological Survey Professional Paper, 525-B:B187-B189, 1967. pdf](#)
- [10] D. Sornette.
Critical Phenomena in Natural Sciences.
Springer-Verlag, Berlin, 1st edition, 2003.
- [11] J. S. Weitz and H. B. Fraser.
Explaining mortality rate plateaus.
[Proc. Natl. Acad. Sci., 98:15383-15386, 2001. pdf](#)

PoCS | @pocsvox
Power-Law Mechanisms, Pt. 1

Random Walks
The First Return Problem
Examples
Variable transformation
Basics
Holtsmark's Distribution
PLIPLI
[References](#)



60 of 60