



What's  
The  
Story?

**Principles of Complex Systems, CSYS/MATH 300**  
**University of Vermont, Fall 2017**  
**Assignment 1 • code name: Bizarre Gardening Accident**

**Dispersed:** Tuesday, August 29, 2017.

**Due:** 11:59 pm, Friday, September 8, 2017.

*Some useful reminders:*

**Deliverator:** Peter Dodds

**Office:** Farrell Hall, second floor, Trinity Campus

**E-mail:** peter.dodds+pocs@uvm.edu

**Office hours:** 1:15 pm to 2:30 pm on Tuesdays and Thursdays, 10:00 am to 11:30 am on Wednesdays

**Course website:** <http://www.uvm.edu/pdodds/teaching/courses/2017-08UVM-300>

All parts are worth 3 points unless marked otherwise. Please show all your workingses clearly and list the names of others with whom you collaborated.

Graduate students are requested to use  $\LaTeX$  (or related  $\TeX$  variant).

**Email submission:** PDF only! Please name your file as follows (where the number is to be padded by a 0 if less than 10 and names are all lowercase):

CSYS300assignment%02d\$firstname-\$lastname.pdf as in

CSYS300assignment06michael-palin.pdf

1. Use a back-of-an-envelope scaling argument to show that maximal rowing speed  $V$  increases as the number of oarspeople  $n$  as  $V \propto N^{1/9}$ .


Assume the following:

- (a) Rowing shells are geometrically similar (isometric). The table below taken from McMahon and Bonner [1] shows that shell width is roughly proportional to shell length  $\ell$ .

Shell dimensions and performances.

No. of oarsmen	Modifying description	Length, $l$ (m)	Beam, $b$ (m)	$l/b$	Boat mass per oarsman (kg)	Time for 2000 m (min)			
						I	II	III	IV
8	Heavyweight	18.28	0.610	30.0	14.7	5.87	5.92	5.82	5.73
8	Lightweight	18.28	0.598	30.6	14.7				
4	With coxswain	12.80	0.574	22.3	18.1				
4	Without coxswain	11.75	0.574	21.0	18.1	6.33	6.42	6.48	6.13
2	Double scull	9.76	0.381	25.6	13.6				
2	Pair-oared shell	9.76	0.356	27.4	13.6	6.87	6.92	6.95	6.77
1	Single scull	7.93	0.293	27.0	16.3	7.16	7.25	7.28	7.17

- (b) The resistance encountered by a shell is due largely to drag on its wetted surface.
  - (c) Drag is proportional to the product of the square of the shell's speed ( $V^2$ ) and the area of the wetted surface ( $\propto \ell^2$  due to shell isometry).
  - (d) Power  $\propto$  drag force  $\times$  speed (in symbols:  $P \propto D_f \times V$ ).
  - (e) Volume displacement of water by a shell is proportional to the number of oarspeople  $N$  (i.e., the team's combined weight).
  - (f) Assume the depth of water displacement by the shell grows isometrically with boat length  $\ell$ .
  - (g) Power is proportional to the number of oarspeople  $N$ .
2. Find the modern day world record times for 2000 metre races and see if this scaling still holds up. Of course, our relationship is approximate as we have neglected numerous factors, the range is extremely small (1–8 oarspeople), and the scaling is very weak (1/9). But see what you can find. The figure below shows data from McMahon and Bonner.



3. Check current weight lifting records for the snatch, clean and jerk, and the total for scaling with body mass (three regressions).  
For weight classes, take the upper limit for the mass of the lifter.
- (a) Does  $2/3$  scaling hold up?
  - (b) Normalized by the appropriate scaling, who holds the overall, rescaled world record?
4. Finish the calculation for the platypus on a pendulum problem so show that a simple pendulum's period  $\tau$  is indeed proportional to  $\sqrt{\ell/g}$ .

Basic plan from lectures: Create a matrix  $\mathbf{A}$  where  $ij$ th entry is the power of dimension  $i$  in the  $j$ th variable, and solve by row reduction to find basis null vectors.

In lectures, we arrived at:

$$\mathbf{A}\vec{x} = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

You only have to take a few steps from here.

**Help—Buckingham  $\pi$  theorem** (runs from 56:14 to 1:12:08, length 15:54):  
<http://www.youtube.com/v/TX-QmpPbWgE?rel=0&start=3374&end=4328>

5. Show that the maximum speed of animals  $V_{\max}$  is proportional to their length  $L$  [2]. Here are five dimensionful parameters:
  - $V_{\max}$ , maximum speed.
  - $\ell$ , animal length.
  - $\rho$ , organismal density.
  - $\sigma$ , maximum applied force per unit area of tissue.
  - $b$ , maximum metabolic rate per unit mass ( $b$  has the dimensions of power per unit mass).

And here are the three dimensions:  $L$ ,  $M$ , and  $T$ .

Use a back-of-the-envelope calculation to express  $V_{\max}/\ell$  in terms of  $\rho$ ,  $\sigma$ , and  $b$ .

Note: It's argued in [2] that these latter three parameters vary little across all organisms (we're mostly thinking about running organisms here), and so finding  $V_{\max}/\ell$  as a function of them indicates that  $V_{\max}/\ell$  is also roughly constant.

6. Use the Buckingham  $\pi$  theorem to reproduce G. I. Taylor's finding the energy of an atom bomb  $E$  is related to the density of air  $\rho$  and the radius of the blast wave  $R$  at time  $t$ :

$$E = \text{constant} \times \rho R^5 / t^2.$$

In constructing the matrix, order parameters as  $E$ ,  $\rho$ ,  $R$ , and  $t$  and dimensions as  $L$ ,  $T$ , and  $M$ .

7. Surface area of allometrically growing Minecraft organisms:

Let's consider animals as parallelepipeds (e.g., the well known box cow), with dimensions  $L_1$ ,  $L_2$ , and  $L_3$  and volume  $V = L_1 \times L_2 \times L_3$ .

As we vary in scale of organism, let's assume the lengths scale with volume as  $L_i = c_i^{-1} V^{\gamma_i}$  where the exponents satisfy  $\gamma_1 + \gamma_2 + \gamma_3 = 1$  and the  $c_i$  are prefactors such that  $c_1 \times c_2 \times c_3 = 1$ . Let's also arrange our organisms so that  $\gamma_1 \leq \gamma_2 \leq \gamma_3$ .

- (a) Show that the scalings  $L_i = c_i^{-1} V^{\gamma_i}$  mean that indeed  $L_1 \times L_2 \times L_3 = V$ .
- (b) Write down the  $\gamma_i$  corresponding to isometric scaling.
- (c) Calculate the surface area  $S$  of our imaginary beings.
- (d) Show how  $S$  behaves as  $V$  becomes large (i.e., which term(s) dominate).
- (e) Which sets of  $\gamma_i$  give the fastest and slowest possible scaling of  $S$  as a function of  $V$ ?

Note: surface area is a big deal for organisms and this calculation will matter later in PoCS and/or CocoNuTs.

## References

- [1] T. A. McMahon and J. T. Bonner. *On Size and Life*. Scientific American Library, New York, 1983.
- [2] N. Meyer-Vernet and J.-P. Rospars. How fast do living organisms move: Maximum speeds from bacteria to elephants and whales. *American Journal of Physics*, pages 719–722, 2015. [pdf](#) 