

Branching Networks II

Last updated: 2023/01/26, 11:37:55 EST

Principles of Complex Systems, Vols. 1, 2, & 3D
CSYS/MATH 300, 303, & 394, 2022-2023 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
Santa Fe Institute | University of Vermont



Licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License.

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

Piracy on the high χ 's:

More: [How river networks move across a landscape](#) (Science Daily)



1 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References



2 of 84

Outline

Horton ⇄ Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References

Piracy on the high χ 's:



"Dynamic Reorganization of River Basins" Willett et al., Science, **343**, 1248765, 2014. [21]



$$\frac{\partial z(x, t)}{\partial t} = U - K A^m \left| \frac{\partial z(x, t)}{\partial x} \right|^n$$

$$z(x) = z_b + \left(\frac{U}{K A_0^m} \right)^{1/n} \chi$$

$$\chi = \int_{x_b}^x \left(\frac{A_0}{A(x')} \right)^{m/n} dx'$$

Can Horton and Tokunaga be happy?

Horton and Tokunaga seem different:

- In terms of network architecture, Horton's laws appear to contain less detailed information than Tokunaga's law.
- Oddly, Horton's laws have **four** parameters and Tokunaga has **two** parameters.
- $R_n, R_a, R_\ell,$ and R_s **versus** T_1 and R_T . One simple redundancy: $R_\ell = R_s$.
[Insert question from assignment 15](#)
- To make a connection, clearest approach is to start with Tokunaga's law ...
- Known result: Tokunaga \rightarrow Horton [18, 19, 20, 9, 2]



8 of 84

Let us make them happy

We need one more ingredient:

Space-fillingness

- A network is **space-filling** if the average distance between adjacent streams is roughly constant.
- Reasonable for river and cardiovascular networks
- For river networks:
Drainage density ρ_{dd} = inverse of typical distance between channels in a landscape.
- In terms of basin characteristics:

$$\rho_{dd} \simeq \frac{\sum \text{stream segment lengths}}{\text{basin area}} = \frac{\sum_{\omega=1}^{\Omega} n_{\omega} \bar{s}_{\omega}}{a_{\Omega}}$$




9 of 84

More with the happy-making thing

Start with Tokunaga's law: $T_k = T_1 R_T^{k-1}$

- Start looking for Horton's stream number law: $n_{\omega}/n_{\omega+1} = R_n$.
- Estimate n_{ω} , the number of streams of order ω in terms of other $n_{\omega'}, \omega' > \omega$.
- Observe that each stream of order ω terminates by either:
 - Running into another stream of order ω and generating a stream of order $\omega + 1$...
▶ $2n_{\omega+1}$ streams of order ω do this
 - Running into and being absorbed by a stream of higher order $\omega' > \omega$...
▶ $n_{\omega'} T_{\omega'-\omega}$ streams of order ω do this



5 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

More with the happy-making thing

Putting things together:



$$n_{\omega} = \underbrace{2n_{\omega+1}}_{\text{generation}} + \sum_{\omega'=\omega+1}^{\Omega} \underbrace{T_{\omega'-\omega} n_{\omega'}}_{\text{absorption}}$$

- Use Tokunaga's law and manipulate expression to find Horton's law for stream numbers follows and hence obtain R_n .
- [Insert question from assignment 16](#)
- Solution:

$$R_n = \frac{(2 + R_T + T_1) \pm \sqrt{(2 + R_T + T_1)^2 - 8R_T}}{2}$$

(The larger value is the one we want.)



8 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

Finding other Horton ratios

Connect Tokunaga to R_s

- Now use uniform drainage density ρ_{dd} .
- Assume side streams are roughly separated by distance $1/\rho_{dd}$.
- For an order ω **stream segment**, expected length is

$$\bar{s}_{\omega} \simeq \rho_{dd}^{-1} \left(1 + \sum_{k=1}^{\omega-1} T_k \right)$$

- Substitute in Tokunaga's law $T_k = T_1 R_T^{k-1}$:

$$\bar{s}_{\omega} \simeq \rho_{dd}^{-1} \left(1 + T_1 \sum_{k=1}^{\omega-1} R_T^{k-1} \right) \propto R_T^{\omega}$$



9 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References



10 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References



11 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References



12 of 84

Horton and Tokunaga are happy

Altogether then:

$$\Rightarrow \bar{s}_\omega / \bar{s}_{\omega-1} = R_T \Rightarrow R_s = R_T$$

Recall $R_\ell = R_s$ so

$$R_\ell = R_s = R_T$$

And from before:

$$R_n = \frac{(2 + R_T + T_1) + \sqrt{(2 + R_T + T_1)^2 - 8R_T}}{2}$$

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

13 of 84

Horton and Tokunaga are happy

Some observations:

- R_n and R_ℓ depend on T_1 and R_T .
- Seems that R_a must as well ...
- Suggests Horton's laws must contain some redundancy
- We'll in fact see that $R_a = R_n$.
- Also: Both Tokunaga's law and Horton's laws can be generalized to relationships between non-trivial statistical distributions. [3, 4]

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

14 of 84

Horton and Tokunaga are happy

The other way round

Note: We can invert the expressions for R_n and R_ℓ to find Tokunaga's parameters in terms of Horton's parameters.

$$R_T = R_\ell,$$

$$T_1 = R_n - R_\ell - 2 + 2R_\ell/R_n.$$

Suggests we should be able to argue that Horton's laws imply Tokunaga's laws (if drainage density is uniform) ...

PoCS
@pocsvox
Branching
Networks II


Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

15 of 84

Horton and Tokunaga are friends

From Horton to Tokunaga [2]



Assume Horton's laws hold for number and length

Start with picture showing an order ω stream and order $\omega - 1$ generating and side streams.

Scale up by a factor of R_ℓ , orders increment to $\omega + 1$ and ω .

Maintain drainage density by adding new order $\omega - 1$ streams

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

16 of 84

Horton and Tokunaga are friends

...and in detail:

- Must retain same drainage density.
- Add an extra $(R_\ell - 1)$ first order streams for each original tributary.
- Since by definition, an order $\omega + 1$ stream segment has T_ω order 1 side streams, we have:

$$T_k = (R_\ell - 1) \left(1 + \sum_{i=1}^{k-1} T_i \right).$$

For large ω , Tokunaga's law is the solution—let's check ...

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

17 of 84

Horton and Tokunaga are friends

Just checking:

Substitute Tokunaga's law $T_i = T_1 R_T^{i-1} = T_1 R_\ell^{i-1}$ into

$$T_k = (R_\ell - 1) \left(1 + \sum_{i=1}^{k-1} T_i \right)$$

$$T_k = (R_\ell - 1) \left(1 + \sum_{i=1}^{k-1} T_1 R_\ell^{i-1} \right)$$

$$= (R_\ell - 1) \left(1 + T_1 \frac{R_\ell^{k-1} - 1}{R_\ell - 1} \right)$$

$$\approx (R_\ell - 1) T_1 \frac{R_\ell^{k-1}}{R_\ell - 1} = T_1 R_\ell^{k-1} \quad \dots \text{yep.}$$


PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

18 of 84

Horton's laws of area and number:



In bottom plots, stream number graph has been flipped vertically.

Highly suggestive that $R_n \equiv R_a$...

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

19 of 84

Measuring Horton ratios is tricky:

- How robust are our estimates of ratios?
- Rule of thumb: discard data for two smallest and two largest orders.

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

20 of 84

Mississippi:

ω range	R_n	R_a	R_ℓ	R_s	R_a/R_n
[2, 3]	5.27	5.26	2.48	2.30	1.00
[2, 5]	4.86	4.96	2.42	2.31	1.02
[2, 7]	4.77	4.88	2.40	2.31	1.02
[3, 4]	4.72	4.91	2.41	2.34	1.04
[3, 6]	4.70	4.83	2.40	2.35	1.03
[3, 8]	4.60	4.79	2.38	2.34	1.04
[4, 6]	4.69	4.81	2.40	2.36	1.02
[4, 8]	4.57	4.77	2.38	2.34	1.05
[5, 7]	4.68	4.83	2.36	2.29	1.03
[6, 7]	4.63	4.76	2.30	2.16	1.03
[7, 8]	4.16	4.67	2.41	2.56	1.12
mean μ	4.69	4.85	2.40	2.33	1.04
std dev σ	0.21	0.13	0.04	0.07	0.03
σ/μ	0.045	0.027	0.015	0.031	0.024

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

21 of 84

ω range	R_n	R_a	R_ℓ	R_s	R_a/R_n
[2, 3]	4.78	4.71	2.47	2.08	0.99
[2, 5]	4.55	4.58	2.32	2.12	1.01
[2, 7]	4.42	4.53	2.24	2.10	1.02
[3, 5]	4.45	4.52	2.26	2.14	1.01
[3, 7]	4.35	4.49	2.20	2.10	1.03
[4, 6]	4.38	4.54	2.22	2.18	1.03
[5, 6]	4.38	4.62	2.22	2.21	1.06
[6, 7]	4.08	4.27	2.05	1.83	1.05
mean μ	4.42	4.53	2.25	2.10	1.02
std dev σ	0.17	0.10	0.10	0.09	0.02
σ/μ	0.038	0.023	0.045	0.042	0.019

Reducing Horton's laws:

Not quite:

- ...But this only a rough argument as Horton's laws do not imply a strict hierarchy
- Need to account for sidebranching.
- Insert question from assignment 16

Reducing Horton's laws:

Rough first effort to show $R_n \equiv R_a$:

- $a_\Omega \propto$ sum of all stream segment lengths in a order Ω basin (assuming uniform drainage density)
- So:

$$a_\Omega \approx \sum_{\omega=1}^{\Omega} n_\omega \bar{s}_\omega / \rho_{dd}$$

$$\propto \sum_{\omega=1}^{\Omega} \underbrace{R_n^{\Omega-\omega}}_{n_\omega} \cdot \underbrace{\bar{s}_1}_{\bar{s}_1} \cdot \underbrace{R_s^{\omega-1}}_{\bar{s}_\omega}$$

$$= \frac{R_n^\Omega}{R_s} \bar{s}_1 \sum_{\omega=1}^{\Omega} \left(\frac{R_s}{R_n} \right)^\omega$$

Equipartitioning:

Intriguing division of area:

- Observe: Combined area of basins of order ω independent of ω .
- Not obvious: basins of low orders not necessarily contained in basin on higher orders.
- Story:
- Reason:

$$R_n \equiv R_a \Rightarrow n_\omega \bar{a}_\omega = \text{const}$$

$$n_\omega \propto (R_n)^{-\omega}$$

$$\bar{a}_\omega \propto (R_a)^\omega \propto n_\omega^{-1}$$

Reducing Horton's laws:

Continued ...

$$a_\Omega \propto \frac{R_n^\Omega}{R_s} \bar{s}_1 \sum_{\omega=1}^{\Omega} \left(\frac{R_s}{R_n} \right)^\omega$$

$$= \frac{R_n^\Omega}{R_s} \bar{s}_1 \frac{R_s}{R_n} \frac{1 - (R_s/R_n)^\Omega}{1 - (R_s/R_n)}$$


$$\sim R_n^{\Omega-1} \bar{s}_1 \frac{1}{1 - (R_s/R_n)} \text{ as } \Omega \nearrow$$

- So, a_Ω is growing like R_n^Ω and therefore:

$$R_n \equiv R_a$$

Equipartitioning:

Some examples:



Neural Reboot: Fwoompf

Scaling laws

The story so far:

- Natural branching networks are **hierarchical, self-similar** structures
- Hierarchy is **mixed**
- Tokunaga's law describes detailed architecture: $T_k = T_1 R_T^{k-1}$.
- We have connected Tokunaga's and Horton's laws
- Only two Horton laws are independent ($R_n = R_a$)
- Only **two** parameters are **independent**: $(T_1, R_T) \Leftrightarrow (R_n, R_s)$

Scaling laws

A little further ...

- Ignore stream ordering for the moment
- Pick a random location on a branching network p .
- Each point p is associated with a basin and a longest stream length
- Q:** What is probability that the p 's drainage basin has area a ? $P(a) \propto a^{-\tau}$ for large a
- Q:** What is probability that the longest stream from p has length ℓ ? $P(\ell) \propto \ell^{-\gamma}$ for large ℓ
- Roughly observed: $1.3 \lesssim \tau \lesssim 1.5$ and $1.7 \lesssim \gamma \lesssim 2.0$

Scaling laws

Probability distributions with power-law decays

- We see them everywhere:
 - Earthquake magnitudes (Gutenberg-Richter law)
 - City sizes (Zipf's law)
 - Word frequency (Zipf's law)^[22]
 - Wealth (maybe not—at least heavy tailed)
 - Statistical mechanics (phase transitions)^[5]
- A big part of the story of complex systems
- Arise from **mechanisms**: growth, randomness, optimization, ...
- Our task is always to illuminate the mechanism ...

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 31 of 84

Scaling laws

Connecting exponents

- We have the detailed picture of branching networks (Tokunaga and Horton)
- Plan: Derive $P(a) \propto a^{-\tau}$ and $P(\ell) \propto \ell^{-\gamma}$ starting with Tokunaga/Horton story^[17, 1, 2]
- Let's work on $P(\ell)$...
- Our first fudge: assume Horton's laws hold throughout a basin of order Ω .
- (We know they deviate from strict laws for low ω and high ω but not too much.)
- Next: place stick between teeth. Bite stick. Proceed.

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 32 of 84

Scaling laws

Finding γ :

- Often useful to work with **cumulative distributions**, especially when dealing with power-law distributions.
- The complementary cumulative distribution turns out to be most useful:

$$P_{>}(\ell_*) = P(\ell > \ell_*) = \int_{\ell=\ell_*}^{\ell_{\max}} P(\ell) d\ell$$

$$P_{>}(\ell_*) = 1 - P(\ell < \ell_*)$$

- Also known as the exceedance probability.

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 33 of 84

Scaling laws

Finding γ :

- The connection between $P(x)$ and $P_{>}(x)$ when $P(x)$ has a power law tail is simple:
- Given $P(\ell) \sim \ell^{-\gamma}$ large ℓ then for large enough ℓ_*

$$P_{>}(\ell_*) = \int_{\ell=\ell_*}^{\ell_{\max}} P(\ell) d\ell$$

$$\sim \int_{\ell=\ell_*}^{\ell_{\max}} \ell^{-\gamma} d\ell$$

$$= \frac{\ell^{-(\gamma-1)}}{-(\gamma-1)} \Big|_{\ell=\ell_*}^{\ell_{\max}}$$

$$\propto \ell_*^{-(\gamma-1)} \text{ for } \ell_{\max} \gg \ell_*$$

Scaling laws

Finding γ :

- Aim:** determine probability of randomly choosing a point on a network with main stream length $> \ell_*$
- Assume some spatial sampling resolution Δ
- Landscape is broken up into grid of $\Delta \times \Delta$ sites
- Approximate $P_{>}(\ell_*)$ as

$$P_{>}(\ell_*) = \frac{N_{>}(\ell_*; \Delta)}{N_{>}(0; \Delta)}$$

where $N_{>}(\ell_*; \Delta)$ is the number of sites with main stream length $> \ell_*$.

- Use Horton's law of stream segments: $\bar{s}_{\omega} / \bar{s}_{\omega-1} = R_s$...

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 34 of 84

Scaling laws

Finding γ :

- We are here:

$$P_{>}(\bar{\ell}_{\omega}) \propto \sum_{\omega'=\omega+1}^{\Omega} (1 \cdot R_n^{\Omega-\omega'}) (\bar{s}_1 \cdot R_s^{\omega'-1})$$

- Cleaning up irrelevant constants:

$$P_{>}(\bar{\ell}_{\omega}) \propto \sum_{\omega'=\omega+1}^{\Omega} \left(\frac{R_s}{R_n} \right)^{\omega'}$$

- Change summation order by substituting $\omega'' = \Omega - \omega'$.
- Sum is now from $\omega'' = 0$ to $\omega'' = \Omega - \omega - 1$ (equivalent to $\omega' = \Omega$ down to $\omega' = \omega + 1$)

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 35 of 84

Scaling laws

Finding γ :

-

$$P_{>}(\bar{\ell}_{\omega}) \propto \sum_{\omega''=0}^{\Omega-\omega-1} \left(\frac{R_s}{R_n} \right)^{\Omega-\omega''} \propto \sum_{\omega''=0}^{\Omega-\omega-1} \left(\frac{R_n}{R_s} \right)^{\omega''}$$

- Since $R_n > R_s$ and $1 \ll \omega \ll \Omega$,

$$P_{>}(\bar{\ell}_{\omega}) \propto \left(\frac{R_n}{R_s} \right)^{\Omega-\omega} \propto \left(\frac{R_n}{R_s} \right)^{-\omega}$$

again using $\sum_{i=0}^{n-1} a^i = (a^n - 1)/(a - 1)$

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 36 of 84

Scaling laws

Finding γ :

- Nearly there:

$$P_{>}(\bar{\ell}_{\omega}) \propto \left(\frac{R_n}{R_s} \right)^{-\omega} = e^{-\omega \ln(R_n/R_s)}$$

- Need to express right hand side in terms of $\bar{\ell}_{\omega}$.
- Recall that $\bar{\ell}_{\omega} \simeq \bar{\ell}_1 R_{\ell}^{\omega-1}$.

$$\bar{\ell}_{\omega} \propto R_{\ell}^{\omega} = R_s^{\omega} = e^{\omega \ln R_s}$$

📄 36 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 37 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 38 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ⇄
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

📄 39 of 84

Scaling laws

Finding γ :

Therefore:

$$P_{>}(\bar{\ell}_\omega) \propto e^{-\omega \ln(R_n/R_s)} = (e^{\omega \ln R_s})^{-\ln(R_n/R_s)/\ln R_s}$$

$$\propto \bar{\ell}_\omega^{-\ln(R_n/R_s)/\ln R_s}$$

$$= \bar{\ell}_\omega^{-(\ln R_n - \ln R_s)/\ln R_s}$$

$$= \bar{\ell}_\omega^{-\ln R_n/\ln R_s + 1}$$

$$= \bar{\ell}_\omega^{-\gamma + 1}$$

Scaling laws

Finding γ :

And so we have:

$$\gamma = \ln R_n / \ln R_s$$

Proceeding in a similar fashion, we can show

$$\tau = 2 - \ln R_s / \ln R_n = 2 - 1/\gamma$$

Insert question from assignment 16

Such connections between exponents are called **scaling relations**

Let's connect to one last relationship: Hack's law

Scaling laws

Hack's law: [6]

$$\ell \propto a^h$$

Typically observed that $0.5 \leq h \leq 0.7$.

Use Horton laws to connect h to Horton ratios:

$$\bar{\ell}_\omega \propto R_s^\omega \text{ and } \bar{a}_\omega \propto R_n^\omega$$

Observe:

$$\bar{\ell}_\omega \propto e^{\omega \ln R_s} \propto (e^{\omega \ln R_n})^{\ln R_s / \ln R_n}$$

$$\propto (R_n^\omega)^{\ln R_s / \ln R_n} \propto \bar{a}_\omega^{\ln R_s / \ln R_n} \Rightarrow h = \ln R_s / \ln R_n$$

PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



40 of 84

PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



41 of 84

PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



42 of 84

We mentioned there were a good number of 'laws': [2]

Relation:	Name or description:
$T_k = T_1 (R_T)^{k-1}$	Tokunaga's law
$\ell \sim L^d$	self-affinity of single channels
$n_\omega / n_{\omega+1} = R_n$	Horton's law of stream numbers
$\ell_{\omega+1} / \ell_\omega = R_\ell$	Horton's law of main stream lengths
$\bar{a}_{\omega+1} / \bar{a}_\omega = R_a$	Horton's law of basin areas
$\bar{s}_{\omega+1} / \bar{s}_\omega = R_s$	Horton's law of stream segment lengths
$L_\perp \sim L^H$	scaling of basin widths
$P(a) \sim a^{-\tau}$	probability of basin areas
$P(\ell) \sim \ell^{-\gamma}$	probability of stream lengths
$\ell \sim a^h$	Hack's law
$a \sim L^D$	scaling of basin areas
$\Lambda \sim a^\beta$	Langbein's law
$\lambda \sim L^\varphi$	variation of Langbein's law

PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



43 of 84

PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



44 of 84

PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



45 of 84

Connecting exponents


Only 3 parameters are independent: e.g., take d , R_n , and R_s

relation:	scaling relation/parameter: [2]
$\ell \sim L^d$	d
$T_k = T_1 (R_T)^{k-1}$	$T_1 = R_n - R_s - 2 + 2R_s/R_n$ $R_T = R_s$
$n_\omega / n_{\omega+1} = R_n$	R_n
$\bar{a}_{\omega+1} / \bar{a}_\omega = R_a$	$R_a = R_n$
$\ell_{\omega+1} / \ell_\omega = R_\ell$	$R_\ell = R_s$
$\ell \sim a^h$	$h = \ln R_s / \ln R_n$
$a \sim L^D$	$D = d/h$
$L_\perp \sim L^H$	$H = d/h - 1$
$P(a) \sim a^{-\tau}$	$\tau = 2 - h$
$P(\ell) \sim \ell^{-\gamma}$	$\gamma = 1/h$
$\Lambda \sim a^\beta$	$\beta = 1 + h$
$\lambda \sim L^\varphi$	$\varphi = d$

A toy model—Scheidegger's model

Random walk basins:

Boundaries of basins are random walks



PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



46 of 84


PoCS @pocsvox Branching Networks II

Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



47 of 84

Scheidegger's model



PoCS @pocsvox Branching Networks II


Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



48 of 84

Scheidegger's model

Directed random networks [11, 12]



$$P(\searrow) = P(\swarrow) = 1/2$$

Functional form of all scaling laws exhibited by exponents differ from real world [15, 16, 14]

Useful and interesting test case

Scheidegger's model

Prob for first return of a random walk in (1+1) dimensions (from CSYS/MATH 300):

$$P(n) \sim \frac{1}{2\sqrt{\pi}} n^{-3/2}$$

and so $P(\ell) \propto \ell^{-3/2}$.

Typical area for a walk of length n is $\propto n^{3/2}$:

$$\ell \propto a^{2/3}$$

Find $\tau = 4/3$, $h = 2/3$, $\gamma = 3/2$, $d = 1$.

Note $\tau = 2 - h$ and $\gamma = 1/h$.

R_n and R_ℓ have not been derived analytically.

PoCS @pocsvox Branching Networks II


Horton Tokunaga Reducing Horton Scaling relations Fluctuations Models Nutshell References



48 of 84

Equipartitioning reexamined:

Recall this story:




PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 49 of 84

A toy model—Scheidegger's model

Directed random networks [11, 12]



$$P(\searrow) = P(\swarrow) = 1/2$$

Flow is directed downwards

PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 52 of 84

Generalizing Horton's laws

Comparison of predicted versus measured main stream lengths for large scale river networks (in 10^3 km):

basin:	$\bar{\ell}_\Omega$	$\bar{\ell}'_\Omega$	σ_ℓ	$\bar{\ell}_\Omega/\bar{\ell}'_\Omega$	$\sigma_\ell/\bar{\ell}'_\Omega$
Mississippi	4.92	11.10	5.60	0.44	0.51
Amazon	5.75	9.18	6.85	0.63	0.75
Nile	6.49	2.66	2.20	2.44	0.83
Congo	5.07	10.13	5.75	0.50	0.57
Kansas	1.07	2.37	1.74	0.45	0.73

	a_{Ω}	\bar{a}'_Ω	σ_a	$a_{\Omega}/\bar{a}'_\Omega$	σ_a/\bar{a}'_Ω
Mississippi	2.74	7.55	5.58	0.36	0.74
Amazon	5.40	9.07	8.04	0.60	0.89
Nile	3.08	0.96	0.79	3.19	0.82
Congo	3.70	10.09	8.28	0.37	0.82
Kansas	0.14	0.49	0.42	0.28	0.86

PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 55 of 84

Equipartitioning

What about

$$P(a) \sim a^{-\tau} \quad ?$$

Since $\tau > 1$, suggests no equipartitioning:

$$aP(a) \sim a^{-\tau+1} \neq \text{const}$$

- $P(a)$ overcounts basins within basins ...
- while stream ordering separates basins ...

PoCS @pocsvox Branching Networks II



Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 50 of 84

Generalizing Horton's laws

$$\bar{\ell}_\omega \propto (R_\ell)^\omega \Rightarrow N(\ell|\omega) = (R_n R_\ell)^{-\omega} F_\ell(\ell/R_\ell^\omega)$$

$$\bar{a}_\omega \propto (R_a)^\omega \Rightarrow N(a|\omega) = (R_n^2)^{-\omega} F_a(a/R_n^\omega)$$




- Scaling collapse works well for intermediate orders
- All moments grow exponentially with order

PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 53 of 84

Combining stream segments distributions:



Stream segments sum to give main stream lengths

$$\ell_\omega = \sum_{\mu=1}^{\mu=\omega} s_\mu$$

$P(\ell_\omega)$ is a convolution of distributions for the s_ω

PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 56 of 84

Fluctuations

Moving beyond the mean:

Both Horton's laws and Tokunaga's law relate average properties, e.g.,

$$\bar{s}_\omega / \bar{s}_{\omega-1} = R_s$$

- Natural generalization to consider relationships between probability distributions
- Yields rich and full description of branching network structure
- See into the heart of randomness ...


PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 51 of 84

Generalizing Horton's laws

How well does overall basin fit internal pattern?



- Actual length = 4920 km (at 1 km res)
- Predicted Mean length = 11100 km
- Predicted Std dev = 5600 km
- Actual length/Mean length = 44 %
- Okay.

PoCS @pocsvox Branching Networks II


Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 54 of 84

Generalizing Horton's laws

Sum of variables $\ell_\omega = \sum_{\mu=1}^{\mu=\omega} s_\mu$ leads to convolution of distributions:

$$N(\ell|\omega) = N(s|1) * N(s|2) * \dots * N(s|\omega)$$



$$N(s|\omega) = \frac{1}{R_n^\omega R_\ell^\omega} F(s/R_\ell^\omega)$$

$$F(x) = e^{-x/\xi}$$

Mississippi: $\xi \approx 900$ m.


PoCS @pocsvox Branching Networks II

Horton ⇄ Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

📊 57 of 84

Generalizing Horton's laws

- Next level up: Main stream length distributions must combine to give overall distribution for stream length



- $P(\ell) \sim \ell^{-\gamma}$
- Another round of convolutions [3]
- Interesting ...




58 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

Mississippi:



- Same data collapse for Mississippi ...



61 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

- Follow stream segments down stream from their beginning
- Probability (or rate) of an order μ stream segment terminating is **constant**:

$$\tilde{p}_\mu \approx 1/(R_s)^{\mu-1} \xi_s$$

- Probability decays exponentially with stream order
- Inter-tributary lengths exponentially distributed
- ⇒ random spatial distribution of stream segments




64 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Horton's laws

- Number and area distributions for the Scheidegger model [3]
- $P(n_{1,6})$ versus $P(a_6)$ for a randomly selected $\omega = 6$ basin.



59 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

So

$$P(T_{\mu, \nu}) = (R_s)^{\mu-\nu-1} P_t [T_{\mu, \nu}/(R_s)^{\mu-\nu-1}]$$

where

$$P_t(z) = \frac{1}{\xi_t} e^{-z/\xi_t}$$

$$P(s_\mu) \Leftrightarrow P(T_{\mu, \nu})$$

- Exponentials arise from randomness.
- Look at joint probability $P(s_\mu, T_{\mu, \nu})$.




62 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

Scheidegger:



- Observe exponential distributions for $T_{\mu, \nu}$
- Scaling collapse works using R_s




60 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

Network architecture:

- Inter-tributary lengths exponentially distributed
- Leads to random spatial distribution of stream segments



62 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

- Now deal with this thing:

$$P(s_\mu, T_{\mu, \nu}) = \tilde{p}_\mu \left(\frac{s_\mu - 1}{T_{\mu, \nu}} \right) p_\nu^{T_{\mu, \nu}} (1 - p_\nu - \tilde{p}_\mu)^{s_\mu - T_{\mu, \nu} - 1}$$


- Set $(x, y) = (s_\mu, T_{\mu, \nu})$ and $q = 1 - p_\nu - \tilde{p}_\mu$, approximate liberally.

- Obtain

$$P(x, y) = N x^{-1/2} [F(y/x)]^x$$

where

$$F(v) = \left(\frac{1-v}{q} \right)^{-(1-v)} \left(\frac{v}{p} \right)^{-v}$$



65 of 84

- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References



66 of 84


- Horton ⇄ Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References

Generalizing Tokunaga's law

PoCS
@pocsvox
Branching
Networks II

Checking form of $P(s_\mu, T_{\mu,\nu})$ works:

Scheidegger:



Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References


67 of 84

Generalizing Tokunaga's law

PoCS
@pocsvox
Branching
Networks II

Checking form of $P(s_\mu, T_{\mu,\nu})$ works:

Scheidegger:



Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References


68 of 84

Generalizing Tokunaga's law

PoCS
@pocsvox
Branching
Networks II

Checking form of $P(s_\mu, T_{\mu,\nu})$ works:

Scheidegger:



Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References


69 of 84

Generalizing Tokunaga's law

PoCS
@pocsvox
Branching
Networks II

Checking form of $P(s_\mu, T_{\mu,\nu})$ works:

Mississippi:




Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

70 of 84

Models

Random subnetworks on a Bethe lattice [13]



- Dominant theoretical concept for several decades.
- Bethe lattices are fun and tractable.
- Led to idea of "Statistical inevitability" of river network statistics [7]
- But Bethe lattices unconnected with surfaces.
- In fact, Bethe lattices \approx infinite dimensional spaces (oops).
- So let's move on ...

PoCS
@pocsvox
Branching
Networks II

Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

72 of 84

Optimal channel networks

PoCS
@pocsvox
Branching
Networks II

Rodríguez-Iturbe, Rinaldo, et al. [10]

Landscapes $h(\vec{x})$ evolve such that energy dissipation $\dot{\epsilon}$ is minimized, where

$$\dot{\epsilon} \propto \int d\vec{r} (\text{flux}) \times (\text{force}) \sim \sum_i a_i \nabla h_i \sim \sum_i a_i^\gamma$$

- Landscapes obtained numerically give exponents near that of real networks.
- **But:** numerical method used matters.
- **And:** Maritan et al. find basic universality classes are that of Scheidegger, self-similar, and a third kind of random network [8]

Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

74 of 84

Theoretical networks

Summary of universality classes:

network	h	d
Non-convergent flow	1	1
Directed random	2/3	1
Undirected random	5/8	5/4
Self-similar	1/2	1
OCN's (I)	1/2	1
OCN's (II)	2/3	1
OCN's (III)	3/5	1
Real rivers	0.5-0.7	1.0-1.2

$$h \Rightarrow \ell \propto a^h \text{ (Hack's law)}$$

$$d \Rightarrow \ell \propto L_{\parallel}^d \text{ (stream self-affinity)}$$

PoCS
@pocsvox
Branching
Networks II


Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

75 of 84

Scheidegger's model

PoCS
@pocsvox
Branching
Networks II

Directed random networks [11, 12]



$$P(\swarrow) = P(\searrow) = 1/2$$

Functional form of all scaling laws exhibited but exponents differ from real world [15, 16, 14]

Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

73 of 84

Nutshell

PoCS
@pocsvox
Branching
Networks II

Branching networks II Key Points:

- Horton's laws and Tokunaga's law all fit together.
- For 2-d networks, these laws are 'planform' laws and ignore slope.
- Abundant scaling relations can be derived.
- Can take R_n , R_ℓ , and d as three independent parameters necessary to describe all 2-d branching networks.
- For scaling laws, only $h = \ln R_\ell / \ln R_n$ and d are needed.
- Laws can be extended nicely to laws of distributions.
- Numerous models of branching network evolution exist: nothing rock solid yet ...?

Horton ⇌
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
References

76 of 84

References I

- [1] H. de Vries, T. Becker, and B. Eckhardt. Power law distribution of discharge in ideal networks. [Water Resources Research](#), 30(12):3541–3543, 1994. [pdf](#)
- [2] P. S. Dodds and D. H. Rothman. Unified view of scaling laws for river networks. [Physical Review E](#), 59(5):4865–4877, 1999. [pdf](#)
- [3] P. S. Dodds and D. H. Rothman. Geometry of river networks. II. Distributions of component size and number. [Physical Review E](#), 63(1):016116, 2001. [pdf](#)

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 77 of 84

References II

- [4] P. S. Dodds and D. H. Rothman. Geometry of river networks. III. Characterization of component connectivity. [Physical Review E](#), 63(1):016117, 2001. [pdf](#)
- [5] N. Goldenfeld. [Lectures on Phase Transitions and the Renormalization Group](#), volume 85 of [Frontiers in Physics](#). Addison-Wesley, Reading, Massachusetts, 1992.
- [6] J. T. Hack. Studies of longitudinal stream profiles in Virginia and Maryland. [United States Geological Survey Professional Paper](#), 294-B:45–97, 1957. [pdf](#)

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 78 of 84

References III

- [7] J. W. Kirchner. Statistical inevitability of Horton's laws and the apparent randomness of stream channel networks. [Geology](#), 21:591–594, 1993. [pdf](#)
- [8] A. Maritan, F. Colaiori, A. Flammini, M. Cieplak, and J. R. Banavar. Universality classes of optimal channel networks. [Science](#), 272:984–986, 1996. [pdf](#)
- [9] S. D. Peckham. New results for self-similar trees with applications to river networks. [Water Resources Research](#), 31(4):1023–1029, 1995.

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 79 of 84

References IV

- [10] I. Rodríguez-Iturbe and A. Rinaldo. [Fractal River Basins: Chance and Self-Organization](#). Cambridge University Press, Cambridge, UK, 1997.
- [11] A. E. Scheidegger. A stochastic model for drainage patterns into an intramontane trench. [Bull. Int. Assoc. Sci. Hydrol.](#), 12(1):15–20, 1967. [pdf](#)
- [12] A. E. Scheidegger. [Theoretical Geomorphology](#). Springer-Verlag, New York, third edition, 1991.

References V

- [13] R. L. Shreve. Infinite topologically random channel networks. [Journal of Geology](#), 75:178–186, 1967. [pdf](#)
- [14] H. Takayasu. Steady-state distribution of generalized aggregation system with injection. [Physical Review Letters](#), 63(23):2563–2565, 1989. [pdf](#)
- [15] H. Takayasu, I. Nishikawa, and H. Tasaki. Power-law mass distribution of aggregation systems with injection. [Physical Review A](#), 37(8):3110–3117, 1988.

References VI

- [16] M. Takayasu and H. Takayasu. Apparent independency of an aggregation system with injection. [Physical Review A](#), 39(8):4345–4347, 1989. [pdf](#)
- [17] D. G. Tarboton, R. L. Bras, and I. Rodríguez-Iturbe. Comment on “On the fractal dimension of stream networks” by Paolo La Barbera and Renzo Rosso. [Water Resources Research](#), 26(9):2243–4, 1990. [pdf](#)
- [18] E. Tokunaga. The composition of drainage network in Toyohira River Basin and the valuation of Horton's first law. [Geophysical Bulletin of Hokkaido University](#), 15:1–19, 1966. [pdf](#)

References VII

- [19] E. Tokunaga. Consideration on the composition of drainage networks and their evolution. [Geographical Reports of Tokyo Metropolitan University](#), 13:G1–27, 1978. [pdf](#)
- [20] E. Tokunaga. Ordering of divide segments and law of divide segment numbers. [Transactions of the Japanese Geomorphological Union](#), 5(2):71–77, 1984.
- [21] S. D. Willett, S. W. McCoy, J. T. Perron, L. Goren, and C.-Y. Chen. Dynamic reorganization of river basins. [Science](#), 343(6175):1248765, 2014. [pdf](#)

References VIII

- [22] G. K. Zipf. [Human Behaviour and the Principle of Least-Effort](#). Addison-Wesley, Cambridge, MA, 1949.

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 80 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 81 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 82 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 83 of 84

PoCS
@pocsvox
Branching
Networks II

Horton ↔
Tokunaga
Reducing Horton
Scaling relations
Fluctuations
Models
Nutshell
[References](#)

 84 of 84