

# Branching Networks II

Last updated: 2023/01/26, 11:44:57 EST

Principles of Complex Systems, Vols. 1, 2, & 3D  
CSYS/MATH 300, 303, & 394, 2022-2023 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center  
Santa Fe Institute | University of Vermont



Licensed under the *Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License*.

PoCS  
@pocsvox  
Branching  
Networks II

Horton ↔  
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



These slides are brought to you by:

PoCS  
@pocsvox

Branching  
Networks II

Sealie & Lambie  
Productions



Horton ↔  
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References



# These slides are also brought to you by:

## Special Guest Executive Producer



 On Instagram at [pratchett\\_the\\_cat](https://www.instagram.com/pratchett_the_cat) 

PoCS  
@pocsvox

Branching  
Networks II

Horton ↔  
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Outline

PoCS  
@pocsvox

Branching  
Networks II

Horton  $\Leftrightarrow$  Tokunaga

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Reducing Horton

Scaling relations

Scaling relations

Fluctuations

Fluctuations

Models

Models

Nutshell

Nutshell

References


References






# Piracy on the high $\chi$ 's:



“Dynamic Reorganization of River Basins” 

Willett et al.,

Science, **343**, 1248765, 2014. [21]



$$\frac{\partial z(x, t)}{\partial t} = U - K A^m \left| \frac{\partial z(x, t)}{\partial x} \right|^n$$

$$z(x) = z_b + \left( \frac{U}{K A_0^m} \right)^{1/n} \chi$$

$$\chi = \int_{x_b}^x \left( \frac{A_0}{A(x')} \right)^{m/n} dx'$$

# Piracy on the high $\chi$ 's:

PoCS  
@pocsvox

Branching  
Networks II

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations


Fluctuations

Models

Nutshell

References

[https://www.youtube.com/watch?v=FnroL1\\_-l2c?rel=0](https://www.youtube.com/watch?v=FnroL1_-l2c?rel=0)

More: [How river networks move across a landscape](#)   
(Science Daily)










# Can Horton and Tokunaga be happy?

## Horton and Tokunaga seem different:

- 🧱 In terms of network architecture, Horton's laws appear to contain less detailed information than Tokunaga's law.
- 🧱 Oddly, Horton's laws have **four** parameters and Tokunaga has **two** parameters.
- 🧱  $R_n, R_a, R_\ell,$  and  $R_s$  **versus**  $T_1$  and  $R_T$ . One simple redundancy:  $R_\ell = R_s$ .  
[Insert question from assignment 15](#) 
- 🧱 To make a connection, clearest approach is to start with Tokunaga's law ...
- 🧱 Known result: Tokunaga  $\rightarrow$  Horton <sup>[18, 19, 20, 9, 2]</sup>

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell





References



# Let us make them happy

We need one more ingredient:

## Space-fillingness

-  A network is **space-filling** if the average distance between adjacent streams is roughly constant.
-  Reasonable for river and cardiovascular networks
-  For river networks:  
**Drainage density**  $\rho_{dd}$  = inverse of typical distance between channels in a landscape.
-  In terms of basin characteristics:

$$\rho_{dd} \simeq \frac{\sum \text{stream segment lengths}}{\text{basin area}} = \frac{\sum_{\omega=1}^{\Omega} n_{\omega} \bar{s}_{\omega}}{a_{\Omega}}$$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References





# More with the happy-making thing

Start with Tokunaga's law:  $T_k = T_1 R_T^{k-1}$

 Start looking for Horton's stream number law:

$$n_\omega / n_{\omega+1} = R_n.$$

 Estimate  $n_\omega$ , the number of streams of order  $\omega$  in terms of other  $n_{\omega'}, \omega' > \omega$ .

 Observe that each stream of order  $\omega$  terminates by either:


1. Running into another stream of order  $\omega$  and generating a stream of order  $\omega + 1$

...

▶  $2n_{\omega+1}$  streams of order  $\omega$  do this

2. Running into and being absorbed by a stream of higher order  $\omega' > \omega$  ...

▶  $n_{\omega'} T_{\omega'-\omega}$  streams of order  $\omega$  do this



Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References






# More with the happy-making thing


Putting things together:



$$n_{\omega} = \underbrace{2n_{\omega+1}}_{\text{generation}} + \sum_{\omega'=\omega+1}^{\Omega} \underbrace{T_{\omega'-\omega}n_{\omega'}}_{\text{absorption}}$$

 Use Tokunaga's law and manipulate expression to find Horton's law for stream numbers follows and hence obtain  $R_n$ .

 [Insert question from assignment 16](#) 

 Solution:

$$R_n = \frac{(2 + R_T + T_1) \pm \sqrt{(2 + R_T + T_1)^2 - 8R_T}}{2}$$

(The larger value is the one we want.)

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Finding other Horton ratios

## Connect Tokunaga to $R_s$

- Now use uniform drainage density  $\rho_{dd}$ .
- Assume side streams are roughly separated by distance  $1/\rho_{dd}$ .
- For an order  $\omega$  **stream segment**, expected length is

$$\bar{s}_\omega \simeq \rho_{dd}^{-1} \left( 1 + \sum_{k=1}^{\omega-1} T_k \right)$$

- Substitute in Tokunaga's law  $T_k = T_1 R_T^{k-1}$ :

$$\bar{s}_\omega \simeq \rho_{dd}^{-1} \left( 1 + T_1 \sum_{k=1}^{\omega-1} R_T^{k-1} \right) \propto R_T^\omega$$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Horton and Tokunaga are happy

Altogether then:



$$\Rightarrow \bar{s}_\omega / \bar{s}_{\omega-1} = R_T \Rightarrow R_s = R_T$$



Recall  $R_\ell = R_s$  so

$$R_\ell = R_s = R_T$$



And from before:

$$R_n = \frac{(2 + R_T + T_1) + \sqrt{(2 + R_T + T_1)^2 - 8R_T}}{2}$$

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



# Horton and Tokunaga are happy

## Some observations:

- 🧱  $R_n$  and  $R_\ell$  depend on  $T_1$  and  $R_T$ .
- 🧱 Seems that  $R_a$  must as well ...
- 🧱 Suggests Horton's laws must contain some redundancy
- 🧱 We'll in fact see that  $R_a = R_n$ .
- 🧱 Also: Both Tokunaga's law and Horton's laws can be generalized to relationships between non-trivial statistical distributions. [3, 4]

Horton ↔  
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell


References





# Horton and Tokunaga are happy

## The other way round


 Note: We can invert the expressions for  $R_n$  and  $R_\ell$  to find Tokunaga's parameters in terms of Horton's parameters.



$$R_T = R_\ell,$$



$$T_1 = R_n - R_\ell - 2 + 2R_\ell/R_n.$$

 Suggests we should be able to argue that Horton's laws imply Tokunaga's laws (if drainage density is uniform) ...

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell


References





# Horton and Tokunaga are friends


## From Horton to Tokunaga [2]



 Assume Horton's laws hold for number and length

 Start with picture showing an order  $\omega$  stream and order  $\omega - 1$  generating and side streams.

 Scale up by a factor of  $R_s$ , orders increment to  $\omega + 1$  and  $\omega$ .

 Maintain drainage density by adding new order  $\omega - 1$  streams

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Horton and Tokunaga are friends

...and in detail:

- Must retain same drainage density.
- Add an extra  $(R_\ell - 1)$  first order streams for each original tributary.
- Since by definition, an order  $\omega + 1$  stream segment has  $T_\omega$  order 1 side streams, we have:

$$T_k = (R_\ell - 1) \left( 1 + \sum_{i=1}^{k-1} T_i \right).$$

- For large  $\omega$ , Tokunaga's law is the solution—let's check ...

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References



# Horton and Tokunaga are friends

Just checking:

 Substitute Tokunaga's law  $T_i = T_1 R_T^{i-1} = T_1 R_\ell^{i-1}$  into

$$T_k = (R_\ell - 1) \left( 1 + \sum_{i=1}^{k-1} T_i \right)$$



$$T_k = (R_\ell - 1) \left( 1 + \sum_{i=1}^{k-1} T_1 R_\ell^{i-1} \right)$$

$$= (R_\ell - 1) \left( 1 + T_1 \frac{R_\ell^{k-1} - 1}{R_\ell - 1} \right)$$

$$\simeq (R_\ell - 1) T_1 \frac{R_\ell^{k-1}}{R_\ell - 1} = T_1 R_\ell^{k-1} \quad \dots\text{yep.}$$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations







Models

Nutshell

References



# Horton's laws of area and number:



Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



In bottom plots, stream number graph has been flipped vertically.



Highly suggestive that  $R_n \equiv R_a \dots$



# Measuring Horton ratios is tricky:

PoCS  
@pocsvox  
Branching  
Networks II

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



How robust are our estimates of ratios?



Rule of thumb: discard data for two smallest and two largest orders.



# Mississippi:

PoCS  
@pocsvox

Branching  
Networks II

$\omega$ range	$R_n$	$R_a$	$R_\ell$	$R_s$	$R_a/R_n$
[2, 3]	5.27	5.26	2.48	2.30	1.00
[2, 5]	4.86	4.96	2.42	2.31	1.02
[2, 7]	4.77	4.88	2.40	2.31	1.02
[3, 4]	4.72	4.91	2.41	2.34	1.04
[3, 6]	4.70	4.83	2.40	2.35	1.03
[3, 8]	4.60	4.79	2.38	2.34	1.04
[4, 6]	4.69	4.81	2.40	2.36	1.02
[4, 8]	4.57	4.77	2.38	2.34	1.05
[5, 7]	4.68	4.83	2.36	2.29	1.03
[6, 7]	4.63	4.76	2.30	2.16	1.03
[7, 8]	4.16	4.67	2.41	2.56	1.12
mean $\mu$	4.69	4.85	2.40	2.33	1.04
std dev $\sigma$	0.21	0.13	0.04	0.07	0.03
$\sigma/\mu$	0.045	0.027	0.015	0.031	0.024

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



$\omega$ range	$R_n$	$R_a$	$R_\ell$	$R_s$	$R_a/R_n$
[2, 3]	4.78	4.71	2.47	2.08	0.99
[2, 5]	4.55	4.58	2.32	2.12	1.01
[2, 7]	4.42	4.53	2.24	2.10	1.02
[3, 5]	4.45	4.52	2.26	2.14	1.01
[3, 7]	4.35	4.49	2.20	2.10	1.03
[4, 6]	4.38	4.54	2.22	2.18	1.03
[5, 6]	4.38	4.62	2.22	2.21	1.06
[6, 7]	4.08	4.27	2.05	1.83	1.05
mean $\mu$	4.42	4.53	2.25	2.10	1.02
std dev $\sigma$	0.17	0.10	0.10	0.09	0.02
$\sigma/\mu$	0.038	0.023	0.045	0.042	0.019

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell


References





# Reducing Horton's laws:

Rough first effort to show  $R_n \equiv R_a$ :

  $a_\Omega \propto$  sum of all stream segment lengths in a order  $\Omega$  basin (assuming uniform drainage density)

 So:

$$\begin{aligned} a_\Omega &\simeq \sum_{\omega=1}^{\Omega} n_\omega \bar{s}_\omega / \rho_{dd} \\ &\propto \sum_{\omega=1}^{\Omega} \underbrace{R_n^{\Omega-\omega} \cdot \hat{1}^{n_\Omega}}_{n_\omega} \underbrace{\bar{s}_1 \cdot R_s^{\omega-1}}_{\bar{s}_\omega} \\ &= \frac{R_n^\Omega}{R_s} \bar{s}_1 \sum_{\omega=1}^{\Omega} \left( \frac{R_s}{R_n} \right)^\omega \end{aligned}$$




# Reducing Horton's laws:

Continued ...



$$\begin{aligned} a_{\Omega} &\propto \frac{R_n^{\Omega}}{R_s} \bar{s}_1 \sum_{\omega=1}^{\Omega} \left( \frac{R_s}{R_n} \right)^{\omega} \\ &= \frac{R_n^{\Omega}}{R_s} \bar{s}_1 \frac{R_s}{R_n} \frac{1 - (R_s/R_n)^{\Omega}}{1 - (R_s/R_n)} \\ &\sim R_n^{\Omega-1} \bar{s}_1 \frac{1}{1 - (R_s/R_n)} \text{ as } \Omega \nearrow \end{aligned}$$

 So,  $a_{\Omega}$  is growing like  $R_n^{\Omega}$  and therefore:

$$R_n \equiv R_a$$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References



# Reducing Horton's laws:

Not quite:

- ...But this only a rough argument as Horton's laws do not imply a strict hierarchy
- Need to account for sidebranching.
- Insert question from assignment 16 

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations


Models


Nutshell


References




## Intriguing division of area:

 Observe: Combined area of basins of order  $\omega$  independent of  $\omega$ .

 Not obvious: basins of low orders not necessarily contained in basin on higher orders.

 Story:

$$R_n \equiv R_a \Rightarrow \boxed{n_\omega \bar{a}_\omega = \text{const}}$$

 Reason:

$$n_\omega \propto (R_n)^{-\omega}$$

$$\bar{a}_\omega \propto (R_a)^\omega \propto n_\omega^{-1}$$

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models




Nutshell

References



# Equipartitioning:

Some examples:



PoCS  
@pocsvox

Branching  
Networks II

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Neural Reboot: Fwoompf

PoCS  
@pocsvox

Branching  
Networks II

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



<https://www.youtube.com/watch?v=5mUs70SqD4o?rel=0>



## The story so far:

- 🧱 Natural branching networks are **hierarchical**, **self-similar** structures
- 🧱 Hierarchy is **mixed**
- 🧱 Tokunaga's law describes detailed architecture:  
 $T_k = T_1 R_T^{k-1}$ .
- 🧱 We have connected Tokunaga's and Horton's laws
- 🧱 Only two Horton laws are independent ( $R_n = R_a$ )
- 🧱 Only **two** parameters are **independent**:  
 $(T_1, R_T) \Leftrightarrow (R_n, R_s)$

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



## A little further ...

- Ignore stream ordering for the moment
- Pick a random location on a branching network  $p$ .
- Each point  $p$  is associated with a basin and a longest stream length
- Q:** What is probability that the  $p$ 's drainage basin has area  $a$ ?  $P(a) \propto a^{-\tau}$  for large  $a$
- Q:** What is probability that the longest stream from  $p$  has length  $\ell$ ?  $P(\ell) \propto \ell^{-\gamma}$  for large  $\ell$
- Roughly observed:  $1.3 \lesssim \tau \lesssim 1.5$  and  $1.7 \lesssim \gamma \lesssim 2.0$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References





## Probability distributions with power-law decays



We see them everywhere:

- Earthquake magnitudes (Gutenberg-Richter law)
- City sizes (Zipf's law)
- Word frequency (Zipf's law)<sup>[22]</sup>
- Wealth (maybe not—at least heavy tailed)
- Statistical mechanics (phase transitions)<sup>[5]</sup>



A big part of the story of complex systems



Arise from **mechanisms**: growth, randomness, optimization, ...



Our task is always to illuminate the mechanism ...

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



## Connecting exponents

- ☰ We have the detailed picture of branching networks (Tokunaga and Horton)
- ☰ Plan: Derive  $P(a) \propto a^{-\tau}$  and  $P(\ell) \propto \ell^{-\gamma}$  starting with Tokunaga/Horton story<sup>[17, 1, 2]</sup>
- ☰ Let's work on  $P(\ell)$  ...
- ☰ Our first fudge: assume Horton's laws hold throughout a basin of order  $\Omega$ .
- ☰ (We know they deviate from strict laws for low  $\omega$  and high  $\omega$  but not too much.)
- ☰ Next: place stick between teeth. Bite stick. Proceed.

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



## Finding $\gamma$ :



Often useful to work with **cumulative distributions**, especially when dealing with power-law distributions.



The complementary cumulative distribution turns out to be most useful:

$$P_{>}(l_*) = P(l > l_*) = \int_{l=l_*}^{l_{\max}} P(l) dl$$



$$P_{>}(l_*) = 1 - P(l < l_*)$$



Also known as the exceedance probability.

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell


References



# Scaling laws

## Finding $\gamma$ :

 The connection between  $P(x)$  and  $P_{>}(x)$  when  $P(x)$  has a power law tail is simple:

 Given  $P(l) \sim l^{-\gamma}$  large  $l$  then for large enough  $l_*$

$$P_{>}(l_*) = \int_{l=l_*}^{l_{\max}} P(l) dl$$

$$\sim \int_{l=l_*}^{l_{\max}} l^{-\gamma} dl$$

$$= \frac{l^{-(\gamma-1)}}{-(\gamma-1)} \Big|_{l=l_*}^{l_{\max}}$$

$$\propto l_*^{-(\gamma-1)} \quad \text{for } l_{\max} \gg l_*$$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations





Models

Nutshell

References




## Finding $\gamma$ :

-  **Aim:** determine probability of randomly choosing a point on a network with main stream length  $> l_*$
-  Assume some spatial sampling resolution  $\Delta$
-  Landscape is broken up into grid of  $\Delta \times \Delta$  sites
-  Approximate  $P_{>}(l_*)$  as

$$P_{>}(l_*) = \frac{N_{>}(l_*; \Delta)}{N_{>}(0; \Delta)}$$


where  $N_{>}(l_*; \Delta)$  is the number of sites with main stream length  $> l_*$ .

-  Use Horton's law of stream segments:  
 $\bar{s}_\omega / \bar{s}_{\omega-1} = R_s \dots$

- Horton  $\Leftrightarrow$  Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References





## Finding $\gamma$ :


 Set  $l_* = \bar{l}_\omega$  for some  $1 \ll \omega \ll \Omega$ .



$$P_{>}(\bar{l}_\omega) = \frac{N_{>}(\bar{l}_\omega; \Delta)}{N_{>}(0; \Delta)} \simeq \frac{\sum_{\omega'=\omega+1}^{\Omega} n_{\omega'} \bar{s}_{\omega'} / \Delta}{\sum_{\omega'=1}^{\Omega} n_{\omega'} \bar{s}_{\omega'} / \Delta}$$

  $\Delta$ 's cancel


 Denominator is  $a_{\Omega} \rho_{dd}$ , a constant.

 So ...using Horton's laws ...


$$P_{>}(\bar{l}_\omega) \propto \sum_{\omega'=\omega+1}^{\Omega} n_{\omega'} \bar{s}_{\omega'} \simeq \sum_{\omega'=\omega+1}^{\Omega} (1 \cdot R_n^{\Omega-\omega'}) (\bar{s}_1 \cdot R_s^{\omega'-1})$$




## Finding $\gamma$ :


 We are here:

$$P_{>}(\bar{l}_{\omega}) \propto \sum_{\omega'=\omega+1}^{\Omega} (1 \cdot R_n^{\Omega-\omega'}) (\bar{s}_1 \cdot R_s^{\omega'-1})$$

 Cleaning up irrelevant constants:

$$P_{>}(\bar{l}_{\omega}) \propto \sum_{\omega'=\omega+1}^{\Omega} \left( \frac{R_s}{R_n} \right)^{\omega'}$$

 Change summation order by substituting  $\omega'' = \Omega - \omega'$ .

 Sum is now from  $\omega'' = 0$  to  $\omega'' = \Omega - \omega - 1$  (equivalent to  $\omega' = \Omega$  down to  $\omega' = \omega + 1$ )

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell


References



## Finding $\gamma$ :



$$P_{>}(\bar{l}_\omega) \propto \sum_{\omega''=0}^{\Omega-\omega-1} \left(\frac{R_s}{R_n}\right)^{\Omega-\omega''} \propto \sum_{\omega''=0}^{\Omega-\omega-1} \left(\frac{R_n}{R_s}\right)^{\omega''}$$

 Since  $R_n > R_s$  and  $1 \ll \omega \ll \Omega$ ,

$$P_{>}(\bar{l}_\omega) \propto \left(\frac{R_n}{R_s}\right)^{\Omega-\omega} \propto \left(\frac{R_n}{R_s}\right)^{-\omega}$$

again using  $\sum_{i=0}^{n-1} a^i = (a^n - 1)/(a - 1)$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References







## Finding $\gamma$ :

 Nearly there:

$$P_{>}(\bar{l}_{\omega}) \propto \left( \frac{R_n}{R_s} \right)^{-\omega} = e^{-\omega \ln(R_n/R_s)}$$

 Need to express right hand side in terms of  $\bar{l}_{\omega}$ .

 Recall that  $\bar{l}_{\omega} \simeq \bar{l}_1 R_{\ell}^{\omega-1}$ .



$$\bar{l}_{\omega} \propto R_{\ell}^{\omega} = R_s^{\omega} = e^{\omega \ln R_s}$$

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Scaling laws

## Finding $\gamma$ :

Therefore:

$$P_{>}(\bar{l}_{\omega}) \propto e^{-\omega \ln(R_n/R_s)} = (e^{\omega \ln R_s})^{-\ln(R_n/R_s)/\ln(R_s)}$$

$$\propto \bar{l}_{\omega}^{-\ln(R_n/R_s)/\ln R_s}$$

$$= \bar{l}_{\omega}^{-(\ln R_n - \ln R_s)/\ln R_s}$$


$$= \bar{l}_{\omega}^{-\ln R_n/\ln R_s + 1}$$

$$= \bar{l}_{\omega}^{-\gamma + 1}$$




# Scaling laws


## Finding $\gamma$ :


 And so we have:


$$\gamma = \ln R_n / \ln R_s$$

 Proceeding in a similar fashion, we can show

$$\tau = 2 - \ln R_s / \ln R_n = 2 - 1/\gamma$$

Insert question from assignment 16 

 Such connections between exponents are called **scaling relations**

 Let's connect to one last relationship: Hack's law

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Scaling laws

Hack's law: [6]



$$l \propto a^h$$



Typically observed that  $0.5 \lesssim h \lesssim 0.7$ .



Use Horton laws to connect  $h$  to Horton ratios:

$$\bar{l}_\omega \propto R_s^\omega \text{ and } \bar{a}_\omega \propto R_n^\omega$$



Observe:

$$\bar{l}_\omega \propto e^{\omega \ln R_s} \propto (e^{\omega \ln R_n})^{\ln R_s / \ln R_n}$$

$$\propto (R_n^\omega)^{\ln R_s / \ln R_n} \propto \bar{a}_\omega^{\ln R_s / \ln R_n} \Rightarrow h = \ln R_s / \ln R_n$$

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# We mentioned there were a good number of 'laws': [2]

**Relation:**   **Name or description:**

$$T_k = T_1 (R_T)^{k-1}$$
$$\ell \sim L^d$$

Tokunaga's law

self-affinity of single channels

$$n_\omega / n_{\omega+1} = R_n$$
$$\ell_{\omega+1} / \ell_\omega = R_\ell$$

Horton's law of stream numbers

Horton's law of main stream lengths

$$\bar{a}_{\omega+1} / \bar{a}_\omega = R_a$$
$$\bar{s}_{\omega+1} / \bar{s}_\omega = R_s$$

Horton's law of basin areas

Horton's law of stream segment lengths

$$L_\perp \sim L^H$$

scaling of basin widths

$$P(a) \sim a^{-\tau}$$

probability of basin areas

$$P(\ell) \sim \ell^{-\gamma}$$

probability of stream lengths

$$\ell \sim a^h$$

Hack's law

$$a \sim L^D$$

scaling of basin areas

$$\Lambda \sim a^\beta$$

Langbein's law

$$\lambda \sim L^\varphi$$

variation of Langbein's law

on  $\leftrightarrow$   
inaga

cing Horton

ng relations

uations

els

hell

rences



# Connecting exponents

Only 3 parameters are independent:  
e.g., take  $d$ ,  $R_n$ , and  $R_s$

relation:	scaling relation/parameter: [2]
$\ell \sim L^d$	$d$
$T_k = T_1 (R_T)^{k-1}$	$T_1 = R_n - R_s - 2 + 2R_s/R_n$ $R_T = R_s$
$n_\omega/n_{\omega+1} = R_n$	$R_n$
$\bar{a}_{\omega+1}/\bar{a}_\omega = R_a$	$R_a = R_n$
$\bar{\ell}_{\omega+1}/\bar{\ell}_\omega = R_\ell$	$R_\ell = R_s$
$\ell \sim a^h$	$h = \ln R_s / \ln R_n$
$a \sim L^D$	$D = d/h$
$L_\perp \sim L^H$	$H = d/h - 1$
$P(a) \sim a^{-\tau}$	$\tau = 2 - h$
$P(\ell) \sim \ell^{-\gamma}$	$\gamma = 1/h$
$\Lambda \sim a^\beta$	$\beta = 1 + h$
$\lambda \sim L^\varphi$	$\varphi = d$




# Scheidegger's model

PoCS  
@pocsvox

Branching  
Networks II

## Directed random networks <sup>[11, 12]</sup>



$$P(\searrow) = P(\swarrow) = 1/2$$



Functional form of all scaling laws exhibited but exponents differ from real world <sup>[15, 16, 14]</sup>



Useful and interesting test case

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell


References




# A toy model—Scheidegger's model

PoCS  
@pocsvox  
Branching  
Networks II

## Random walk basins:

 Boundaries of basins are random walks



Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell


References





# Scheidegger's model

PoCS  
@pocsvox  
Branching  
Networks II



Horton  $\leftrightarrow$   
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



# Scheidegger's model

Prob for first return of a random walk in (1+1) dimensions (from CSYS/MATH 300):



$$P(n) \sim \frac{1}{2\sqrt{\pi}} n^{-3/2}.$$

and so  $P(\ell) \propto \ell^{-3/2}$ .



Typical area for a walk of length  $n$  is  $\propto n^{3/2}$ :

$$\ell \propto a^{2/3}.$$



Find  $\tau = 4/3$ ,  $h = 2/3$ ,  $\gamma = 3/2$ ,  $d = 1$ .



Note  $\tau = 2 - h$  and  $\gamma = 1/h$ .



$R_n$  and  $R_\ell$  have not been derived analytically.

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References






# Equipartitioning reexamined:

Recall this story:

PoCS  
@pocsvox

Branching  
Networks II



Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations


Fluctuations

Models


Nutshell

References





 What about

$$P(a) \sim a^{-\tau} \quad ?$$

 Since  $\tau > 1$ , suggests no equipartitioning:

$$aP(a) \sim a^{-\tau+1} \neq \text{const}$$

  $P(a)$  overcounts basins within basins ...

 while stream ordering separates basins ...

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



## Moving beyond the mean:

- Both Horton's laws and Tokunaga's law relate average properties, e.g.,

$$\bar{s}_\omega / \bar{s}_{\omega-1} = R_s$$

- Natural generalization to consider relationships between **probability distributions**
- Yields rich and full description of branching network structure
- See into the heart of randomness ...

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References





# A toy model—Scheidegger's model

PoCS  
@pocsvox

Branching  
Networks II

Directed random networks <sup>[11, 12]</sup>



$$P(\searrow) = P(\swarrow) = 1/2$$



Flow is directed downwards

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Generalizing Horton's laws

$\bar{\ell}_\omega \propto (R_\ell)^\omega \Rightarrow N(\ell|\omega) = (R_n R_\ell)^{-\omega} F_\ell(\ell/R_\ell^\omega)$

$\bar{a}_\omega \propto (R_a)^\omega \Rightarrow N(a|\omega) = (R_n^2)^{-\omega} F_a(a/R_n^\omega)$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton



Scaling relations

Fluctuations

Models

Nutshell

References





Scaling collapse works well for intermediate orders


All **moments** grow exponentially with order




# Generalizing Horton's laws


 How well does overall basin fit internal pattern?




 Actual length = **4920 km** (at 1 km res)

 Predicted Mean length = **11100 km**

 Predicted Std dev = **5600 km**

 Actual length/Mean length = **44 %**

 Okay.

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References





# Generalizing Horton's laws

Comparison of predicted versus measured main stream lengths for large scale river networks (in  $10^3$  km):

basin:	$l_\Omega$	$\bar{l}_\Omega$	$\sigma_l$	$l_\Omega/\bar{l}_\Omega$	$\sigma_l/\bar{l}_\Omega$
Mississippi	4.92	11.10	5.60	0.44	0.51
Amazon	5.75	9.18	6.85	0.63	0.75
Nile	6.49	2.66	2.20	2.44	0.83
Congo	5.07	10.13	5.75	0.50	0.57
Kansas	1.07	2.37	1.74	0.45	0.73
	$a_\Omega$	$\bar{a}_\Omega$	$\sigma_a$	$a_\Omega/\bar{a}_\Omega$	$\sigma_a/\bar{a}_\Omega$
Mississippi	2.74	7.55	5.58	0.36	0.74
Amazon	5.40	9.07	8.04	0.60	0.89
Nile	3.08	0.96	0.79	3.19	0.82
Congo	3.70	10.09	8.28	0.37	0.82
Kansas	0.14	0.49	0.42	0.28	0.86

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations


Models


Nutshell

References




# Combining stream segments distributions:



 Stream segments  
sum to give main  
stream lengths



$$l_\omega = \sum_{\mu=1}^{\mu=\omega} s_\mu$$

  $P(l_\omega)$  is a  
convolution of  
distributions for  
the  $s_\omega$

Horton  $\leftrightarrow$   
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



# Generalizing Horton's laws



Sum of variables  $\ell_\omega = \sum_{\mu=1}^{\mu=\omega} s_\mu$  leads to convolution of distributions:

$$N(\ell|\omega) = N(s|1) * N(s|2) * \dots * N(s|\omega)$$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton


Scaling relations

Fluctuations

Models

Nutshell

References



$$N(s|\omega) = \frac{1}{R_n^\omega R_l^\omega} F(s/R_\ell^\omega)$$


$$F(x) = e^{-x/\xi}$$

Mississippi:  $\xi \simeq 900$  m.



# Generalizing Horton's laws

- Next level up: Main stream length distributions must combine to give overall distribution for stream length



$$P(l) \sim l^{-\gamma}$$



Another round of convolutions <sup>[3]</sup>



Interesting ...




# Generalizing Horton's laws



Number and area distributions for the Scheidegger model [3]



$P(n_{1,6})$  versus  $P(a_6)$  for a randomly selected  $\omega = 6$  basin.



Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models



Nutshell

References



# Generalizing Tokunaga's law

Scheidegger:



Horton  $\leftrightarrow$   
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



Observe exponential distributions for  $T_{\mu,\nu}$





Scaling collapse works using  $R_s$



# Generalizing Tokunaga's law

Mississippi:



- Horton  $\leftrightarrow$  Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References



Same data collapse for Mississippi ...



# Generalizing Tokunaga's law


So


$$P(T_{\mu,\nu}) = (R_s)^{\mu-\nu-1} P_t [T_{\mu,\nu}/(R_s)^{\mu-\nu-1}]$$

where

$$P_t(z) = \frac{1}{\xi_t} e^{-z/\xi_t}.$$

$$P(s_\mu) \Leftrightarrow P(T_{\mu,\nu})$$

 Exponentials arise from randomness.

 Look at joint probability  $P(s_\mu, T_{\mu,\nu})$ .

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References





# Generalizing Tokunaga's law

PoCS  
@pocsvox

Branching  
Networks II

## Network architecture:

- Inter-tributary lengths exponentially distributed
- Leads to random spatial distribution of stream segments



—  $\mu$   
—  $\mu - 1$   
- - -  $\mu - 2$

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# Generalizing Tokunaga's law


- Follow stream segments down stream from their beginning
- Probability (or rate) of an order  $\mu$  stream segment terminating is **constant**:

$$\tilde{p}_\mu \simeq 1/(R_s)^{\mu-1} \xi_s$$

- Probability decays exponentially with stream order
- Inter-tributary lengths exponentially distributed
- $\Rightarrow$  random spatial distribution of stream segments





# Generalizing Tokunaga's law


 Joint distribution for generalized version of Tokunaga's law:


$$P(s_\mu, T_{\mu,\nu}) = \tilde{p}_\mu \binom{s_\mu - 1}{T_{\mu,\nu}} p_\nu^{T_{\mu,\nu}} (1 - p_\nu - \tilde{p}_\mu)^{s_\mu - T_{\mu,\nu} - 1}$$

where

  $p_\nu$  = probability of absorbing an order  $\nu$  side stream

  $\tilde{p}_\mu$  = probability of an order  $\mu$  stream terminating

 Approximation: depends on distance units of  $s_\mu$

 In each unit of distance along stream, there is one chance of a side stream entering or the stream terminating.

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations


Models

Nutshell


References




# Generalizing Tokunaga's law

 Now deal with this thing:

$$P(s_\mu, T_{\mu,\nu}) = \tilde{p}_\mu \binom{s_\mu - 1}{T_{\mu,\nu}} p_\nu^{T_{\mu,\nu}} (1 - p_\nu - \tilde{p}_\mu)^{s_\mu - T_{\mu,\nu} - 1}$$

 Set  $(x, y) = (s_\mu, T_{\mu,\nu})$  and  $q = 1 - p_\nu - \tilde{p}_\mu$ , approximate liberally.

 Obtain

$$P(x, y) = Nx^{-1/2} [F(y/x)]^x$$

where

$$F(v) = \left(\frac{1-v}{q}\right)^{-(1-v)} \left(\frac{v}{p}\right)^{-v}.$$

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell



References



# Generalizing Tokunaga's law

☒ Checking form of  $P(s_\mu, T_{\mu,\nu})$  works:

Scheidegger:



Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell



References



# Generalizing Tokunaga's law

☒ Checking form of  $P(s_\mu, T_{\mu,\nu})$  works:

Scheidegger:




Horton  $\leftrightarrow$   
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



# Generalizing Tokunaga's law

☒ Checking form of  $P(s_\mu, T_{\mu,\nu})$  works:

Scheidegger:



Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell


References



# Generalizing Tokunaga's law

Checking form of  $P(s_\mu, T_{\mu,\nu})$  works:


Mississippi:







## Random subnetworks on a Bethe lattice <sup>[13]</sup>



- ❏ Dominant theoretical concept for several decades.
- ❏ Bethe lattices are fun and tractable.
- ❏ Led to idea of “Statistical inevitability” of river network statistics <sup>[7]</sup>
- ❏ But Bethe lattices unconnected with surfaces.
- ❏ In fact, Bethe lattices  $\simeq$  infinite dimensional spaces (oops).
- ❏ So let's move on ...

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models


Nutshell

References



# Scheidegger's model

## Directed random networks [11, 12]



$$P(\searrow) = P(\swarrow) = 1/2$$



Functional form of all scaling laws exhibited but exponents differ from real world [15, 16, 14]

- Horton  $\leftrightarrow$  Tokunaga
- Reducing Horton
- Scaling relations
- Fluctuations
- Models
- Nutshell
- References



Rodríguez-Iturbe, Rinaldo, et al. [10]

- Landscapes  $h(\vec{x})$  evolve such that energy dissipation  $\dot{\epsilon}$  is minimized, where

$$\dot{\epsilon} \propto \int d\vec{r} \text{ (flux)} \times \text{ (force)} \sim \sum_i a_i \nabla h_i \sim \sum_i a_i^\gamma$$

- Landscapes obtained numerically give exponents near that of real networks.
- But:** numerical method used matters.
- And:** Maritan et al. find basic universality classes are that of Scheidegger, self-similar, and a third kind of random network [8]

Horton  $\leftrightarrow$   
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



## Summary of universality classes:

network	h	d
Non-convergent flow	1	1
Directed random	2/3	1
Undirected random	5/8	5/4
Self-similar	1/2	1
OCN's (I)	1/2	1
OCN's (II)	2/3	1
OCN's (III)	3/5	1
Real rivers	0.5-0.7	1.0-1.2

$h \Rightarrow \ell \propto a^h$  (Hack's law).

$d \Rightarrow \ell \propto L_{\parallel}^d$  (stream self-affinity).

Horton  $\Leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



## Branching networks II Key Points:

- 🧱 Horton's laws and Tokunaga's law all fit together.
- 🧱 For 2-d networks, these laws are 'planform' laws and ignore slope.
- 🧱 Abundant scaling relations can be derived.
- 🧱 Can take  $R_n$ ,  $R_\ell$ , and  $d$  as three independent parameters necessary to describe all 2-d branching networks.
- 🧱 For scaling laws, only  $h = \ln R_\ell / \ln R_n$  and  $d$  are needed.
- 🧱 Laws can be extended nicely to laws of distributions.
- 🧱 Numerous models of branching network evolution exist: nothing rock solid yet ...?

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# References I

PoCS  
@pocsvox

Branching  
Networks II

- [1] H. de Vries, T. Becker, and B. Eckhardt.  
Power law distribution of discharge in ideal  
networks.  
[Water Resources Research, 30\(12\):3541–3543,  
1994. pdf](#)
- [2] P. S. Dodds and D. H. Rothman.  
Unified view of scaling laws for river networks.  
[Physical Review E, 59\(5\):4865–4877, 1999. pdf](#)
- [3] P. S. Dodds and D. H. Rothman.  
Geometry of river networks. II. Distributions of  
component size and number.  
[Physical Review E, 63\(1\):016116, 2001. pdf](#)

Horton ⇔  
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References



# References II



- [4] P. S. Dodds and D. H. Rothman.  
Geometry of river networks. III. Characterization  
of component connectivity.  
[Physical Review E, 63\(1\):016117, 2001. pdf](#)
- [5] N. Goldenfeld.  
Lectures on Phase Transitions and the  
Renormalization Group, volume 85 of Frontiers in  
Physics.  
Addison-Wesley, Reading, Massachusetts, 1992.
- [6] J. T. Hack.  
Studies of longitudinal stream profiles in Virginia  
and Maryland.  
[United States Geological Survey Professional  
Paper, 294-B:45-97, 1957. pdf](#)

Horton ↔  
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References






# References III

- [7] J. W. Kirchner.  
Statistical inevitability of Horton's laws and the  
apparent randomness of stream channel  
networks.  
[Geology](#), 21:591–594, 1993. [pdf](#) 
- [8] A. Maritan, F. Colaiori, A. Flammini, M. Cieplak,  
and J. R. Banavar.  
Universality classes of optimal channel networks.  
[Science](#), 272:984–986, 1996. [pdf](#) 
- [9] S. D. Peckham.  
New results for self-similar trees with applications  
to river networks.  
[Water Resources Research](#), 31(4):1023–1029,  
1995.

Horton  $\leftrightarrow$   
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



# References IV

- [10] I. Rodríguez-Iturbe and A. Rinaldo.  
Fractal River Basins: Chance and Self-Organization.  
Cambridge University Press, Cambridge, UK,  
1997.
- [11] A. E. Scheidegger.  
A stochastic model for drainage patterns into an  
intramontane trench.  
Bull. Int. Assoc. Sci. Hydrol., 12(1):15–20, 1967.  
[pdf](#) 
- [12] A. E. Scheidegger.  
Theoretical Geomorphology.  
Springer-Verlag, New York, third edition, 1991.



# References V

- [13] R. L. Shreve.  
Infinite topologically random channel networks.  
[Journal of Geology, 75:178–186, 1967.](#) pdf ↗
- [14] H. Takayasu.  
Steady-state distribution of generalized  
aggregation system with injection.  
[Physical Review Letters, 63\(23\):2563–2565, 1989.](#)  
pdf ↗
- [15] H. Takayasu, I. Nishikawa, and H. Tasaki.  
Power-law mass distribution of aggregation  
systems with injection.  
[Physical Review A, 37\(8\):3110–3117, 1988.](#)

Horton ⇔  
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



# References VI

- [16] M. Takayasu and H. Takayasu.  
Apparent independency of an aggregation system  
with injection.  
[Physical Review A, 39\(8\):4345–4347, 1989. pdf](#)
- [17] D. G. Tarboton, R. L. Bras, and I. Rodríguez-Iturbe.  
Comment on “On the fractal dimension of stream  
networks” by Paolo La Barbera and Renzo Rosso.  
[Water Resources Research, 26\(9\):2243–4, 1990.](#)  
[pdf](#)
- [18] E. Tokunaga.  
The composition of drainage network in Toyohira  
River Basin and the valuation of Horton’s first law.  
[Geophysical Bulletin of Hokkaido University,](#)  
[15:1–19, 1966. pdf](#)

Horton ↔  
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



## References VII

- [19] E. Tokunaga.  
Consideration on the composition of drainage  
networks and their evolution.  
[Geographical Reports of Tokyo Metropolitan  
University, 13:G1-27, 1978. pdf](#)
- [20] E. Tokunaga.  
Ordering of divide segments and law of divide  
segment numbers.  
[Transactions of the Japanese Geomorphological  
Union, 5\(2\):71-77, 1984.](#)
- [21] S. D. Willett, S. W. McCoy, J. T. Perron, L. Goren,  
and C.-Y. Chen.  
Dynamic reorganization of river basins.  
[Science, 343\(6175\):1248765, 2014. pdf](#)

Horton ↔  
Tokunaga  
Reducing Horton  
Scaling relations  
Fluctuations  
Models  
Nutshell  
References



- [22] G. K. Zipf.  
Human Behaviour and the Principle of  
Least-Effort.  
Addison-Wesley, Cambridge, MA, 1949.

Horton  $\leftrightarrow$   
Tokunaga

Reducing Horton

Scaling relations

Fluctuations

Models

Nutshell

References

