

# Mechanisms for Generating Power-Law Size Distributions, Part 1

Last updated: 2022/08/27, 23:54:10 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D  
 CSYS/MATH 300, 303, & 394, 2022–2023 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center  
 Santa Fe Institute | University of Vermont



Licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License.

## Outline

Random Walks

The First Return Problem

Random River Networks

Scaling Relations

Death and Sports

Fractional Brownian Motion

References

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



1 of 46

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References




2 of 46

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



5 of 46



## Mechanisms:

A powerful story in the rise of complexity:


- structure arises out of randomness.
- Exhibit A: Random walks

The essential random walk:

- One spatial dimension.
- Time and space are discrete
- Random walker (e.g., a zombie texter) starts at origin  $x = 0$ .
- Step at time  $t$  is  $\epsilon_t$ :

$$\epsilon_t = \begin{cases} +1 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/2 \end{cases}$$

A few random random walks:



PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



6 of 46

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



7 of 46

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



8 of 46

## Random walks:

Displacement after  $t$  steps:

$$x_t = \sum_{i=1}^t \epsilon_i$$

Expected displacement:

$$\langle x_t \rangle = \left\langle \sum_{i=1}^t \epsilon_i \right\rangle = \sum_{i=1}^t \langle \epsilon_i \rangle = 0$$

- At any time step, we 'expect' our zombie texter to be back at their starting place.
- Obviously fails for odd number of steps...
- But as time goes on, the chance of our texting undead friend lurching back to  $x=0$  must diminish, right?

Variances sum:

$$\text{Var}(x_t) = \text{Var}\left(\sum_{i=1}^t \epsilon_i\right) = \sum_{i=1}^t \text{Var}(\epsilon_i) = \sum_{i=1}^t 1 = t$$

\* Sum rule = a good reason for using the variance to measure spread; only works for independent distributions.

So typical displacement from the origin scales as:

$$\sigma = t^{1/2}$$

- A non-trivial scaling law arises out of additive aggregation or accumulation.

## Random walk basics:

Counting random walks:

- Each specific random walk of length  $t$  appears with a chance  $1/2^t$ .
- We'll be more interested in how many random walks end up at the same place.
- Define  $N(i, j, t)$  as # distinct walks that start at  $x = i$  and end at  $x = j$  after  $t$  time steps.
- Random walk must displace by  $+(j - i)$  after  $t$  steps.

Insert question from assignment 5

$$N(i, j, t) = \binom{t}{(t+j-i)/2}$$

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



9 of 46

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References




10 of 46

PoCS  
 @pocsvox  
 Power-Law  
 Mechanisms, Pt. 1

Random Walks  
 The First Return  
 Problem  
 Random River  
 Networks  
 Scaling Relations  
 Death and Sports  
 Fractional  
 Brownian Motion  
 References



12 of 46



## How does $P(x_t)$ behave for large $t$ ?

- Take time  $t = 2n$  to help ourselves.
- $x_{2n} \in \{0, \pm 2, \pm 4, \dots, \pm 2n\}$
- $x_{2n}$  is even so set  $x_{2n} = 2k$ .
- Using our expression  $N(i, j, t)$  with  $i = 0, j = 2k$ , and  $t = 2n$ , we have

$$\Pr(x_{2n} \equiv 2k) \propto \binom{2n}{n+k}$$

- For large  $n$ , the binomial deliciously approaches the Normal Distribution of Snoredom:

$$\Pr(x_t \equiv x) \simeq \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}}$$

Insert question from assignment 5

- The whole is different from the parts. #nutritious
- See also: [Stable Distributions](#)

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

13 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1


Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

14 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

15 of 46



PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

16 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

17 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

18 of 46

## Applied knot theory:



"Designing tie knots by random walks"  
Fink and Mao,  
Nature, **398**, 31-32, 1999. [6]




Figure 1 All diagrams are drawn in the frame of reference of the mirror image of the actual tie. a. The two ways of beginning a knot, L and R. For knots beginning with L, the tie must begin inside-out. b. The four-in-hand, denoted by the sequence L<sub>2</sub>R<sub>2</sub>L<sub>2</sub>C<sub>2</sub>T. c. A knot may be represented by a persistent random walk on a triangular lattice. The example shown is the four-in-hand, indicated by the walk 111a.

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

19 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References


20 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1


Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

21 of 46

## Universality is also not left-handed:



- This is Diffusion: the most essential kind of spreading (more later).
- View as Random Additive Growth Mechanism.



PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

14 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

15 of 46

## Applied knot theory:

h	γ	γ/h	K(h, γ)	s	b	Name	Sequence
3	1	0.33	1	0	0		L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> T
4	1	0.25	1	-1	1	Four-in-hand	L <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> T
5	2	0.40	2	-1	0	Pratt knot	L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> T
6	2	0.33	4	0	0	Half-Windsor	L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> T
7	2	0.29	6	-1	1		L <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> T
7	3	0.43	4	0	1		L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> C <sub>2</sub> L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> T
8	2	0.25	8	0	2		L <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> T
8	3	0.38	12	-1	0	Windsor	L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> T
9	3	0.33	24	0	0		L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> L <sub>2</sub> R <sub>2</sub> C <sub>2</sub> T
9	4	0.44	8	-1	2		L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> C <sub>2</sub> L <sub>2</sub> C <sub>2</sub> R <sub>2</sub> L <sub>2</sub> C <sub>2</sub> T

- $h$  = number of moves
- $\gamma$  = number of center moves
- $K(h, \gamma) = \frac{2^{\gamma-1} (h-\gamma-2)}{\gamma-1}$
- $s = \sum_{i=1}^h x_i$  where  $x = -1$  for L and +1 for R.
- $b = \frac{1}{2} \sum_{i=2}^{h-1} |\omega_i + \omega_{i-1}|$  where  $\omega = \pm 1$  represents winding direction.

## Random walks #crazytownbananapants

### The problem of first return:


- What is the probability that a random walker in one dimension returns to the origin for the first time after  $t$  steps?
- Will our zombie texter always return to the origin?
- What about higher dimensions?

### Reasons for caring:

- We will find a power-law size distribution with an interesting exponent.
- Some physical structures may result from random walks.
- We'll start to see how different scalings relate to each other.

## So many things are connected:

### Pascal's Triangle



- Could have been the Pyramid of Pingala<sup>1</sup> or the Triangle of Khayyam, Jia Xian, Tartaglia, ...

- Binomials tend towards the Normal.
- Counting encoded in algebraic forms (and much more).
- $(h+t)^n = \sum_{k=0}^n \binom{n}{k} h^k t^{n-k}$  where  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$
- $(h+t)^3 = hhh + hht + hth + thh + htt + tht + tth + ttt$

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

15 of 46

## Random walks are even weirder than you might think...

- $\xi_{r,t}$  = the probability that by time step  $t$ , a random walk has crossed the origin  $r$  times.
- Think of a coin flip game with ten thousand tosses.
- If you are behind early on, what are the chances you will make a comeback?
- The most likely number of lead changes is... 0.
- In fact:  $\xi_{0,t} > \xi_{1,t} > \xi_{2,t} > \dots$
- Even crazier:  
The expected time between tied scores =  $\infty$

See Feller, Intro to Probability Theory, Volume I [5]


PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References


18 of 46

<sup>1</sup>Stigler's Law of Eponymy showing excellent form again.

## For random walks in 1-d:



- A **return** to origin can only happen when  $t = 2n$ .
- In example above, returns occur at  $t = 8, 10$ , and  $14$ .
- Call  $P_{fr}(2n)$  the probability of **first return** at  $t = 2n$ .
- Probability calculation  $\equiv$  Counting problem (combinatorics/statistical mechanics).
- Idea:** Transform first return problem into an easier return problem.



- Can assume zombie texter first lurches to  $x = 1$ .
- Observe walk first returning at  $t = 16$  stays at or above  $x = 1$  for  $1 \leq t \leq 15$  (dashed red line).
- Now want walks that can return many times to  $x = 1$ .
- $P_{fr}(2n) = 2 \cdot \frac{1}{2} Pr(x_t \geq 1, 1 \leq t \leq 2n-1, \text{ and } x_1 = x_{2n-1} = 1)$
- The  $\frac{1}{2}$  accounts for  $x_{2n} = 2$  instead of 0.
- The 2 accounts for texters that first lurch to  $x = -1$ .

## Counting first returns:

### Approach:

- Move to counting numbers of walks.
- Return to probability at end.
- Again,  $N(i, j, t)$  is the # of possible walks between  $x = i$  and  $x = j$  taking  $t$  steps.
- Consider **all paths** starting at  $x = 1$  and ending at  $x = 1$  after  $t = 2n - 2$  steps.
- Idea:** If we can compute the number of walks that hit  $x = 0$  at least once, then we can subtract this from the total number to find the ones that maintain  $x \geq 1$ .
- Call walks that drop below  $x = 1$  **excluded walks**.
- We'll use a method of images to identify these excluded walks.

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

22 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References


23 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

24 of 46

## Examples of excluded walks:



### Key observation for excluded walks:

- For any path starting at  $x=1$  that hits 0, there is a unique matching path starting at  $x=-1$ .
- Matching path first mirrors and then tracks after first reaching  $x=0$ .
- # of  $t$ -step paths starting and ending at  $x=1$  and hitting  $x=0$  at least once = # of  $t$ -step paths starting at  $x=-1$  and ending at  $x=1 = N(-1, 1, t)$
- So  $N_{\text{first return}}(2n) = N(1, 1, 2n-2) - N(-1, 1, 2n-2)$

## Probability of first return:

Insert question from assignment 5

Find

$$N_{fr}(2n) \sim \frac{2^{2n-3/2}}{\sqrt{2\pi n^{3/2}}}$$

- Normalized number of paths gives probability.
- Total number of possible paths =  $2^{2n}$ .

$$P_{fr}(2n) = \frac{1}{2^{2n}} N_{fr}(2n) \approx \frac{1}{2^{2n}} \frac{2^{2n-3/2}}{\sqrt{2\pi n^{3/2}}} = \frac{1}{\sqrt{2\pi}} (2n)^{-3/2} \propto t^{-3/2}$$

- We have  $P(t) \propto t^{-3/2}$ ,  $\gamma = 3/2$ .
- Same scaling holds for continuous space/time walks.
- $P(t)$  is normalizable.
- Recurrence:** Random walker always returns to origin
- But mean, variance, and all higher moments are infinite. **#totalmadness**
- Even though walker must return, expect a long wait...
- One moral:** Repeated gambling against an infinitely wealthy opponent must lead to ruin.

### Higher dimensions

- Walker in  $d = 2$  dimensions must also return
- Walker may not return in  $d \geq 3$  dimensions
- Associated human genius: [George Pólya](#)

27 of 46

## Random walks


### On finite spaces:

- In any finite homogeneous space, a random walker will visit every site with equal probability
- Call this probability the **Invariant Density** of a dynamical system
- Non-trivial Invariant Densities arise in chaotic systems.

### On networks:

- On networks, a random walker visits each node with frequency  $\propto$  node degree **#groovy**
- Equal probability still present: walkers traverse **edges** with equal frequency. **#totallygroovy**

## Scheidegger Networks [17, 4]



- Random directed network on triangular lattice.
- Toy model of real networks.
- 'Flow' is southeast or southwest with equal probability.

## Scheidegger networks

- Creates basins with random walk boundaries.
- Observe** that subtracting one random walk from another gives random walk with increments:

$$\epsilon_t = \begin{cases} +1 & \text{with probability } 1/4 \\ 0 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/4 \end{cases}$$

- Random walk with probabilistic pauses.
- Basin termination = first return random walk problem.
- Basin length  $\ell$  distribution:  $P(\ell) \propto \ell^{-3/2}$
- For real river networks, generalize to  $P(\ell) \propto \ell^{-\gamma}$ .

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

25 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

26 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

27 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

28 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

29 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References

30 of 46

## Connections between exponents:

- For a basin of length  $\ell$ , width  $\propto \ell^{1/2}$
- Basin area  $a \propto \ell \cdot \ell^{1/2} = \ell^{3/2}$
- Invert:  $\ell \propto a^{2/3}$
- $d\ell \propto d(a^{2/3}) = 2/3 a^{-1/3} da$
- $\Pr(\text{basin area} = a) da$   
 $= \Pr(\text{basin length} = \ell) d\ell$   
 $\propto \ell^{-3/2} d\ell$   
 $\propto (a^{2/3})^{-3/2} a^{-1/3} da$   
 $= a^{-4/3} da$   
 $= a^{-\tau} da$

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



31 of 46

## Connections between exponents:

- Both basin area and length obey power law distributions
- Observed for real river networks
- Reportedly:  $1.3 < \tau < 1.5$  and  $1.5 < \gamma < 2$

## Generalize relationship between area and length:

- Hack's law<sup>[10]</sup>:

$$\ell \propto a^h.$$

- For real, large networks<sup>[13]</sup>  $h \simeq 0.5$  (isometric scaling)
- Smaller basins possibly  $h > 1/2$  (allometric scaling).
- Models exist with interesting values of  $h$ .
- Plan: Redo calc with  $\gamma$ ,  $\tau$ , and  $h$ .

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



32 of 46

## Connections between exponents:

- Given

$$\ell \propto a^h, P(a) \propto a^{-\tau}, \text{ and } P(\ell) \propto \ell^{-\gamma}$$

- $d\ell \propto d(a^h) = ha^{h-1} da$
- Find  $\tau$  in terms of  $\gamma$  and  $h$ .
- $\Pr(\text{basin area} = a) da$   
 $= \Pr(\text{basin length} = \ell) d\ell$   
 $\propto \ell^{-\gamma} d\ell$   
 $\propto (a^h)^{-\gamma} a^{h-1} da$   
 $= a^{-(1+h(\gamma-1))} da$

$$\tau = 1 + h(\gamma - 1)$$

- Excellent example of the **Scaling Relations** found between exponents describing power laws for many systems.

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



33 of 46

## Connections between exponents:

With more detailed description of network structure,  $\tau = 1 + h(\gamma - 1)$  simplifies to:<sup>[3]</sup>

$$\tau = 2 - h$$

and

$$\gamma = 1/h$$

- Only one exponent is independent (take  $h$ ).
- Simplifies system description.
- Expect Scaling Relations where power laws are found.
- Need only characterize **Universality** class with independent exponents.

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



34 of 46

## Death ...

### Failure:

- A very simple model of failure/death
- $x_t$  = entity's 'health' at time  $t$
- Start with  $x_0 > 0$ .
- Entity fails when  $x$  hits 0.



"Explaining mortality rate plateaus"  
Weitz and Fraser,  
Proc. Natl. Acad. Sci., **98**, 15383-15386,  
2001. <sup>[18]</sup>

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



35 of 46

## ... and the NBA:

### Basketball and other sports<sup>[2]</sup>:

- Three arcsine laws (Lévy<sup>[12]</sup>) for continuous-time random walk last time  $T$ :

$$\frac{1}{\pi} \frac{1}{\sqrt{t(T-t)}}$$

The arcsine distribution applies for:  
(1) fraction of time positive, (2) the last time the walk changes sign, and (3) the time the maximum is achieved.

- Well approximated by basketball score lines<sup>[8, 2]</sup>.
- Australian Rules Football has some differences<sup>[11]</sup>.


## More than randomness

- Can generalize to Fractional Random Walks<sup>[15, 16, 14]</sup>
- Fractional Brownian Motion, Lévy flights
- See Montroll and Shlesinger for example:<sup>[14]</sup>  
"On  $1/f$  noise and other distributions with long tails."  
Proc. Natl. Acad. Sci., 1982.
- In 1-d, standard deviation  $\sigma$  scales as


$$\sigma \sim t^\alpha$$

$\alpha = 1/2$  — diffusive  
 $\alpha > 1/2$  — superdiffusive  
 $\alpha < 1/2$  — subdiffusive

- Extensive memory of path now matters...



- First big studies of movement and interactions of people.
- Brockmann *et al.*<sup>[1]</sup> "Where's George" study.
- Beyond Lévy: Superdiffusive in space but with long waiting times.
- Tracking movement via cell phones<sup>[9]</sup> and Twitter<sup>[7]</sup>.



PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



36 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



37 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References




38 of 46

PoCS  
@pocsvox  
Power-Law  
Mechanisms, Pt. 1

Random Walks  
The First Return  
Problem  
Random River  
Networks  
Scaling Relations  
Death and Sports  
Fractional  
Brownian Motion  
References



39 of 46



## References III

- [8] A. Gabel and S. Redner.  
Random walk picture of basketball scoring.  
[Journal of Quantitative Analysis in Sports](#), 8:1–20, 2012.
- [9] M. C. González, C. A. Hidalgo, and A.-L. Barabási.  
Understanding individual human mobility patterns.  
[Nature](#), 453:779–782, 2008. [pdf](#)
- [10] J. T. Hack.  
Studies of longitudinal stream profiles in Virginia and Maryland.  
[United States Geological Survey Professional Paper](#), 294-B:45–97, 1957. [pdf](#)

## References IV

- [11] D. P. Kiley, A. J. Reagan, L. Mitchell, C. M. Danforth, and P. S. Dodds.  
The game story space of professional sports: Australian Rules Football.  
[Physical Review E](#), 93, 2016. [pdf](#)
- [12] P. Lévy and M. Loeve.  
[Processus stochastiques et mouvement brownien](#).  
Gauthier-Villars Paris, 1965.
- [13] D. R. Montgomery and W. E. Dietrich.  
Channel initiation and the problem of landscape scale.  
[Science](#), 255:826–30, 1992. [pdf](#)

## References V



- [14] E. W. Montroll and M. F. Shlesinger.  
[On the wonderful world of random walks](#), volume XI of [Studies in statistical mechanics](#), chapter 1, pages 1–121.  
New-Holland, New York, 1984.
- [15] E. W. Montroll and M. W. Shlesinger.  
On  $1/f$  noise and other distributions with long tails.  
[Proc. Natl. Acad. Sci.](#), 79:3380–3383, 1982. [pdf](#)
- [16] E. W. Montroll and M. W. Shlesinger.  
Maximum entropy formalism, fractals, scaling phenomena, and  $1/f$  noise: a tale of tails.  
[J. Stat. Phys.](#), 32:209–230, 1983.

## References VI

- [17] A. E. Scheidegger.  
The algebra of stream-order numbers.  
[United States Geological Survey Professional Paper](#), 525-B:B187–B189, 1967. [pdf](#)
- [18] J. S. Weitz and H. B. Fraser.  
Explaining mortality rate plateaus.  
[Proc. Natl. Acad. Sci.](#), 98:15383–15386, 2001. [pdf](#)



## References I

- [1] D. Brockmann, L. Hufnagel, and T. Geisel.  
The scaling laws of human travel.  
[Nature](#), pages 462–465, 2006. [pdf](#)
- [2] A. Clauset, M. Kogan, and S. Redner.  
Safe leads and lead changes in competitive team sports.  
[Phys. Rev. E](#), 91:062815, 2015. [pdf](#)
- [3] P. S. Dodds and D. H. Rothman.  
Unified view of scaling laws for river networks.  
[Physical Review E](#), 59(5):4865–4877, 1999. [pdf](#)
- [4] P. S. Dodds and D. H. Rothman.  
Scaling, universality, and geomorphology.  
[Annu. Rev. Earth Planet. Sci.](#), 28:571–610, 2000. [pdf](#)

## References II

- [5] W. Feller.  
[An Introduction to Probability Theory and Its Applications](#), volume I.  
John Wiley & Sons, New York, third edition, 1968.
- [6] T. M. Fink and Y. Mao.  
Designing tie knots by random walks.  
[Nature](#), 398:31–32, 1999. [pdf](#)
- [7] M. R. Frank, L. Mitchell, P. S. Dodds, and C. M. Danforth.  
Happiness and the patterns of life: A study of geolocated Tweets.  
[Nature Scientific Reports](#), 3:2625, 2013. [pdf](#)