


Mechanisms for Generating Power-Law Size Distributions, Part 2

Last updated: 2023/08/22, 11:48:23 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D
CSYS/MATH 6701, 6713, & a pretend number,
2023–2024 | @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
Santa Fe Institute | University of Vermont



These slides are brought to you by:

Sealie & Lambie
Productions




The PoCSverse
Power-Law
Mechanisms, Pt. 2
2 of 20

Variable
transformation


Basics
Holtmark's Distribution
PLIPLO

References




These slides are also brought to you by:

Special Guest Executive Producer



On Instagram at [pratchett_the_cat](https://www.instagram.com/pratchett_the_cat/)



Outline

The PoCSverse
Power-Law
Mechanisms, Pt. 2
4 of 20

Variable
transformation

Basics
Holtsmark's Distribution
PLIPLO

References


Variable transformation

Basics

Holtsmark's Distribution

PLIPLO

References



Variable
transformation



Basics

Holtsmark's Distribution


PLIPLO

References

The Boggoracle Speaks:



THE UNEXPECTED
TRANSFIGURATION



Variable Transformation

The PoCSverse
Power-Law
Mechanisms, Pt. 2
8 of 20

Variable
transformation

Basics
Holtsmark's Distribution
PLIPLO

References


Understand power laws as arising from

1. Elementary distributions (e.g., exponentials).
2. Variables connected by power relationships.

- Random variable X with known distribution P_x
- Second random variable Y with $y = f(x)$.

$$\begin{aligned} P_Y(y)dy &= \\ \sum_{x|f(x)=y} P_X(x)dx &= \\ \sum_{y|f(x)=y} P_X(f^{-1}(y)) \frac{dy}{|f'(f^{-1}(y))|} \end{aligned}$$

- Often easier to do by hand...



General Example

-Assume relationship between x and y is 1-1.

-Power-law relationship between variables:

$$y = cx^{-\alpha}, \alpha > 0$$

-Look at y large and x small




$$dy = d(cx^{-\alpha})$$

$$= c(-\alpha)x^{-\alpha-1}dx$$

invert: $dx = \frac{-1}{c\alpha}x^{\alpha+1}dy$

$$dx = \frac{-1}{c\alpha} \left(\frac{y}{c}\right)^{-(\alpha+1)/\alpha} dy$$

$$dx = \frac{-c^{1/\alpha}}{\alpha} y^{-1-1/\alpha} dy$$



Now make transformation:

$$P_y(y)dy = P_x(x)dx$$


$$P_y(y)dy = P_x\left(\left(\frac{y}{c}\right)^{-1/\alpha}\right)\overbrace{\frac{c^{1/\alpha}}{\alpha}y^{-1-1/\alpha}dy}^{dx}$$

>If $P_x(x) \rightarrow$ non-zero constant as $x \rightarrow 0$ then

$$P_y(y) \propto y^{-1-1/\alpha} \text{ as } y \rightarrow \infty.$$

If $P_x(x) \rightarrow x^\beta$ as $x \rightarrow 0$ then

$$P_y(y) \propto y^{-1-1/\alpha-\beta/\alpha} \text{ as } y \rightarrow \infty.$$



Example

The PoCSverse
Power-Law
Mechanisms, Pt. 2
11 of 20

Variable
transformation

Basics
Holtsmark's Distribution
PLIPLO


References

Exponential distribution

Given $P_x(x) = \frac{1}{\lambda} e^{-x/\lambda}$ and $y = cx^{-\alpha}$, then

$$P(y) \propto y^{-1-1/\alpha} + O(y^{-1-2/\alpha})$$

- ➊ Exponentials arise from randomness (easy) ...
- ➋ More later when we cover robustness.



Gravity

- Select a random point in the universe \vec{x} .
- Measure the force of gravity $F(\vec{x})$.
- Observe that $P_F(F) \sim F^{-5/2}$.
- Distribution named after Holtsmark who was thinking about electrostatics and plasma [1].
- Again, the humans naming things after humans, poorly.¹



¹Stigler's Law of Eponomy ↗

Matter is concentrated in stars: [2]

- ⬢ F is distributed unevenly
- ⬢ Probability of being a distance r from a single star at $\vec{x} = \vec{0}$:

$$P_r(r)dr \propto r^2 dr$$

- ⬢ Assume stars are distributed randomly in space (oops?)
- ⬢ Assume only one star has significant effect at \vec{x} .
- ⬢ Law of gravity:

$$F \propto r^{-2}$$

- ⬢ invert:

$$r \propto F^{-\frac{1}{2}}$$

- ⬢ Connect differentials: $dr \propto dF^{-\frac{1}{2}} \propto F^{-\frac{3}{2}} dF$



Transformation:

Using $r \propto F^{-1/2}$, $dr \propto F^{-3/2}dF$, and $P_r(r) \propto r^2$



$$P_F(F)dF = P_r(r)dr$$



$$\propto P_r(\text{const} \times F^{-1/2})F^{-3/2}dF$$



$$\propto (F^{-1/2})^2 F^{-3/2}dF$$



$$= F^{-1-3/2}dF$$



$$= F^{-5/2}dF.$$



Gravity:

The PoCSverse
Power-Law
Mechanisms, Pt. 2
16 of 20

Variable
transformation

Basics
Holtmark's Distribution
PLIPLO

References

$$P_F(F) = F^{-5/2} dF$$




$$\gamma = 5/2$$

- Mean is finite.
- Variance = ∞ .
- A **wild** distribution.
- Upshot:** Random sampling of space usually safe but can end badly...



Todo: Build Dalek army.



Extreme Caution!

The PoCSverse
Power-Law
Mechanisms, Pt. 2
19 of 20

Variable
transformation
Basics
Holtsmark's Distribution
PLIPLO

References

- ⬢ PLIPLO = Power law in, power law out
- ⬢ Explain a power law as resulting from another unexplained power law.
- ⬢ Yet another homunculus argument ↗...
- ⬢ Don't do this!!! (slap, slap)
- ⬢ MIWO = Mild in, Wild out is the stuff.
- ⬢ In general: We need mechanisms!



References I

References

- [1] J. Holtsmark.
Über die verbreiterung von spektrallinien.
Ann. Phys., 58:577–630, 1919. pdf ↗
- [2] D. Sornette.
Critical Phenomena in Natural Sciences.
Springer-Verlag, Berlin, 1st edition, 2003.

