

Mechanisms for Generating Power-Law Size Distributions, Part 1

Last updated: 2023/08/22, 11:48:21 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D
 CSYS/MATH 6701, 6713, & a pretend number,
 2023–2024 | @pocsvs


Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
 Santa Fe Institute | University of Vermont



Licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License.

- The PoCSverse Power-Law Mechanisms, Pt. 1 1 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References



- The PoCSverse Power-Law Mechanisms, Pt. 1 5 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

Random walks:

Displacement after t steps:

$$x_t = \sum_{i=1}^t \epsilon_i$$

Expected displacement:

$$\langle x_t \rangle = \left\langle \sum_{i=1}^t \epsilon_i \right\rangle = \sum_{i=1}^t \langle \epsilon_i \rangle = 0$$

- ☞ At any time step, we 'expect' our zombie texter to be back at their starting place.
- ☞ Obviously fails for odd number of steps...
- ☞ But as time goes on, the chance of our texting undead friend lurching back to $x=0$ must diminish, right?

- The PoCSverse Power-Law Mechanisms, Pt. 1 8 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

Outline

- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

- The PoCSverse Power-Law Mechanisms, Pt. 1 2 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

Mechanisms:

A powerful story in the rise of complexity:

- ☞ structure arises out of randomness.
- ☞ Exhibit A: Random walks

The essential random walk:

- ☞ One spatial dimension.
- ☞ Time and space are discrete
- ☞ Random walker (e.g., a zombie texter) starts at origin $x = 0$.
- ☞ Step at time t is ϵ_t :

$$\epsilon_t = \begin{cases} +1 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/2 \end{cases}$$

- The PoCSverse Power-Law Mechanisms, Pt. 1 6 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

Variance sum: \square^*

$$\begin{aligned} \text{Var}(x_t) &= \text{Var}\left(\sum_{i=1}^t \epsilon_i\right) \\ &= \sum_{i=1}^t \text{Var}(\epsilon_i) = \sum_{i=1}^t 1 = t \end{aligned}$$

* Sum rule = a good reason for using the variance to measure spread; only works for independent distributions.

So typical displacement from the origin scales as:

$$\sigma = t^{1/2}$$


- ☞ A non-trivial scaling law arises out of additive aggregation or accumulation.

- The PoCSverse Power-Law Mechanisms, Pt. 1 9 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References



- The PoCSverse Power-Law Mechanisms, Pt. 1 4 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

A few random random walks:



- The PoCSverse Power-Law Mechanisms, Pt. 1 7 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

Random walk basics:

Counting random walks:

- ☞ Each **specific** random walk of length t appears with a chance $1/2^t$.
- ☞ We'll be more interested in how many random walks end up at the same place.
- ☞ Define $N(i, j, t)$ as # distinct walks that start at $x = i$ and end at $x = j$ after t time steps.
- ☞ Random walk must displace by $+(j - i)$ after t steps.
- ☞ Insert assignment question \square

$$N(i, j, t) = \binom{t}{(t + j - i)/2}$$

- The PoCSverse Power-Law Mechanisms, Pt. 1 11 of 45
- Random Walks
- The First Return Problem
- Random River Networks
- Scaling Relations
- Death and Sports
- Fractional Brownian Motion
- References

How does $P(x_t)$ behave for large t ?

- Take time $t = 2n$ to help ourselves.
- $x_{2n} \in \{0, \pm 2, \pm 4, \dots, \pm 2n\}$
- x_{2n} is even so set $x_{2n} = 2k$.
- Using our expression $N(i, j, t)$ with $i = 0, j = 2k$, and $t = 2n$, we have

$$\Pr(x_{2n} \equiv 2k) \propto \binom{2n}{n+k}$$


- For large n , the binomial deliciously approaches the Normal Distribution of Snoredom:

$$\Pr(x_t \equiv x) \approx \frac{1}{\sqrt{2\pi t}} e^{-\frac{x^2}{2t}}$$

Insert assignment question

- The whole is different from the parts. #nutritious
- See also: [Stable Distributions](#)


Universality is also not left-handed:



- This is [Diffusion](#): the most essential kind of spreading (more later).
- View as Random Additive Growth Mechanism.

So many things are connected:

Pascal's Triangle




- Could have been the Pyramid of Pingala¹ or the Triangle of Khayyam, Jia Xian, Tartaglia, ...


- Binomials tend towards the Normal.
- Counting encoded in algebraic forms (and much more).
- $(h+t)^n = \sum_{k=0}^n \binom{n}{k} h^k t^{n-k}$ where $\binom{n}{k} = \frac{n!}{k!(n-k)!}$
- $(h+t)^3 = hhh + hht + hth + thh + htt + tht + tth + ttt$

¹Stigler's Law of Eponymy showing excellent form again.

The PoCServe Power-Law Mechanisms, Pt. 1 12 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References



The PoCServe Power-Law Mechanisms, Pt. 1 13 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References



The PoCServe Power-Law Mechanisms, Pt. 1 14 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References

Random walks are even weirder than you might think...

- $\xi_{r,t}$ = the probability that by time step t , a random walk has crossed the origin r times.
- Think of a coin flip game with ten thousand tosses.
- If you are behind early on, what are the chances you will make a comeback?
- The most likely number of lead changes is... 0.
- In fact: $\xi_{0,t} > \xi_{1,t} > \xi_{2,t} > \dots$
- Even crazier:
The expected time between tied scores = ∞

See Feller, Intro to Probability Theory, Volume I [5]

The PoCServe Power-Law Mechanisms, Pt. 1 15 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References

Applied knot theory:



"Designing tie knots by random walks" Fink and Mao, Nature, 398, 31-32, 1999. [6]




Figure 1 All diagrams are drawn in the frame of reference of the mirror image of the actual tie. a. The two ways of beginning a knot, L_0 and R_0 . For knots beginning with L_0 , the tie must begin inside-out. b. The four-in-hand, denoted by the sequence $L_0 R_0 L_0 C_0 T$. c. A knot may be represented by a persistent random walk on a triangular lattice. The example shown is the four-in-hand, indicated by the walk $111a$.

The PoCServe Power-Law Mechanisms, Pt. 1 16 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References

Applied knot theory:

h	γ	γ/h	$K(h, \gamma)$	s	b	Name	Sequence
3	1	0.33	1	0	0		$L_0 R_0 C_0 T$
4	1	0.25	1	-1	1	Four-in-hand	$L_0 R_0 L_0 C_0 T$
5	2	0.40	2	-1	0	Pratt knot	$L_0 C_0 R_0 L_0 C_0 T$
6	2	0.33	4	0	0		$L_0 R_0 C_0 L_0 R_0 C_0 T$
7	2	0.29	6	-1	1	Half-Windsor	$L_0 R_0 L_0 C_0 R_0 L_0 C_0 T$
7	3	0.43	4	0	1		$L_0 C_0 R_0 C_0 L_0 R_0 C_0 T$
8	2	0.25	8	0	2		$L_0 R_0 L_0 C_0 R_0 L_0 C_0 T$
8	3	0.38	12	-1	0	Windsor	$L_0 C_0 R_0 L_0 C_0 R_0 L_0 C_0 T$
9	3	0.33	24	0	0		$L_0 R_0 C_0 L_0 R_0 C_0 L_0 R_0 C_0 T$
9	4	0.44	8	-1	2		$L_0 C_0 R_0 C_0 L_0 C_0 R_0 L_0 C_0 T$

Knots are characterized by half-winding number h , centre number γ , centre fraction γ/h , knots per class $K(h, \gamma)$, symmetry s , balance b , name and sequence.

- h = number of moves
- γ = number of center moves
- $K(h, \gamma) = \frac{2^{\gamma-1} (h-\gamma-2)}{\gamma-1}$
- $s = \sum_{i=1}^h x_i$ where $x = -1$ for L and $+1$ for R .
- $b = \frac{1}{2} \sum_{i=2}^{h-1} |\omega_i + \omega_{i-1}|$ where $\omega = \pm 1$ represents winding direction.

Random walks #crazytownbananapants

The problem of first return:

- What is the probability that a random walker in one dimension returns to the origin for the first time after t steps?
- Will our zombie texter always return to the origin?
- What about higher dimensions?

Reasons for caring:


- We will find a power-law size distribution with an interesting exponent.
- Some physical structures may result from random walks.
- We'll start to see how different scalings relate to each other.

The PoCServe Power-Law Mechanisms, Pt. 1 18 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References


The PoCServe Power-Law Mechanisms, Pt. 1 19 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References

The PoCServe Power-Law Mechanisms, Pt. 1 20 of 45
 Random Walks
 The First Return Problem
 Random River Networks
 Scaling Relations
 Death and Sports
 Fractional Brownian Motion
 References

For random walks in 1-d:



- A **return** to origin can only happen when $t = 2n$.
- In example above, returns occur at $t = 8, 10,$ and 14 .
- Call $P_{fr}(2n)$ the probability of **first return** at $t = 2n$.
- Probability calculation \equiv Counting problem (combinatorics/statistical mechanics).
- Idea:** Transform first return problem into an easier return problem.




- Can assume zombie texter first lurches to $x = 1$.
- Observe walk first returning at $t = 16$ stays at or above $x = 1$ for $1 \leq t \leq 15$ (dashed red line).
- Now want walks that can return many times to $x = 1$.
- $P_{fr}(2n) = 2 \cdot \frac{1}{2} Pr(x_t \geq 1, 1 \leq t \leq 2n - 1, \text{ and } x_1 = x_{2n-1} = 1)$
- The $\frac{1}{2}$ accounts for $x_{2n} = 2$ instead of 0.
- The 2 accounts for texters that first lurch to $x = -1$.

Counting first returns:

Approach:

- Move to counting numbers of walks.
- Return to probability at end.
- Again, $N(i, j, t)$ is the # of possible walks between $x = i$ and $x = j$ taking t steps.
- Consider **all paths** starting at $x = 1$ and ending at $x = 1$ after $t = 2n - 2$ steps.
- Idea:** If we can compute the number of walks that hit $x = 0$ at least once, then we can subtract this from the total number to find the ones that maintain $x \geq 1$.
- Call walks that drop below $x = 1$ **excluded walks**.
- We'll use a method of images to identify these excluded walks.

Examples of excluded walks:



Key observation for excluded walks:

- For any path starting at $x=1$ that hits 0, there is a unique matching path starting at $x=-1$.
- Matching path first mirrors and then tracks after first reaching $x=0$.
- # of t -step paths starting and ending at $x=1$ and hitting $x=0$ at least once = # of t -step paths starting at $x=-1$ and ending at $x=1 = N(-1, 1, t)$
- So $N_{\text{first return}}(2n) = N(1, 1, 2n - 2) - N(-1, 1, 2n - 2)$

Probability of first return:

Insert assignment question [↗](#):

- Find

$$N_{fr}(2n) \sim \frac{2^{2n-3/2}}{\sqrt{2\pi n^{3/2}}}$$

- Normalized number of paths gives probability.
- Total number of possible paths = 2^{2n} .

$$P_{fr}(2n) = \frac{1}{2^{2n}} N_{fr}(2n) \approx \frac{1}{2^{2n}} \frac{2^{2n-3/2}}{\sqrt{2\pi n^{3/2}}} = \frac{1}{\sqrt{2\pi}} (2n)^{-3/2} \propto t^{-3/2}$$

- We have $P(t) \propto t^{-3/2}$, $\gamma = 3/2$.
- Same scaling holds for continuous space/time walks.
- $P(t)$ is normalizable.
- Recurrence:** Random walker always returns to origin
- But mean, variance, and all higher moments are infinite. **#totalmadness**
- Even though walker must return, expect a long wait...
- One moral:** Repeated gambling against an infinitely wealthy opponent must lead to ruin.

Higher dimensions [↗](#):

- Walker in $d = 2$ dimensions must also return
- Walker may not return in $d \geq 3$ dimensions
- Associated human **genius:** [George Pólya](#) [↗](#)

Random walks


On finite spaces:

- In any finite homogeneous space, a random walker will visit every site with equal probability
- Call this probability the **Invariant Density** of a dynamical system
- Non-trivial Invariant Densities arise in chaotic systems.

On networks:

- On networks, a random walker visits each node with frequency \propto node degree **#groovy**
- Equal probability still present: walkers traverse **edges** with equal frequency. **#totallygroovy**

Scheidegger Networks ^[17, 4]



- Random directed network on triangular lattice.
- Toy model of real networks.
- 'Flow' is southeast or southwest with equal probability.

Scheidegger networks

- Creates basins with random walk boundaries.
- Observe** that subtracting one random walk from another gives random walk with increments:

$$\epsilon_t = \begin{cases} +1 & \text{with probability } 1/4 \\ 0 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/4 \end{cases}$$

- Random walk with probabilistic pauses.
- Basin termination = first return random walk problem.
- Basin length ℓ distribution: $P(\ell) \propto \ell^{-3/2}$
- For real river networks, generalize to $P(\ell) \propto \ell^{-\gamma}$.

Connections between exponents:

- For a basin of length ℓ , width $\propto \ell^{1/2}$
- Basin area $a \propto \ell \cdot \ell^{1/2} = \ell^{3/2}$
- Invert: $\ell \propto a^{2/3}$
- $d\ell \propto d(a^{2/3}) = 2/3 a^{-1/3} da$
- Pr**(basin area = a) da
 $= \mathbf{Pr}$ (basin length = ℓ) $d\ell$
 $\propto \ell^{-3/2} d\ell$
 $\propto (a^{2/3})^{-3/2} a^{-1/3} da$
 $= a^{-4/3} da$
 $= a^{-\tau} da$

The PoCSverse
Power-Law
Mechanisms, Pt. 1
30 of 45
Random Walks
The First Return
Problem
Random River
Networks
[Scaling Relations](#)
Death and Sports
Fractional
Brownian Motion
References

Connections between exponents:

With more detailed description of network structure, $\tau = 1 + h(\gamma - 1)$ simplifies to: [3]

$$\tau = 2 - h$$

and

$$\gamma = 1/h$$

- Only one exponent is independent (take h).
- Simplifies system description.
- Expect Scaling Relations where power laws are found.
- Need only characterize [Universality](#) class with independent exponents.

The PoCSverse
Power-Law
Mechanisms, Pt. 1
33 of 45
Random Walks
The First Return
Problem
Random River
Networks
[Scaling Relations](#)
Death and Sports
Fractional
Brownian Motion
References

More than randomness

Can generalize to Fractional Random Walks [15, 16, 14]

[Fractional Brownian Motion](#), [Lévy flights](#)

See Montroll and Shlesinger for example: [14]

“On $1/f$ noise and other distributions with long tails.”
Proc. Natl. Acad. Sci., 1982.

In 1-d, standard deviation σ scales as


$$\sigma \sim t^\alpha$$

$\alpha = 1/2$ — diffusive

$\alpha > 1/2$ — superdiffusive

$\alpha < 1/2$ — subdiffusive

Extensive memory of path now matters...



The PoCSverse
Power-Law
Mechanisms, Pt. 1
36 of 45
Random Walks
The First Return
Problem
Random River
Networks
Scaling Relations
Death and Sports
[Fractional Brownian Motion](#)
References

Connections between exponents:

- Both basin area and length obey power law distributions
- Observed for real river networks
- Reportedly: $1.3 < \tau < 1.5$ and $1.5 < \gamma < 2$

Generalize relationship between area and length:

Hack's law [10]:

$$\ell \propto a^h.$$

- For real, large networks [13] $h \simeq 0.5$ (isometric scaling)
- Smaller basins possibly $h > 1/2$ (allometric scaling).
- Models exist with interesting values of h .
- Plan:** Redo calc with γ , τ , and h .

The PoCSverse
Power-Law
Mechanisms, Pt. 1
31 of 45
Random Walks
The First Return
Problem
Random River
Networks
[Scaling Relations](#)
Death and Sports
Fractional
Brownian Motion
References

Death ...

Failure:

- A very simple model of failure/death
- x_t = entity's 'health' at time t
- Start with $x_0 > 0$.
- Entity fails when x hits 0.



“Explaining mortality rate plateaus”
Weitz and Fraser,
Proc. Natl. Acad. Sci., **98**, 15383–15386,
2001. [18]

The PoCSverse
Power-Law
Mechanisms, Pt. 1
34 of 45
Random Walks
The First Return
Problem
Random River
Networks
Scaling Relations
[Death and Sports](#)
Fractional
Brownian Motion
References

Connections between exponents:

Given

$$\ell \propto a^h, P(a) \propto a^{-\tau}, \text{ and } P(\ell) \propto \ell^{-\gamma}$$

- $d\ell \propto d(a^h) = ha^{h-1} da$
- Find τ in terms of γ and h .
- Pr**(basin area = a) da
 $= \mathbf{Pr}$ (basin length = ℓ) $d\ell$
 $\propto \ell^{-\gamma} d\ell$
 $\propto (a^h)^{-\gamma} a^{h-1} da$
 $= a^{-(1+h(\gamma-1))} da$

$$\tau = 1 + h(\gamma - 1)$$

- Excellent example of the [Scaling Relations](#) found between exponents describing power laws for many systems.

The PoCSverse
Power-Law
Mechanisms, Pt. 1
32 of 45
Random Walks
The First Return
Problem
Random River
Networks
[Scaling Relations](#)
Death and Sports
Fractional
Brownian Motion
References

... and the NBA:

Basketball and other sports [2]:

Three arcsine laws (Lévy [12]) for continuous-time random walk last time T :


$$\frac{1}{\pi} \frac{1}{\sqrt{t(T-t)}}$$

The arcsine distribution applies for:


- (1) fraction of time positive,
- (2) the last time the walk changes sign,
- and (3) the time the maximum is achieved.

- Well approximated by basketball score lines [8, 2].
- Australian Rules Football has some differences [11].

The PoCSverse
Power-Law
Mechanisms, Pt. 1
35 of 45
Random Walks
The First Return
Problem
Random River
Networks
Scaling Relations
[Death and Sports](#)
Fractional
Brownian Motion
References



The PoCSverse
Power-Law
Mechanisms, Pt. 1
38 of 45
Random Walks
The First Return
Problem
Random River
Networks
Scaling Relations
Death and Sports
[Fractional Brownian Motion](#)
References



References III

- [8] A. Gabel and S. Redner.
Random walk picture of basketball scoring.
[Journal of Quantitative Analysis in Sports](#), 8:1–20, 2012.
- [9] M. C. González, C. A. Hidalgo, and A.-L. Barabási.
Understanding individual human mobility patterns.
[Nature](#), 453:779–782, 2008. [pdf](#)
- [10] J. T. Hack.
Studies of longitudinal stream profiles in Virginia and Maryland.
[United States Geological Survey Professional Paper](#), 294-B:45–97, 1957. [pdf](#)

References V

- [14] E. W. Montroll and M. F. Shlesinger.
On the wonderful world of random walks,
volume XI of Studies in statistical mechanics,
chapter 1, pages 1–121.
New-Holland, New York, 1984.
- [15] E. W. Montroll and M. W. Shlesinger.
On $1/f$ noise and other distributions with long tails.
[Proc. Natl. Acad. Sci.](#), 79:3380–3383, 1982. [pdf](#)
- [16] E. W. Montroll and M. W. Shlesinger.
Maximum entropy formalism, fractals, scaling phenomena, and $1/f$ noise: a tale of tails.
[J. Stat. Phys.](#), 32:209–230, 1983.

References I

- [1] D. Brockmann, L. Hufnagel, and T. Geisel.
The scaling laws of human travel.
[Nature](#), pages 462–465, 2006. [pdf](#)
- [2] A. Clauset, M. Kogan, and S. Redner.
Safe leads and lead changes in competitive team sports.
[Phys. Rev. E](#), 91:062815, 2015. [pdf](#)
- [3] P. S. Dodds and D. H. Rothman.
Unified view of scaling laws for river networks.
[Physical Review E](#), 59(5):4865–4877, 1999. [pdf](#)
- [4] P. S. Dodds and D. H. Rothman.
Scaling, universality, and geomorphology.
[Annu. Rev. Earth Planet. Sci.](#), 28:571–610, 2000. [pdf](#)

References IV

- [11] D. P. Kiley, A. J. Reagan, L. Mitchell, C. M. Danforth, and P. S. Dodds.
The game story space of professional sports: Australian Rules Football.
[Physical Review E](#), 93, 2016. [pdf](#)
- [12] P. Lévy and M. Loeve.
Processus stochastiques et mouvement brownien.
Gauthier-Villars Paris, 1965.
- [13] D. R. Montgomery and W. E. Dietrich.
Channel initiation and the problem of landscape scale.
[Science](#), 255:826–30, 1992. [pdf](#)

References VI

- [17] A. E. Scheidegger.
The algebra of stream-order numbers.
[United States Geological Survey Professional Paper](#), 525-B:B187–B189, 1967. [pdf](#)
- [18] J. S. Weitz and H. B. Fraser.
Explaining mortality rate plateaus.
[Proc. Natl. Acad. Sci.](#), 98:15383–15386, 2001. [pdf](#)

References II

- [5] W. Feller.
An Introduction to Probability Theory and Its Applications, volume I.
John Wiley & Sons, New York, third edition, 1968.
- [6] T. M. Fink and Y. Mao.
Designing tie knots by random walks.
[Nature](#), 398:31–32, 1999. [pdf](#)
- [7] M. R. Frank, L. Mitchell, P. S. Dodds, and C. M. Danforth.
Happiness and the patterns of life: A study of geolocated Tweets.
[Nature Scientific Reports](#), 3:2625, 2013. [pdf](#)