Biological Contagion

Principles of Complex Systems, Vols. 1 & 2
CSYS/MATH 300 and 303, 2021–2022| @pocsvox

Prof. Peter Sheridan Dodds | @peterdodds

Computational Story Lab | Vermont Complex Systems Center
Vermont Advanced Computing Core | University of Vermont

Outline

Introduction

Simple disease spreading models
- Background
- Prediction
- More models
- Toy metapopulation models
- Model output
- Nutshell
- Other kinds of prediction
- Next

References

🎲 An awful recording: Wikipedia's list of epidemics ⧉ from 430 BC on.

(Screenshot of Wikipedia's "List of epidemics" page)

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next
References

Contagion

A confusion of contagions:

🎲 Did Harry Potter spread like a virus?
🎲 Can disinformation be "infectious"?
🎲 Suicide, violence?
🎲 Morality? Evil? Laziness? Stupidity? Happiness?
🎲 Religion?
🎲 Democracy …?
🎲 Language? The alphabet? [10]
🎲 Stories?

Contagion

Naturomorphisms

🎲 "The feeling was contagious."
🎲 "The news spread like wildfire."
🎲 "Freedom is the most contagious virus known to man."
   —Hubert H. Humphrey, Johnson's vice president
🎲 "Nothing is so contagious as enthusiasm."
   —Samuel Taylor Coleridge

Optimism according to Ambrose Bierce: ⧉

The doctrine that everything is beautiful, including what is ugly, everything good, especially the bad, and everything right that is wrong. … It is hereditary, but fortunately not contagious.

Social contagion

Eric Hoffer, 1902–1983

There is a grandeur in the uniformity of the mass. When a fashion, a dance, a song, a slogan or a joke sweeps like wildfire from one end of the continent to the other, and a hundred million people roar with laughter, sway their bodies in unison, hum one song or break forth in anger and denunciation, there is the overpowering feeling that in this country we have come nearer the brotherhood of man than ever before.

🎲 Hoffer ⧉ was an interesting fellow...

The spread of fanaticism

Hoffer's most famous work: "The True Believer: Thoughts On The Nature Of Mass Movements" (1951) [12]

Aphorisms-aplenty:

🎲 "We can be absolutely certain only about things we do not understand."
🎲 "Mass movements can rise and spread without belief in a God, but never without belief in a devil."
🎲 "Where freedom is real, equality is the passion of the masses. Where equality is real, freedom is the passion of a small minority."


Imitation

"When people are free to do as they please, they usually imitate each other."
—Eric Hoffer "The Passionate State of Mind" [13]

despair.com

The collective...

"Never Underestimate the Power of Stupid People in Large Groups."

despair.com

Examples of non-disease spreading:

Interesting infections:

- Spreading of certain buildings in the US:

Marbleization of the US:

The most terrifying contagious outbreak?

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
  Background
  Prediction
  More models
  Toy metapopulation
  models
  Model output
  Nutshell
  Other kinds of prediction
  Next

References

Contagion

Definitions

- (1) The spreading of a quality or quantity between individuals in a population.
- (2) A disease itself:
  the plague, a blight, the dreaded lurgi, ...
- from Latin: *con* = 'with' + *tangere* 'to touch.'
- Contagion has unpleasant overtones...
- Just Spreading might be a more neutral word
- But contagion is kind of exciting...

Contagions

Two main classes of contagion

1. Infectious diseases:
   tuberculosis, HIV, ebola, SARS, influenza, zombification, ...

2. Social contagion:
   fashion, word usage, rumors, uprisings, religion, stories about zombies, ...


PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
  Background
  Prediction
  More models
  Toy metapopulation
  models
  Model output
  Nutshell
  Other kinds of prediction
  Next

References

Community—S2E6: Epidemiology

Mathematical Epidemiology

The standard SIR model [18]

- = basic model of disease contagion
- Three states:
  1. S = Susceptible
  2. I = Infective/Infectious
  3. R = Recovered or Removed or Refractory
- $S(t) + I(t) + R(t) = 1$
- Presumes random interactions (mass-action principle)
- Interactions are independent (no memory)
- Discrete and continuous time versions

Mathematical Epidemiology

Discrete time automata example:

Transition Probabilities:

β for being infected given contact with infected
r for recovery
ρ for loss of immunity

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
  Background
  Prediction
  More models
  Toy metapopulation
  models
  Model output
  Nutshell
  Other kinds of prediction
  Next

References

Mathematical Epidemiology

Original models attributed to

- 1920's: Reed and Frost
- 1920's/1930's: Kermack and McKendrick [14, 16, 15]
- Coupled differential equations with a mass-action principle

---

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Reproduction Number R_0

Discrete version:

- Set up: One Infective in a randomly mixing population of Susceptibles
- At time $t = 0$, single infective random bumps into a Susceptible
- Probability of transmission = β
- At time $t = 1$, single Infective remains infected with probability $1 - r$
- At time $t = k$, single Infective remains infected with probability $(1-r)^k$

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Independent Interaction models

For the continuous version

- Second equation:

$$\frac{d}{dt}I = \beta SI - rI$$

$$\frac{d}{dt}I = (\beta S - r)I$$

- Number of infectives grows initially if

$$\beta S(0) - r > 0 \Rightarrow \beta S(0) > r \Rightarrow \beta S(0)/r > 1$$

where $S(0) \simeq 1$.

- Same story as for discrete model.

Independent Interaction models

Differential equations for continuous model

$$\frac{d}{dt}S = -\beta IS + \rho R$$

$$\frac{d}{dt}I = \beta IS - rI$$

$$\frac{d}{dt}R = rI - \rho R$$

β, r, and ρ are now rates.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Reproduction Number R_0

Discrete version:

- Expected number infected by original infective:

$$R_0 = \beta + (1-r)\beta + (1-r)^2\beta + (1-r)^3\beta + \ldots$$

$$= \beta\left(1 + (1-r) + (1-r)^2 + (1-r)^3 + \ldots\right)$$

$$= \beta\frac{1}{1-(1-r)} = \beta/r$$

For $S(0) \simeq 1$ initial susceptibles
$(1 - S(0) = R(0)$ = fraction initially immune):

$$R_0 = S(0)\beta/r$$

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Independent Interaction models

Many variants of the SIR model:

- SIS: susceptible-infective-susceptible
- SIRS: susceptible-infective-recovered-susceptible
- compartment models (age or gender partitions)
- more categories such as 'exposed' (SEIRS)
- recruitment (migration, birth)

Reproduction Number R_0

Reproduction Number R_0

- R_0 = expected number of infected individuals resulting from a single initial infective
- Epidemic threshold: If $R_0 > 1$, 'epidemic' occurs.
- Exponential take off: R_0^n where n is the number of generations.
- Fantastically awful notation convention: R_0 and the R in SIR.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Independent Interaction models

Example of epidemic threshold:

Fraction infected (y-axis, 0 to 1) vs R_0 (x-axis, 0 to 4).

- Continuous phase transition.
- Fine idea from a simple model.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Watch someone else pretend to save the world:

MARION COTILLARD   MATT DAMON   LAURENCE FISHBURNE   JUDE LAW   GWYNETH PALTROW   KATE WINSLET

NOTHING SPREADS LIKE FEAR

CONTAGION

Save the world yourself: ⬈

- And you can be the virus. ⬈
- Also contagious?: Cooperative games …

Neural reboot—Save another pretend world with Vax: ⬈

Lesson 4: Quarantine

Vaccines take time to 'kick in' so they're ineffective if an infection has already begun to spread.

Start >

VAX!

Networks Epidemics Vaccines Quarantine

Pandemic severity index (PSI) ⬈

- Classification during/post pandemic:

- Category based.
- 1–5 scale.
- Modeled on the Saffir-Simpson hurricane scale ⬈.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

For novel diseases:

1. Can we predict the size of an epidemic?
2. How important is the reproduction number R_0?

R_0 approximately same for all of the following:

- 1918-19 "Spanish Flu" $\sim$ 75,000,000 world-wide, 500,000 deaths in US.
- 1957-58 "Asian Flu" $\sim$ 2,000,000 world-wide, 70,000 deaths in US.
- 1968-69 "Hong Kong Flu" $\sim$ 1,000,000 world-wide, 34,000 deaths in US.
- 2003 "SARS Epidemic" $\sim$ 800 deaths world-wide.

Size distributions

As we know, heavy-tailed size distributions are somewhat prevalent in complex systems:

- earthquakes (Gutenberg-Richter law)
- city sizes, forest fires, war fatalities
- wealth distributions
- 'popularity' (books, music, websites, ideas)
- Epidemics?

Power law distributions are common but not obligatory…

Really, what about epidemics?

- Simply hasn't attracted much attention.
- Data not as clean as for other phenomena.

Feeling Ill in Iceland ⬈

Caseload recorded monthly for range of diseases in Iceland, 1888-1990

- Treat outbreaks separated in time as 'novel' diseases.

Really not so good at all in Iceland

Epidemic size distributions $N(S)$ for Measles, Rubella, and Whooping Cough.

Spike near $S = 0$, relatively flat otherwise.

Measles & Pertussis

Insert plots:

Complementary cumulative frequency distributions:

$$N(\Psi' > \Psi) \propto \Psi^{-\gamma+1}$$

Limited scaling with a possible break.

Power law distributions

Measured values of γ:

- measles: 1.40 (low Ψ) and 1.13 (high Ψ)
- pertussis: 1.39 (low Ψ) and 1.16 (high Ψ)

- Expect $2 \le \gamma < 3$ (finite mean, infinite variance)
- When $\gamma < 1$, can't normalize
- Distribution is quite flat.

Resurgence—example of SARS

- Epidemic slows...
  then an infective moves to a new context.
- Epidemic discovers new 'pools' of susceptibles:
  Resurgence.
- Importance of rare, stochastic events.

Community—S2E6: Epidemiology

The challenge

So... can a simple model produce

1. broad epidemic distributions
   and
2. resurgence ?

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
  Background
  Prediction
  More models
  Toy metapopulation
  models
  Model output
  Nutshell
 Other kinds of prediction
  Next
References

Size distributions

Simple models typically produce bimodal or unimodal size distributions.

- This includes network models:
  random, small-world, scale-free, ...
- Exceptions:
  1. Forest fire models
 2. Sophisticated metapopulation models

Burning through the population

Forest fire models: [19]

- Rhodes & Anderson, 1996
- The physicist's approach:
  "if it works for magnets, it'll work for people..."

A bit of a stretch:

1. Epidemics ≡ forest fires
   spreading on 3-d and 5-d lattices.
2. Claim Iceland and Faroe Islands exhibit power law distributions for outbreaks.
3. Original forest fire model not completely understood.

Size distributions

From Rhodes and Anderson, 1996.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
  Background
  Prediction
  More models
  Toy metapopulation
  models
  Model output
  Nutshell
  Other kinds of prediction
  Next
References

"The hidden geometry of complex, network-driven contagion phenomena"
Brockmann and Helbing,
Science, **342**, 1337–1342, 2013. [5]

Fig. 1. Complexity in global, network-driven contagion phenomena. (A) The global mobility network (GMN). Gray lines represent passenger flows along direct connections between 4069 airports worldwide. Geographic regions are distinguished by color [classified according to network modularity maximization (39)]. **(B)** Temporal snapshot of a simulated global pandemic with initial outbreak location (OL) in Hong Kong (HKG). The simulation is based on the metapopulation model defined by Eq. 3 with parameters $R_0 = 1.5$, $\beta = 0.285$ day^{-1}, $\gamma = 2.8 \times 10^{-1}$ day^{-1}, $\varepsilon = 10^{-3}$. Red symbols depict locations with epidemic arrival times in the time window 105 days $\leq T_a \leq$ 110 days. Because of the multiscale structure of the underlying network, the spatial distribution of disease prevalence (i.e., the fraction of infected individuals) lacks geometric coherence. No clear wave-front is visible, and based on this dynamic state, the OL cannot be easily deduced. **(C)** For the same simulation as in (B), the panel depicts arrival times T_a as a function of geographic distance D_g from the OL [nodes are colored according to geographic region as in (A)] for each of the 4069 nodes in the network. On a global scale, T_a weakly correlates with geographic distance D_g ($R^2 = 0.34$). A linear fit yields an average global spreading speed of $v_g = 331$ km/day (see also fig. S7). Using D_g and v_g to estimate arrival times for specific locations, however, does not work well owing to the strong variability of the arrival times for a given geographic distance. The red horizontal bar corresponds to the arrival time window shown in (B). **(D)** Arrival times versus geographic distance from the source (Mexico) for the 2009 H1N1 pandemic. Symbols represent 140 affected countries, and symbol size quantifies total traffic per country. Arrival times are defined as the date of the first confirmed case in a given country after the initial outbreak on 17 March 2009. As in the simulated scenario, arrival time and geographic distance are only weakly correlated ($R^2 = 0.0394$). **(E)** In analogy to (D), the panel depicts the arrival times versus geographic distance from the source (China) of the 2003 SARS epidemic for 29 affected countries worldwide. Arrival times are taken from WHO published data (2). As in (C) and (D), arrival time correlates weakly with geographic distance.

Sophisticated metapopulation models:

- Multiscale models suggested earlier by others but not formalized (Bailey [1], Cliff and Haggett [6], Ferguson et al.)
- Community based mixing (two scales)—Longini. [17]
- Eubank et al.'s EpiSims/TRANSIMS—city simulations. [9]
- Spreading through countries—Airlines: Germann et al., Colizza et al. [7]

- GLEAM: Global pandemic simulations by Vespignani et al.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
  Background
  Prediction
  More models
  Toy metapopulation
  models
  Model output
  Nutshell
 Other kinds of prediction
  Next
References

Fig. 2. Understanding global contagion phenomena using effective distance. (A) The structure of the shortest path tree (in gray) from Hong Kong (central node). Radial distance represents effective distance D_{eff} as defined by Eqs. 4 and 5. Nodes are colored according to the same scheme as in Fig. 1A. (B) The sequence (from left to right) of panels depicts the time course of a simulated model disease with initial outbreak in Hong Kong (HKG), for the same parameter set as used in Fig. 1B. Prevalence is reflected by the redness of the symbols. Each panel compares the state of the system in the conventional geographic representation (bottom) with the effective distance representation (top). The complex spatial pattern in the conventional view is equivalent to a homogeneous wave that propagates outwards at constant effective speed in the effective distance representation. (C) Epidemic arrival time T_a versus effective distance D_{eff} for the same simulated epidemic as in (B). In contrast to geographic distance (Fig. 1C), effective distance correlates strongly with arrival time ($R^2 = 0.973$), i.e., effective distance is an excellent predictor of arrival times. (D and E) Linear relationship between effective distance and arrival time for the 2009 H1N1 pandemic (D) and the 2003 SARS epidemic (E). The arrival time data are the same as in Fig. 1, D and E. The effective distance was computed from the projected global mobility network between countries. As in the model system, we observe a strong correlation between arrival time and effective distance.

Fig. 3. Qualitative outbreak reconstruction based on effective distance. (A) Spatial distribution of prevalence $j_n(t)$ at time $T = 81$ days for OL Chicago (parameters $\beta = 0.28$ day^{-1}, $R_0 = 1.9$, $\gamma = 2.8 \times 10^{-1}$ day^{-1}, and $\varepsilon = 10^{-3}$). After this time, it is difficult, if not impossible, to determine the correct OL from snapshots of the dynamics. (B) Candidate OLs chosen from different geographic regions. (C) Panels depict the state of the system shown in (A) from the perspective of each candidate OL, using each OL's shortest path tree representation. Only the actual OL (ORD, circled in blue) produces a circular wavefront. Even for comparable North American airports (Atlanta (ATL), Toronto (YYZ), and Mexico City (MEX)), the wavefronts are not nearly as concentric. Effective distances thus permit the extraction of the correct OL, based on information on the mobility network and a single snapshot of the dynamics.

Community—S2E6: Epidemiology

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

Size distributions

- Vital work but perhaps hard to generalize from...
- ⇒ Create a simple model involving multiscale travel
- Very big question: What is N?
- Should we model SARS in Hong Kong as spreading in a neighborhood, in Hong Kong, Asia, or the world?
- For simple models, we need to know the final size beforehand...

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

Improving simple models

Contexts and Identities—Bipartite networks

- boards of directors
- movies
- transportation modes (subway)

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

Improving simple models

Idea for social networks: incorporate identity

Identity is formed from attributes such as:

- Geographic location
- Type of employment
- Age
- Recreational activities

Groups are crucial...

- formed by people with at least one similar attribute
- Attributes ⇔ Contexts ⇔ Interactions ⇔ Networks. [23]

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

Infer interactions/network from identities

Distance makes sense in identity/context space.

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

Generalized context space

(Blau & Schwartz [3], Simmel [20], Breiger [4])

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

A toy agent-based model:

"Multiscale, resurgent epidemics in a hierarchcial metapopulation model" ⧉
Watts et al.,
Proc. Natl. Acad. Sci., **102**, 11157–11162, 2005. [24]

Geography: allow people to move between contexts

- Locally: standard SIR model with random mixing
- discrete time simulation
- β = infection probability
- γ = recovery probability
- P = probability of travel
- Movement distance: $\Pr(d) \propto \exp(-d/\xi)$
- ξ = typical travel distance

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
More models
Toy metapopulation
models
Model output
Nutshell
Other kinds of prediction
Next

References

A toy agent-based model

Schematic:

$b=2$
$l=3$
$x_{ij}=2$
$n=8$
i
j

Model output

- Define P_0 = Expected number of infected individuals leaving initially infected context.
- Need $P_0 > 1$ for disease to spread (independent of R_0).
- Limit epidemic size by restricting frequency of travel and/or range

---

Model output

Varying ξ:

- Transition in expected final size based on typical movement distance (sensible)

Model output

Varying P_0:

- Transition in expected final size based on typical number of infectives leaving first group (also sensible)
- Travel advisories: ξ has larger effect than P_0.

Example model output: size distributions

- Flat distributions are possible for certain ξ and P.
- Different R_0's may produce similar distributions
- Same epidemic sizes may arise from different R_0's

Model output—resurgence

Standard model:

Model output—resurgence

Standard model with transport:

The upshot

Simple multiscale population structure
+
stochasticity

leads to

resurgence
+
broad epidemic size distributions

Nutshelling

- For the hierarchical movement model, epidemic size is highly unpredictable
- Model is more complicated than SIR but still simple.
- We haven't even included normal social responses such as travel bans and self-quarantine.
- The reproduction number R_0 is not terribly useful.
- R_0, however measured, is not informative about
  1. how likely the observed epidemic size was,
  2. and how likely future epidemics will be.
- Problem: R_0 summarises one epidemic after the fact and enfolds movement, the price of bananas, everything.

Conclusions

- 🦠 Disease's spread is highly sensitive to population structure.
- 🦠 Rare events may matter enormously: e.g., an infected individual taking an international flight.
- 🦠 More support for controlling population movement:
  e.g., travel advisories, quarantine

Nutshelling

What to do:
- 🦠 Need to separate movement from disease
- 🦠 R_0 needs a friend or two.
- 🦠 Need $R_0 > 1$ and $P_0 > 1$ and ξ sufficiently large for disease to have a chance of spreading
- 🦠 And in general: keep building up the kitchen sink models.

More wondering:
- 🦠 Exactly how important are rare events in disease spreading?
- 🦠 Again, what is N?

Krugman, 1998: "Why most economists' predictions are wrong."

"The growth of the Internet will slow drastically, as the flaw in "Metcalfe's law"—which states that the number of potential connections in a network is proportional to the square of the number of participants—becomes apparent: most people have nothing to say to each other! By 2005 or so, it will become clear that the Internet's impact on the economy has been no greater than the fax machine's."[1]

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Economics, Schmeconomics

Alan Greenspan (September 18, 2007):

"I've been dealing with these big mathematical models of forecasting the economy ...

If I could figure out a way to determine whether or not people are more fearful or changing to more euphoric,

I don't need any of this other stuff.

I could forecast the economy better than any way I know."

http://wikipedia.org

Economics, Schmeconomics

Greenspan continues:

"The trouble is that we can't figure that out. I've been in the forecasting business for 50 years. I'm no better than I ever was, and nobody else is. Forecasting 50 years ago was as good or as bad as it is today. And the reason is that human nature hasn't changed. We can't improve ourselves."

Jon Stewart:

"You just bummed the @*!# out of me."

wildbluffmedia.com

- 🦠 From the Daily Show (September 18, 2007)
- 🦠 The full episode is here:
  http://www.cc.com/video-clips/cenrt5/the-daily-show-with-jon-st

Predicting social catastrophe isn't easy...

"Greenspan Concedes Error on Regulation"

- 🦠 ...humbled Mr. Greenspan admitted that he had put too much faith in the self-correcting power of free markets ...
- 🦠 "Those of us who have looked to the self-interest of lending institutions to protect shareholders' equity, myself included, are in a state of shocked disbelief"
- 🦠 Rep. Henry A. Waxman: "Do you feel that your ideology pushed you to make decisions that you wish you had not made?"
- 🦠 Mr. Greenspan conceded: "Yes, I've found a flaw. I don't know how significant or permanent it is. But I've been very distressed by that fact."

New York Times, October 23, 2008

PoCS
@pocsvox
Biological
Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next

References

Economics, Schmeconomics

James K. Galbraith:

NYT But there are at least 15,000 professional economists in this country, and you're saying only two or three of them foresaw the mortgage crisis? [JKG] Ten or 12 would be closer than two or three.

NYT What does that say about the field of economics, which claims to be a science? [JKG] It's an enormous blot on the reputation of the profession. There are thousands of economists. Most of them teach. And most of them teach a theoretical framework that has been shown to be fundamentally useless.

From the New York Times, 11/02/2008

Other attempts to use SIR and co. elsewhere:

- 🦠 Adoption of ideas/beliefs (Goffman & Newell, 1964)[11]
- 🦠 Spread of rumors (Daley & Kendall, 1965)[8]
- 🦠 Diffusion of innovations (Bass, 1969)[2]
- 🦠 Spread of fanatical behavior (Castillo-Chávez & Song, 2003)
- 🦠 Spread of Feynmann diagrams (Bettencourt et al., 2006)

Social contagion:

- 🦠 SIR may apply sometimes ...
- 🦠 But we need new fundamental models.
- 🦠 Next up: Thresholds.

We really should know social contagion is different but ...

"It's contagious: Rethinking a metaphor dialogically"
Warren and Power,
Culture & Psychology, **21**, 359–379, 2015.[22]

- 🦠 "Facebook will lose 80% of users by 2017, say Princeton researchers" (Guardian, 2014)

"Epidemiological modeling of online social network dynamics"
Spechler and Cannarella,
Availabe online at http://arxiv.org/abs/1401.4208, 2014.[21]

The Facebook Data Science team's response :

Mike Develin, Lada Adamic, and Sean Taylor.

References I

[1] N. T. J. Bailey.
The Mathematical Theory of Infectious Diseases and Its Applications.
Griffin, London, Second edition, 1975.

[2] F. Bass.
A new product growth model for consumer durables.
Manage. Sci., 15:215–227, 1969. pdf

[3] P. M. Blau and J. E. Schwartz.
Crosscutting Social Circles.
Academic Press, Orlando, FL, 1984.

[4] R. L. Breiger.
The duality of persons and groups.
Social Forces, 53(2):181–190, 1974. pdf

References II

[5] D. Brockmann and D. Helbing.
The hidden geometry of complex, network-driven contagion phenomena.
Science, 342:1337–1342, 2013. pdf

[6] A. D. Cliff, P. Haggett, J. K. Ord, and G. R. Versey.
Spatial diffusion: an historical geography of epidemics in an island community.
Cambridge University Press, Cambridge, UK, 1981.

[7] V. Colizza, A. Barrat, M. Barthelmey, A.-J. Valleron, and A. Vespignani.
Modeling the worldwide spread of pandemic influenza: Baseline case and containment interventions.
PLoS Med., 4:e13, 2007. pdf

PoCS
@pocsvox
Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next
References

References III

[8] D. J. Daley and D. G. Kendall.
Stochastic rumours.
J. Inst. Math. Appl., 1:42–55, 1965.

[9] S. Eubank, H. Guclu, V. S. A. Kumar, M. V. Marathe, A. Srinivasan, Z. Toroczkai, and N. Wang.
Modelling disease outbreaks in realistic urban social networks.
Nature, 429:180–184, 2004. pdf

[10] J. Gleick.
The Information: A History, A Theory, A Flood.
Pantheon, 2011.

[11] W. Goffman and V. A. Newill.
Generalization of epidemic theory: An application to the transmission of ideas.
Nature, 204:225–228, 1964. pdf

References IV

[12] E. Hoffer.
The True Believer: On The Nature Of Mass Movements.
Harper and Row, New York, 1951.

[13] E. Hoffer.
The Passionate State of Mind: And Other Aphorisms.
Buccaneer Books, 1954.

[14] W. O. Kermack and A. G. McKendrick.
A contribution to the mathematical theory of epidemics.
Proc. R. Soc. Lond. A, 115:700–721, 1927. pdf

References V

[15] W. O. Kermack and A. G. McKendrick.
A contribution to the mathematical theory of epidemics. III. Further studies of the problem of endemicity.
Proc. R. Soc. Lond. A, 141(843):94–122, 1927. pdf

[16] W. O. Kermack and A. G. McKendrick.
Contributions to the mathematical theory of epidemics. II. The problem of endemicity.
Proc. R. Soc. Lond. A, 138(834):55–83, 1927. pdf

[17] I. M. Longini.
A mathematical model for predicting the geographic spread of new infectious agents.
Math. Biosci., 90:367–383, 1988.

PoCS
@pocsvox
Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next
References

References VI

[18] J. D. Murray.
Mathematical Biology.
Springer, New York, Third edition, 2002.

[19] C. J. Rhodes and R. M. Anderson.
Power laws governing epidemics in isolated populations.
Nature, 381:600–602, 1996. pdf

[20] G. Simmel.
The number of members as determining the sociological form of the group. I.
American Journal of Sociology, 8:1–46, 1902.

References VII

[21] J. A. Spechler and J. Cannarella.
Epidemiological modeling of online social network dynamics.
Availabe online at http://arxiv.org/abs/1401.4208, 2014. pdf

[22] Z. J. Warren and S. A. Power.
It's contagious: Rethinking a metaphor dialogically.
Culture & Psychology, 21:359–379, 2015. pdf

[23] D. J. Watts, P. S. Dodds, and M. E. J. Newman.
Identity and search in social networks.
Science, 296:1302–1305, 2002. pdf

References VIII

[24] D. J. Watts, R. Muhamad, D. Medina, and P. S. Dodds.
Multiscale, resurgent epidemics in a hierarchcial metapopulation model.
Proc. Natl. Acad. Sci., 102(32):11157–11162, 2005. pdf

PoCS
@pocsvox
Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
More models
Toy metapopulation models
Model output
Nutshell
Other kinds of prediction
Next
References