

Lognormals and friends

Last updated: 2024/10/14, 16:39:46 EDT

Principles of Complex Systems, Vols. 1, 2, & 3D
CSYS/MATH 6701, 6713, & a pretend number, 2024–2025

Prof. Peter Sheridan Dodds

Computational Story Lab | Vermont Complex Systems Center
Santa Fe Institute | University of Vermont



Licensed under the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/)



These slides are brought to you by:

Sealie & Lambie
Productions



The PoCverse
Lognormals and
friends
2 of 26


Lognormals



Empirical Confusability
Random Multiplicative Growth
Model
Random Growth with Variable
Lifespan

References

These slides are also brought to you by:

Special Guest Executive Producer



 On Instagram at [pratchett_the_cat](https://www.instagram.com/pratchett_the_cat) 

The PoCverse
Lognormals and
friends
3 of 26

Lognormals

Empirical Confusability
Random Multiplicative Growth
Model
Random Growth with Variable
Lifespan

References

Outline

The PoCSverse
Lognormals and
friends
4 of 26

Lognormals

Empirical Confusability
Random Multiplicative Growth
Model
Random Growth with Variable
Lifespan

References

Lognormals


Empirical Confusability

Random Multiplicative Growth Model

Random Growth with Variable Lifespan

References


There are other 'heavy-tailed' distributions:



1. The Log-normal distribution 

$$P(x) = \frac{1}{x\sqrt{2\pi\sigma}} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

2. Weibull distributions 


$$P(x)dx = \frac{k}{\lambda} \left(\frac{x}{\lambda}\right)^{\mu-1} e^{-(x/\lambda)^\mu} dx$$


CCDF = stretched exponential 


3. Also: Gamma distribution , Erlang distribution , and more.

The lognormal distribution:


$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

 $\ln x$ is distributed according to a normal distribution with mean μ and variance σ .

 Appears in economics and biology where growth increments are distributed normally.


 Standard form reveals the mean μ and variance σ^2 of the underlying normal distribution:

$$P(x) = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$$

 For lognormals:

$$\mu_{\text{lognormal}} = e^{\mu + \frac{1}{2}\sigma^2}, \quad \text{median}_{\text{lognormal}} = e^{\mu},$$

$$\sigma_{\text{lognormal}} = (e^{\sigma^2} - 1)e^{2\mu + \sigma^2}, \quad \text{mode}_{\text{lognormal}} = e^{\mu - \sigma^2}.$$

 All moments of lognormals are **finite**.

Derivation from a normal distribution

Take Y as distributed normally:



$$P(y)dy = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right) dy$$

Set $Y = \ln X$:



Transform according to $P(x)dx = P(y)dy$:




$$\frac{dy}{dx} = 1/x \Rightarrow dy = dx/x$$





$$\Rightarrow P(x)dx = \frac{1}{x\sqrt{2\pi}\sigma} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right) dx$$

Confusion between lognormals and pure power laws



Near agreement
over four orders of
magnitude!

 For lognormal (blue), $\mu = 0$ and $\sigma = 10$.

 For power law (red), $\gamma = 1$ and $c = 0.03$.

What's happening:

$$\begin{aligned}\ln P(x) &= \ln \left\{ \frac{1}{x\sqrt{2\pi}\sigma} \exp \left(-\frac{(\ln x - \mu)^2}{2\sigma^2} \right) \right\} \\ &= -\ln x - \ln\sqrt{2\pi}\sigma - \frac{(\ln x - \mu)^2}{2\sigma^2}\end{aligned}$$

$$= -\frac{1}{2\sigma^2} (\ln x)^2 + \left(\frac{\mu}{\sigma^2} - 1 \right) \ln x - \ln\sqrt{2\pi}\sigma - \frac{\mu^2}{2\sigma^2}.$$

If the first term is relatively small,

$$\boxed{\ln P(x) \sim -\left(1 - \frac{\mu}{\sigma^2}\right) \ln x + \text{const.}} \Rightarrow \boxed{\gamma = 1 - \frac{\mu}{\sigma^2}}$$


Lognormals


Empirical Confusability


Random Multiplicative Growth
Model

Random Growth with Variable
Lifespan


References

 If $\mu < 0, \gamma > 1$ which is totally cool.

 If $\mu > 0, \gamma < 1$, not so much.


 If $\sigma^2 \gg 1$ and μ ,

$$\ln P(x) \sim -\ln x + \text{const.}$$

 Expect -1 scaling to hold until $(\ln x)^2$ term becomes significant compared to $(\ln x)$:

$$-\frac{1}{2\sigma^2} (\ln x)^2 \simeq 0.05 \left(\frac{\mu}{\sigma^2} - 1 \right) \ln x$$

$$\Rightarrow \log_{10} x \lesssim 0.05 \times 2(\sigma^2 - \mu) \log_{10} e \simeq 0.05(\sigma^2 - \mu)$$

 \Rightarrow If you find a -1 exponent,
you may have a lognormal distribution...

Generating lognormals:

Random multiplicative growth:



$$x_{n+1} = rx_n$$

where $r > 0$ is a random growth variable



(Shrinkage is allowed)



In log space, growth is by addition:

$$\ln x_{n+1} = \ln r + \ln x_n$$



$\Rightarrow \ln x_n$ is normally distributed




$\Rightarrow x_n$ is lognormally distributed

Lognormals or power laws?

🧱 Gibrat ^[2] (1931) uses preceding argument to explain lognormal distribution of firm sizes ($\gamma \simeq 1$).


🧱 But Robert Axtell ^[1] (2001) shows a power law fits the data very well with $\gamma = 2$, not $\gamma = 1$ (!)


🧱 Problem of data censusing (missing small firms).




$$\text{Freq} \propto (\text{size})^{-\gamma}$$
$$\gamma \simeq 2$$

🧱 One piece in Gibrat's model seems okay empirically: Growth rate r appears to be independent of firm size. ^[1].


 Axtel cites Malcai et al.'s (1999) argument [5] for why power laws appear with exponent $\gamma \simeq 2$


 The set up: N entities with size $x_i(t)$

 Generally:

$$x_i(t + 1) = rx_i(t)$$

where r is drawn from some happy distribution

 Same as for lognormal but one extra piece.

 Each x_i cannot drop too low with respect to the other sizes:

$$x_i(t + 1) = \max(rx_i(t), c \langle x_i \rangle)$$

Some math later...

Insert assignment question 



$$\text{Find } P(x) \sim x^{-\gamma}$$



where γ is implicitly given by

$$N = \frac{(\gamma - 2)}{(\gamma - 1)} \left[\frac{(c/N)^{\gamma-1} - 1}{(c/N)^{\gamma-1} - (c/N)} \right]$$

N = total number of firms.



$$\text{Now, if } c/N \ll 1 \text{ and } \gamma > 2 \quad N = \frac{(\gamma - 2)}{(\gamma - 1)} \left[\frac{-1}{-(c/N)} \right]$$



$$\text{Which gives } \gamma \sim 1 + \frac{1}{1 - c}$$



Groovy... c small $\Rightarrow \gamma \simeq 2$

The second tweak

Ages of firms/people/... may not be the same

☎ Allow the number of updates for each size x_i to vary

☎ Example: $P(t)dt = ae^{-at} dt$ where $t = \text{age}$.

☎ Back to no bottom limit: each x_i follows a lognormal

☎ Sizes are distributed as ^[6]

$$P(x) = \int_{t=0}^{\infty} ae^{-at} \frac{1}{x\sqrt{2\pi t}} \exp\left(-\frac{(\ln x - \mu)^2}{2t}\right) dt$$

(Assume for this example that $\sigma \sim t$ and $\mu = \ln m$)

☎ Now averaging different lognormal distributions.

Averaging lognormals



$$P(x) = \int_{t=0}^{\infty} a e^{-at} \frac{1}{x \sqrt{2\pi t}} \exp\left(-\frac{(\ln \frac{x}{m})^2}{2t}\right) dt$$



Insert fabulous calculation (team is spared).



Some enjoyable suffering leads to:

$$P(x) \propto x^{-1} e^{-\sqrt{2\lambda} (\ln \frac{x}{m})^2}$$

The second tweak



$$P(x) \propto x^{-1} e^{-\sqrt{2\lambda(\ln \frac{x}{m})^2}}$$



Depends on sign of $\ln \frac{x}{m}$, i.e., whether $\frac{x}{m} > 1$ or $\frac{x}{m} < 1$.




$$P(x) \propto \begin{cases} x^{-1+\sqrt{2\lambda}} & \text{if } \frac{x}{m} < 1 \\ x^{-1-\sqrt{2\lambda}} & \text{if } \frac{x}{m} > 1 \end{cases}$$



'Break' in scaling (not uncommon)



Double-Pareto distribution 








First noticed by Montroll and Shlesinger [7, 8]



Later: Huberman and Adamic [3, 4]: Number of pages per website




Summary of these exciting developments:

-  Lognormals and power laws can be **awfully** similar
-  Random Multiplicative Growth leads to lognormal distributions
-  Enforcing a minimum size leads to a power law tail
-  With no minimum size but a distribution of lifetimes, the double Pareto distribution appears
-  Take-home message: Be careful out there...

References I

- [1] R. Axtell.
Zipf distribution of U.S. firm sizes.
Science, 293(5536):1818–1820, 2001. [pdf](#) 
- [2] R. Gibrat.
Les inégalités économiques.
Librairie du Recueil Sirey, Paris, France, 1931.
- [3] B. A. Huberman and L. A. Adamic.
Evolutionary dynamics of the World Wide Web.
Technical report, Xerox Palo Alto Research Center, 1999.
- [4] B. A. Huberman and L. A. Adamic.
The nature of markets in the World Wide Web.
Quarterly Journal of Economic Commerce, 1:5–12, 2000.

References II

- [5] O. Malcai, O. Biham, and S. Solomon.
Power-law distributions and lévy-stable intermittent fluctuations in stochastic systems of many autocatalytic elements.
[Phys. Rev. E](#), 60(2):1299–1303, 1999. pdf 
- [6] M. Mitzenmacher.
A brief history of generative models for power law and lognormal distributions.
[Internet Mathematics](#), 1:226–251, 2003. pdf 
- [7] E. W. Montroll and M. W. Shlesinger.
On $1/f$ noise and other distributions with long tails.
[Proc. Natl. Acad. Sci.](#), 79:3380–3383, 1982. pdf 

- [8] E. W. Montroll and M. W. Shlesinger.
Maximum entropy formalism, fractals, scaling phenomena,
and $1/f$ noise: a tale of tails.
[J. Stat. Phys.](#), 32:209–230, 1983.