

Optimal Supply Networks II: Blood, Water, and Truthicide

Last updated: 2026/03/11, 16:11:41 EDT

DOI:

Principles of Complex Systems,
Vols. 1, 2, 3D, 4 Fourever, V for Vendetta

Prof. Peter Sheridan Dodds

Computational Story Lab | Vermont Complex Systems Institute
University of Vermont | Santa Fe Institute



Licensed under the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/)

The PoCVerse
Optimal Supply
Networks II
1 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



These slides are brought to you by:

Sealie & Lambie Productions

The PoCverse
Optimal Supply
Networks II
2 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument







Conclusion

References



These slides are also brought to you by:

Special Guest Executive Producer



The PoCverse
Optimal Supply
Networks II
3 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

 On Instagram at [pratchett_the_cat](https://www.instagram.com/pratchett_the_cat) 



Outline

Metabolism and Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

The PoCSverse
Optimal Supply
Networks II
4 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Stories—The Fraction Assassin:

The PoCverse
Optimal Supply
Networks II
5 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Law and Order, Special Science Edition: Truthicide Department

“In the scientific integrity system known as peer review, the people are represented by two highly overlapping yet equally important groups: the independent scientists who review papers and the scientists who punish those who publish garbage. This is one of their stories.”



Animal power

The PoCVerse
Optimal Supply
Networks II
7 of 125

Metabolism and
Truthicide

Fundamental biological and ecological constraint:

$$P = c M^\alpha$$

P = basal metabolic rate

M = organismal body mass



Does 1 elephant equal 1 million shrews in a elephant suit in a trenchcoat?

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



$$P = c M^\alpha$$

Prefactor c depends on **body plan** and **body temperature**:

Birds	39–41 °C
Eutherian Mammals	36–38 °C
Marsupials	34–36 °C
Monotremes	30–31 °C



What one might expect:

$\alpha = 2/3$ because ...

 Dimensional analysis suggests an energy balance surface law:

$$P \propto S \propto V^{2/3} \propto M^{2/3}$$

 Assumes isometric scaling (not quite the spherical cow).

 **Lognormal fluctuations:**

Gaussian fluctuations in $\log_{10} P$ around $\log_{10} cM^\alpha$.

 Stefan-Boltzmann law  for radiated energy:

$$\frac{dE}{dt} = \sigma \epsilon S T^4 \propto S$$



The prevailing belief of the Church of Quarterology:

The PoC'sverse
Optimal Supply
Networks II
10 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

$$\alpha = 3/4$$

$$P \propto M^{3/4}$$

Huh?



The prevailing belief of the Church of Quarterology:

The PoCVerse
Optimal Supply
Networks II
11 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

Most obvious concern:

$$3/4 - 2/3 = 1/12$$

-  An exponent higher than $2/3$ points suggests a fundamental inefficiency in biology.
-  Organisms must somehow be running 'hotter' than they need to balance heat loss.



Related putative scalings:

Wait! There's more!:

-  number of capillaries $\propto M^{3/4}$
-  time to reproductive maturity $\propto M^{1/4}$
-  heart rate $\propto M^{-1/4}$
-  cross-sectional area of aorta $\propto M^{3/4}$
-  population density $\propto M^{-3/4}$



The great 'law' of heartbeats:

Assuming:

-  Average lifespan $\propto M^{\beta}$
-  Average heart rate $\propto M^{-\beta}$
-  Irrelevant but perhaps $\beta = 1/4$.

Then:

$$\begin{aligned} \text{Average number of heart beats in a lifespan} \\ \simeq (\text{Average lifespan}) \times (\text{Average heart rate}) \\ \propto M^{\beta-\beta} \\ \propto M^0 \end{aligned}$$

-  Number of heartbeats per life time is independent of organism size!
-  ≈ 1.5 billion



From earlier in PoCS:



“How fast do living organisms move: Maximum speeds from bacteria to elephants and whales” [↗](#)
 Meyer-Vernet and Rospars,
 American Journal of Physics, **83**, 719–722, 2015. [36]




Fig. 1. Maximum relative speed versus body mass for 202 running species (157 mammals plotted in magenta and 45 non-mammals plotted in green), 127 swimming species and 91 micro-organisms (plotted in blue). The sources of the data are given in Ref. 16. The solid line is the maximum relative speed [Eq. (13)] estimated in Sec. III. The human world records are plotted as asterisks (upper for running and lower for swimming). Some examples of organisms of various masses are sketched in black (drawings by François Meyer).



Abstract

Speed is a key trait for many animals, but the relationship between speed and body size is not well understood. We use a general scaling law to reveal why the largest animals are not the fastest. We find that the relationship between speed and body size is not linear, but follows a power law. This means that the largest animals are not the fastest, but the relationship between speed and body size is not as simple as it seems.

"A general scaling law reveals why the largest animals are not the fastest"

Hirt et al.,

Nature Ecology & Evolution, **1**, 1116, 2017. [24]

The PoCVerse
Optimal Supply
Networks II
15 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References




Figure 2 | Empirical data and time-dependent model fit for the allometric scaling of maximum speed. **a.** Comparison of scaling for the different locomotion modes (flying, running, swimming). **b–d.** Taxonomic differences are illustrated separately for flying (**b**; $n=55$), running (**c**; $n=458$) and swimming (**d**; $n=109$) animals. Overall model fit: $R^2=0.893$. The residual variation does not exhibit a signature of taxonomy (only a weak effect of thermoregulation; see Methods).





“A general scaling law reveals why the largest animals are not the fastest” 

Hirt et al.,

Nature Ecology & Evolution, **1**, 1116, 2017. ^[24]

The PoC Sverse
Optimal Supply
Networks II
16 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References




Figure 1 | Concept of time-dependent and mass-dependent realized maximum speed of animals. **a**, Acceleration of animals follows a saturation curve (solid lines) approaching the theoretical maximum speed (dotted lines) depending on body mass (colour code). **b**, The time available for acceleration increases with body mass following a power law. **c**, **d**, This critical time determines the realized maximum speed (**c**), yielding a hump-shaped increase of maximum speed with body mass (**d**).



Theoretical story:




Figure 4 | Predicting the maximum speed of extinct species with the time-dependent model. The model prediction (grey line) is fitted to data of extant species (grey circles) and extended to higher body masses. Speed data for dinosaurs (green triangles) come from detailed morphological model calculations (values in Table 1) and were not used to obtain model parameters.



Maximum speed increases with size: $v_{\max} = aM^b$



Takes a while to get going: $v(t) = v_{\max}(1 - e^{-kt})$



$k \sim F_{\max}/M \sim cM^{d-1}$
Literature: $0.75 \lesssim d \lesssim 0.94$



Acceleration time = depletion time for anaerobic energy: $\tau \sim fM^g$
Literature: $0.76 \lesssim g \lesssim 1.27$



$v_{\max} = aM^b (1 - e^{-hM^i})$



$i = d - 1 + g$ and $h = cf$



Literature search for for maximum speeds of running, flying and swimming animals.



Search terms: “maximum speed”, “escape speed” and “sprint speed”.

Note: [36] not cited.



A theory is born:

The PoCverse
Optimal Supply
Networks II
19 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

1840's: Sarrus and Rameaux^[45] first suggested $\alpha = 2/3$.



A theory grows:

1883: Rubner^[43] found $\alpha \simeq 2/3$.



The PoCverse
Optimal Supply
Networks II
20 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Theory meets a different ‘truth’:

1930's: Brody, Benedict study mammals. [6]
Found $\alpha \simeq 0.73$ (standard).



The PoCverse
Optimal Supply
Networks II
21 of 125
Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Our hero faces a shadowy cabal:

The PoCverse
Optimal Supply
Networks II
22 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



- 1932: Kleiber analyzed 13 mammals. ^[26]
- Found $\alpha = 0.76$ and suggested $\alpha = 3/4$.
- Scaling law of Metabolism became known as Kleiber's Law  (2011 Wikipedia entry is embarrassing).
- 1961 book: "The Fire of Life. An Introduction to Animal Energetics". ^[27]



When a cult becomes a religion:

1950/1960: Hemmingsen [21, 22]

Extension to unicellular organisms.

$\alpha = 3/4$ assumed true.



The PoCverse
Optimal Supply
Networks II
23 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Quarterology spreads throughout the land:

The Cabal assassinates 2/3-scaling:

- 1964: Troon, Scotland.
 - 3rd Symposium on Energy Metabolism.
 - $\alpha = 3/4$ made official ...
- ... 29 to zip.



- But the Cabal slipped up by publishing the conference proceedings ...
- “Energy Metabolism; Proceedings of the 3rd symposium held at Troon, Scotland, May 1964,” Ed. Sir Kenneth Blaxter ^[4]

The PoCverse
Optimal Supply
Networks II
24 of 125
Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



An unsolved truthicide:

The PoCverse
Optimal Supply
Networks II
26 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

So many questions ...

Did the truth kill a theory? Or did a theory kill the truth?

Or was the truth killed by just a lone, lowly hypothesis?

Does this go all the way to the top?
To the National Academies of Science?

Is $2/3$ -scaling really dead?

Could $2/3$ -scaling have faked its own death?

What kind of people would vote on scientific facts?



Modern Quarterology, Post Truthicide

The PoCSeve
Optimal Supply
Networks II
27 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories


Geometric argument

Conclusion

References



$3/4$ is held by many to be the one true exponent.



*In the Beat of a Heart: Life, Energy, and the
Unity of Nature*—by John Whitfield



But: much controversy ...



See ‘Re-examination of the “ $3/4$ -law” of metabolism’
by the Heretical Unbelievers Dodds, Rothman, and
Weitz^[14], and ensuing madness ...



Some data on metabolic rates

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



Heusner's data
(1991)^[23]



391 Mammals



blue line: 2/3




red line: 3/4.



($B = P$)



Some data on metabolic rates



 Passerine vs. non-passerine issue ...

 Bennett and
Harvey's data
(1987)^[3]

 398 birds

 blue line: $2/3$

 red line: $3/4$.

 ($B = P$)



Linear regression

The PoCVerse
Optimal Supply
Networks II
30 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

Important:

🧱 Ordinary Least Squares (OLS) Linear regression is only appropriate for analyzing a dataset $\{(x_i, y_i)\}$ when we know the x_i are measured without error.

🧱 Here we assume that measurements of mass M have less error than measurements of metabolic rate B .

🧱 Linear regression assumes Gaussian errors.



Measuring exponents

The PoCVerse
Optimal Supply
Networks II
31 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

More on regression:

If (a) we don't know what the errors of either variable are,

or (b) no variable can be considered independent,

then we need to use

Standardized Major Axis Linear Regression. ^[44, 42]

(aka Reduced Major Axis = RMA.)



Measuring exponents

The PoCVerse
Optimal Supply
Networks II
32 of 125
Metabolism and
Truthicide

For Standardized Major Axis Linear Regression:

$$\text{slope}_{\text{SMA}} = \frac{\text{standard deviation of } y \text{ data}}{\text{standard deviation of } x \text{ data}}$$

-  Very simple!
-  Minimization of sum of areas of triangles induced by vertical and horizontal residuals with best fit line.
-  The only linear regression that is Scale invariant .
-  Attributed to Nobel Laureate economist Paul Samuelson  ^[44] but discovered independently by others.
-  #somuchwin

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Measuring exponents

Relationship to ordinary least squares regression is simple:

$$\begin{aligned}\text{slope}_{\text{SMA}} &= r^{-1} \times \text{slope}_{\text{OLS } y \text{ on } x} \\ &= r \times \text{slope}_{\text{OLS } x \text{ on } y}\end{aligned}$$

where r = standard correlation coefficient:

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}$$



Groovy upshot: If (1) a paper uses OLS regression when RMA would be appropriate, and (2) r is reported, we can figure out the RMA slope. ^[42, 30]

The PoCVerse
Optimal Supply
Networks II
33 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References






FIG. 4. Observed correlation of calculated windspeed and airspeed in gliding Black-browed albatrosses showing regression and r.m.a. lines. Figure altered from Pennycuik (1982), figure 9.

LINEAR RELATIONS IN BIOMECHANICS

TABLE II

Calculated statistics of airspeed V_a and windspeed V_w in the Black-browed albatross *Diomedea melanophris* in gliding flight, after Pennycuik (1982)

number of data n	737		
means $\bar{x}, \bar{y}$	-3.14	13.35	ms^{-1}
variances S_{xx}, S_{yy}	13.91	8.218	$(\text{ms}^{-1})^2$
covariance S_{xy}	-4.653		
correlation ρ	-0.435		

model of speed correction: $V_a = x + \beta V_w$

model	intercept α	gradient β	range (95%)
$y(x)$ regression	12.30	-0.334	-0.384 to -0.284
r.m.a.	10.93	-0.769	-0.894 to -0.661
$x(y)$ regression	7.80	-1.766	-2.076 to -1.536
s.r. $b_x = 0.5$	10.66	-0.855	-0.997 to -0.737
$b_x = 1$ or m.a.	11.59	-0.560	-0.648 to -0.479
$b_x = 2$	12.00	-0.431	-0.496 to -0.367

Disparity between slopes for y on x and x on y regressions is a factor of r^2 (r^{-2})

(Rayner uses ρ for r .)

Here: $r^2 = .435^2 = 0.189$, and
 $r^{-2} = .435^{-2} = 2.29^2 = 5.285$.

See also: LaBarbera^[30] (who resigned ...)



Heusner's data, 1991 (391 Mammals)

range of M	N	$\hat{\alpha}$
≤ 0.1 kg	167	0.678 ± 0.038
≤ 1 kg	276	0.662 ± 0.032
≤ 10 kg	357	0.668 ± 0.019
≤ 25 kg	366	0.669 ± 0.018
≤ 35 kg	371	0.675 ± 0.018
≤ 350 kg	389	0.706 ± 0.016
≤ 3670 kg	391	0.710 ± 0.021

The PoCserve
Optimal Supply
Networks II
35 of 125
Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Bennett and Harvey, 1987 (398 birds)

$M_{\max}$	N	$\hat{\alpha}$
≤ 0.032	162	0.636 ± 0.103
≤ 0.1	236	0.602 ± 0.060
≤ 0.32	290	0.607 ± 0.039
≤ 1	334	0.652 ± 0.030
≤ 3.2	371	0.655 ± 0.023
≤ 10	391	0.664 ± 0.020
≤ 32	396	0.665 ± 0.019
≤ 100	398	0.664 ± 0.019

The PoCverse
Optimal Supply
Networks II
36 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Fluctuations—Things look normal ...

The PoCVerse
Optimal Supply
Networks II
37 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



 $P(B|M) = 1/M^{2/3} f(B/M^{2/3})$

 Use a Kolmogorov-Smirnov test.



Hypothesis testing

The PoCVerse
Optimal Supply
Networks II
38 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

Test to see if α' is consistent with our data $\{(M_i, B_i)\}$:

$$H_0 : \alpha = \alpha' \text{ and } H_1 : \alpha \neq \alpha'.$$

-  Assume each B_i (now a random variable) is normally distributed about $\alpha' \log_{10} M_i + \log_{10} c$.
-  Follows that the measured α for one realization obeys a t distribution with $N - 2$ degrees of freedom.
-  Calculate a p -value: probability that the measured α is as least as different to our hypothesized α' as we observe.
-  See, for example, DeGroot and Scherish, “Probability and Statistics.” ^[11]



Revisiting the past—mammals

The PoCverse
Optimal Supply
Networks II
39 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

Full mass range:

	N	$\hat{\alpha}$	$p_{2/3}$	$p_{3/4}$
Kleiber	13	0.738	$< 10^{-6}$	0.11
Brody	35	0.718	$< 10^{-4}$	$< 10^{-2}$
Heusner	391	0.710	$< 10^{-6}$	$< 10^{-5}$
Bennett and Harvey	398	0.664	0.69	$< 10^{-15}$



Revisiting the past—mammals

$M \leq 10$ kg:

	N	$\hat{\alpha}$	$p_{2/3}$	$p_{3/4}$
Kleiber	5	0.667	0.99	0.088
Brody	26	0.709	$< 10^{-3}$	$< 10^{-3}$
Heusner	357	0.668	0.91	$< 10^{-15}$

$M \geq 10$ kg:

	N	$\hat{\alpha}$	$p_{2/3}$	$p_{3/4}$
Kleiber	8	0.754	$< 10^{-4}$	0.66
Brody	9	0.760	$< 10^{-3}$	0.56
Heusner	34	0.877	$< 10^{-12}$	$< 10^{-7}$

The PoCverse
Optimal Supply
Networks II
40 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Analysis of residuals

The PoCVerse
Optimal Supply
Networks II
41 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

1. Presume an exponent of your choice: $2/3$ or $3/4$.
2. Fit the prefactor ($\log_{10} c$) and then examine the residuals:

$$r_i = \log_{10} B_i - (\alpha' \log_{10} M_i - \log_{10} c).$$

3. H_0 : residuals are uncorrelated
 H_1 : residuals are correlated.
4. Measure the correlations in the residuals and compute a p -value.



Analysis of residuals

The PoCVerse
Optimal Supply
Networks II
42 of 125
Metabolism and
Truthicide

We use the spiffing Spearman Rank-Order Correlation Coefficient ↗

Basic idea:

🧱 Given $\{(x_i, y_i)\}$, rank the $\{x_i\}$ and $\{y_i\}$ separately from smallest to largest. Call these ranks R_i and S_i .

🧱 Now calculate correlation coefficient for ranks, r_s :



$$r_s = \frac{\sum_{i=1}^n (R_i - \bar{R})(S_i - \bar{S})}{\sqrt{\sum_{i=1}^n (R_i - \bar{R})^2} \sqrt{\sum_{i=1}^n (S_i - \bar{S})^2}}$$

🧱 Perfect correlation: x_i 's and y_i 's both increase monotonically.

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Analysis of residuals

The PoCverse
Optimal Supply
Networks II
43 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

We assume all rank orderings are equally likely:

-  r_s is distributed according to a Student's t -distribution  with $N - 2$ degrees of freedom.
-  Excellent feature: Non-parametric—real distribution of x 's and y 's doesn't matter.
-  Bonus: works for non-linear monotonic relationships as well.
-  See Numerical Recipes in C/Fortran  which contains many good things. ^[40]



Analysis of residuals—mammals

The PoCserve
Optimal Supply
Networks II
44 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



Analysis of residuals—birds

The PoCserve
Optimal Supply
Networks II
45 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



- (a) $M < 0.1$ kg,
- (b) $M < 1$ kg,
- (c) $M < 10$ kg,
- (d) all birds.



Other approaches to measuring exponents:

 Clauset, Shalizi, Newman: “Power-law distributions in empirical data” ^[10]
SIAM Review, 2009.

 See Clauset’s page on [measuring power law exponents](#) 
(code, other goodies).

 See [this collection of tweets](#)  for related amusement.



Impure scaling?:

The PoCverse
Optimal Supply
Networks II
47 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

So: The exponent $\alpha = 2/3$ works for all birds and mammals up to 10–30 kg

For mammals $> 10\text{--}30$ kg, maybe we have a new scaling regime

Possible connection?: Economos (1983)—limb length break in scaling around 20 kg^[15]

But see later: non-isometric growth leads to **lower** metabolic scaling. Oops.



The widening gyre:

Now we're really confused (empirically):

- White and Seymour, 2005: unhappy with large herbivore measurements^[57]. Pro 2/3: Find $\alpha \simeq 0.686 \pm 0.014$.
- Glazier, BioScience (2006)^[19]: “The 3/4-Power Law Is Not Universal: Evolution of Isometric, Ontogenetic Metabolic Scaling in Pelagic Animals.”
- Glazier, Biol. Rev. (2005)^[18]: “Beyond the 3/4-power law’: variation in the intra- and interspecific scaling of metabolic rate in animals.”
- Savage et al., PLoS Biology (2008)^[46] “Sizing up allometric scaling theory” Pro 3/4: problems claimed to be finite-size scaling.



Somehow, optimal river networks are connected:

The PoCverse
Optimal Supply
Networks II
49 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



a = drainage basin
area



l = length of longest
(main) stream



$L = L_{\parallel}$ =
longitudinal length
of basin



Mysterious allometric scaling in river networks

The PoCVerse
Optimal Supply
Networks II
50 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



1957: J. T. Hack^[20]

“Studies of Longitudinal Stream Profiles in Virginia and Maryland”

$$\ell \sim a^h$$

$$h \sim 0.6$$



Anomalous scaling: we would expect $h = 1/2 \dots$



Subsequent studies: $0.5 \lesssim h \lesssim 0.6$



Another quest to find **universality/god** ...




A catch: studies done on small scales.



Large-scale networks:

(1992) Montgomery and Dietrich ^[37]:



 **Composite data set:** includes everything from unchanneled valleys up to world's largest rivers.

 **Estimated fit:**

$$L \simeq 1.78a^{0.49}$$

 **Mixture of basin and main stream lengths.**

The PoCVerse
Optimal Supply
Networks II
51 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



World's largest rivers only:

The PoCServe
Optimal Supply
Networks II
52 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents


River networks

Earlier theories

Geometric argument

Conclusion

References



Data from Leopold (1994) ^[32, 13]



Estimate of Hack exponent: $h = 0.50 \pm 0.06$



Earlier theories (1973–):

Building on the surface area idea:

- 🧱 McMahon (70's, 80's): Elastic Similarity ^[33, 35]
- 🧱 Idea is that organismal shapes scale allometrically with $1/4$ powers (like trees ...)
- 🧱 Disastrously, cites Hemmingsen ^[22] for surface area data.
- 🧱 Appears to be true for ungulate legs ... ^[34]
- 🧱 Metabolism and shape never properly connected.



"Size and shape in biology" ↗

T. McMahon,

Science, **179**, 1201–1204, 1973. [33]




Fig. 3. (a) Chest circumference, d_c , plotted against body weight, W , for five species of primates. The broken lines represent the standard error in this least-squares fit [adapted from (21)]. The model proposed here, whereby each length, l , increases as the $3/4$ power of diameter, d , is illustrated for two weights differing by a factor of 16. (b) Body surface area plotted against weight for vertebrates. The animal data are reasonably well fitted by the stretched cylinder model [adapted from (8)].



Sharks: “We already told you. No.”




“The geometry of life: Testing the scaling of whole-organism surface area and volume using sharks”

Gayford et al.,

Royal Society Open Science, **12**, 242205, 2025. ^[17]

The Geometric Scaling of Sharks

The biggest study of geometric scaling in large animals found that life follows math's rules. As a 3D object grows while maintaining its shape, its surface area scales as the two-thirds power of volume. This is also true for a diverse set of shark species.



Quanta Magazine article: “Shark Data Suggests Animals Scale Like Geometric Objects”

The PoCVerse
Optimal Supply
Networks II
55 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks


Earlier theories

Geometric argument

Conclusion

References





assuming a specific gravity of 1.0. Naturally, the inclination of this line corresponds to a proportionality power of 0.67.

Of the unicellular organisms represented in fig. 1 not a few are spherical in shape (the bacterium *Sarcella*, *Saccharomyces*, marine eggs); and most of the others have surfaces exceeding those of spheres of equal volume by rarely more than what corresponds to 0.1 decade in the log-ordinate system (*Planobacterium phosphoreum*: 12 %, i.e. 0.05 decade, *Escherichia coli*: 34 %, i.e. 0.13 decade, the ciliates *Golfidium* and *Paramecium*: 10–22 %, i.e. about 0.08–0.09 decade; calculated on the basis of data of PÜRMER, 1924, table 7 on p. 190, and HAVERTY, 1928, table 11). Similar figures probably hold for other ciliates. Only the flagellates represented (*Trypanosoma*, *Astasia* *Neobell*) and certain amoebae are likely to deviate by higher figures. The surface values of the unicellular organisms represented in fig. 1 will, therefore, fall either on, or in most other cases less than 0.1 decade above, a line representing the relation between surface and volume of spheres.

It will be seen from fig. 10 that the points representing the body surfaces of the metazoan animals in question are grouped parallel to the sphere line; that is, also corresponding to a proportionality power of 0.67. An average line through the points would fall about 0.30 logarithmic decade above the sphere line, meaning that on the average the body surface is roughly 2 (anti-log. 0.30) times higher in the animals under study than in spheres of equal weight or volume. In organisms of extreme shapes as the python (10⁴ g) and the beech trees (especially marked in fig. 3) the surface is about 3 and 10 times, respectively, greater than in a sphere of equal weight and volume. These facts agree well with the values 8–11.8 for the constant *k* in the formula

$$\text{body surface in cm}^2 = k \cdot \text{body weight}^{0.67}$$

as tabularized by BISCOE (1928, p. 375) for various birds and mammals weighing 8 g–14 kg; because this is about double the value of *k* for sphere surface (4.85). The value of *k* (13.95) found by RAUHA (1940) for *Acacia* is 2.9 times 4.85, and this corresponds well with the above mentioned figure 3 for the much larger python of similar shape.



Hemmingsen's "fit" is for a 2/3 power, notes possible 10 kg transition. [?]



p 46: "The energy metabolism thus definitely varies interspecifically over similar wide weight ranges with a higher power of the body weight than the body surface."



Earlier theories (1977):

The PoCVerse
Optimal Supply
Networks II
57 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

Building on the surface area idea ...

 Blum (1977) ^[5] speculates on four-dimensional biology:

$$P \propto M^{(d-1)/d}$$

 $d = 3$ gives $\alpha = 2/3$

 $d = 4$ gives $\alpha = 3/4$

 So we need another dimension ...





 Obviously, a bit silly... ^[47]



Nutrient delivering networks:

1960's: Rashevsky considers blood networks and finds a $2/3$ scaling.

1997: West *et al.* [54] use a network story to find $3/4$ scaling.



Nutrient delivering networks:

The PoCverse
Optimal Supply
Networks II
59 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

West et al.'s assumptions:

1. hierarchical network
2. capillaries (delivery units) invariant
3. network impedance is minimized via evolution

Claims:

-  $P \propto M^{3/4}$
-  networks are fractal
-  quarter powers everywhere



Impedance measures:



Poiseuille flow (outer branches):

$$Z = \frac{8\mu}{\pi} \sum_{k=0}^N \frac{\ell_k}{r_k^4 N_k}$$



Pulsatile flow (main branches):

$$Z \propto \sum_{k=0}^N \frac{h_k^{1/2}}{r_k^{5/2} N_k}$$



Wheel out Lagrange multipliers ...



Poiseuille gives $P \propto M^1$ with a logarithmic correction.



Pulsatile calculation explodes into flames.



Not so fast ...

Actually, model shows:

 $P \propto M^{3/4}$ does not follow for pulsatile flow

 networks are not necessarily fractal.

Do find:

 Murray's cube law (1927) for outer branches: [38]

$$r_0^3 = r_1^3 + r_2^3$$

 Impedance is distributed evenly.

 Can still assume networks are fractal.



Connecting network structure to α

1. Ratios of network parameters:

$$R_n = \frac{n_{k+1}}{n_k}, R_\ell = \frac{\ell_{k+1}}{\ell_k}, R_r = \frac{r_{k+1}}{r_k}$$

Note: $R_\ell, R_r < 1$, inverse of stream ordering definition.

2. Number of capillaries $\propto P \propto M^\alpha$.

$$\Rightarrow \boxed{\alpha = -\frac{\ln R_n}{\ln R_r^2 R_\ell}}$$

(also problematic due to prefactor issues)

Obliviously soldiering on, we could assert:

🌀 area-preservingness: $R_r = R_n^{-1/2}$

🌀 space-fillingness: $R_\ell = R_n^{-1/3}$

$$\Rightarrow \alpha = 3/4$$



Data from real networks:

Network	R_n	R_r	R_ℓ	$-\frac{\ln R_r}{\ln R_n}$	$-\frac{\ln R_\ell}{\ln R_n}$	α
West <i>et al.</i>	–	–	–	1/2	1/3	3/4
rat (PAT)	2.76	1.58	1.60	0.45	0.46	0.73
cat (PAT) (Turcotte <i>et al.</i> [51])	3.67	1.71	1.78	0.41	0.44	0.79
dog (PAT)	3.69	1.67	1.52	0.39	0.32	0.90
pig (LCX)	3.57	1.89	2.20	0.50	0.62	0.62
pig (RCA)	3.50	1.81	2.12	0.47	0.60	0.65
pig (LAD)	3.51	1.84	2.02	0.49	0.56	0.65
human (PAT)	3.03	1.60	1.49	0.42	0.36	0.83
human (PAT)	3.36	1.56	1.49	0.37	0.33	0.94



Attempts to look at actual networks:



“Testing foundations of biological scaling theory
using automated measurements of vascular
networks” 

Newberry, Newberry, and Newberry,
PLoS Comput Biol, **11**, e1004455, 2015. ^[39]




“” 

Newberry et al.,
PLoS Comput Biol, **11**, e1004455, . ^[?]



Some people understand it's truly a disaster:



“Power, Sex, Suicide: Mitochondria and the Meaning
of Life” 
by Nick Lane (2005). ^[31]

“As so often happens in science, the apparently solid foundations of
a field turned to rubble on closer inspection.”



Let's never talk about this again:



“The fourth dimension of life: Fractal geometry and allometric scaling of organisms” 

West, Brown, and Enquist,
Science, **284**, 1677–1679, 1999. ^[55]

-  No networks: Scaling argument for energy exchange area a .
-  Distinguish between biological and physical length scales (distance between mitochondria versus cell radius).
-  Buckingham π action. ^[9]
-  Arrive at $a \propto M^{D/D+1}$ and $\ell \propto M^{1/D}$.
-  New disaster: after going on about fractality of a , then state $v \propto a\ell$ in general.



“It was the epoch of belief, it was the epoch of incredulity”



“A General Model for the Origin of Allometric Scaling Laws in Biology” ↗

West, Brown, and Enquist,
Science, **276**, 122–126, 1997. [54]



“Nature” ↗

West, Brown, and Enquist,
Nature, **400**, 664–667, 1999. [56]



“The fourth dimension of life: Fractal geometry and allometric scaling of organisms” ↗

West, Brown, and Enquist,
Science, **284**, 1677–1679, 1999. [55]

The PoCverse
Optimal Supply
Networks II
67 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Really, quite confused:

Whole 2004 issue of Functional Ecology addresses the problem:

-  J. Kozlowski, M. Konrzewski. “Is West, Brown and Enquist’s model of allometric scaling mathematically correct and biologically relevant?” Functional Ecology 18: 283–9, 2004. ^[29]
-  J. H. Brown, G. B. West, and B. J. Enquist. “Yes, West, Brown and Enquist’s model of allometric scaling is both mathematically correct and biologically relevant.” Functional Ecology 19: 735–738, 2005. ^[7]
-  J. Kozlowski, M. Konrzewski. “West, Brown and Enquist’s model of allometric scaling again: the same questions remain.” Functional Ecology 19: 739–743, 2005.





"Curvature in metabolic scaling"

Kolokotronis, Savage, Deeds, Fontana, and Rummer.

Nature, ~~464~~, 753, 2010. [28]

Let's try a quadratic:

$$\log_{10} P \sim \log_{10} c + \alpha_1 \log_{10} M + \alpha_2 \log_{10} M^2$$



Yah:




Figure 1 | Curvature in metabolic scaling. **a**, Linear (red) and quadratic (blue) fits (not including temperature) of $\log_{10}B$ versus $\log_{10}M$. The orca (green square) and Asian elephant (ref. 4; turquoise square at larger mass) are not included in the fit, but are predicted well. Differences in the quality of fit are best seen in terms of the conditional mean of the error, estimated by the lowess (locally-weighted scatterplot smoothing) fit of the residuals (Supplementary Information). See Table 1 for the values of the coefficients obtained from the fit. **b**, Slope of the quadratic fit (including temperature) with pointwise 95% confidence intervals (blue). The slope of the power-law fit (red) and models with fixed 2/3 and 3/4 exponents (black) are included for comparison. This panel suggests that exponents estimated by assuming a power law will be highly sensitive to the mass range of the data set used, as shown in Fig. 2.



“This raises the question of whether the theory can be adapted to agree with the data”¹




Figure 2 | Scaling exponent depends on mass range. **a**, Slope estimated by linear regression within a three log-unit mass range (smaller near the boundaries). Values on the abscissa denote mean $\log_{10}M$ within the range. When the 95% confidence regions (dashed lines) include the 2/3 or 3/4 lines, the local slope is consistent with a 2/3 or 3/4 exponent, respectively. These cases are indicated by the shaded regions (2/3 on the left and 3/4 on the right). **b**, Slope estimated by using all data points with $M < x$. The shaded region is consistent with 2/3 slope estimates. **c**, Slope estimated by using all data points with $M > x$. The shaded region is consistent with 3/4 slope

estimates. **d**, Exponents estimated for eight historical data sets using linear regression (black filled circles): Lovegrove¹³, Lovegrove¹⁴, White¹⁸, White¹⁹, Sieg¹⁵, McNab¹⁶, and Savage¹⁷ using species average data ("Savage") and binned data ("Savage" bin). Exponents predicted using coefficients from quadratic fits to McNab's (red), Sieg's (green), or Savage's (blue) data and the first three moments of $\log_{10}M$ (Supplementary Information). Thick lines represent uncorrected 95% confidence intervals. Thin lines are multiplicity corrected intervals.

The PoCVerse
Optimal Supply
Networks II
71 of 125
Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument


Conclusion

References



¹Already raised and fully established 9 years earlier. [14]

Evolution has generally made things bigger¹



“The Phantom Tollbooth”  
by Norton Juster (1961). ^[25]



Regression starting at low M makes sense



Regression starting at high M makes ...no sense



¹Yes, yes, yes: insular dwarfism  with the shrinkage 

Still going:



“A general model for metabolic scaling in self-similar asymmetric networks” 


Brummer, Brummer, and Enquist,
PLoS Comput Biol, **13**, e1005394, 2017. [8]

Wut?:

“Most importantly, we show that the $3/4$ metabolic scaling exponent from Kleiber’s Law can still be attained within many asymmetric networks.”



Oh no:



“Scale: The Universal Laws of Growth, Innovation, Sustainability, and the Pace of Life in Organisms, Cities, Economies, and Companies”  
by Geoffrey B. West (2017). ^[53]

Amazon reviews excerpts (so, so not fair but ...):



“Full of intriguing, big ideas but amazingly sloppy both in details and exposition, especially considering the author is a theoretical physicist.”



“The beginning is terrible. He shows four graphs to illustrate scaling relationships, none of which have intelligible scales”



“(he actually repeats several times that businesses can die but are not really an animal - O RLY?)”

The PoCverse
Optimal Supply
Networks II
74 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument





Conclusion

References



Simple supply networks:

The PoCserve
Optimal Supply
Networks II
75 of 125
Metabolism and
Truthicide



Banavar et al.,
Nature,
(1999) ^[1].



Flow rate
argument.



Ignore
impedance.



Very general
attempt to find
most efficient
transportation
networks.

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Simple supply networks

The PoCVerse
Optimal Supply
Networks II
76 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Banavar *et al.* find ‘most efficient’ networks with

$$P \propto M^{d/(d+1)}$$



...but also find

$$V_{\text{network}} \propto M^{(d+1)/d}$$



$d = 3$:

$$V_{\text{blood}} \propto M^{4/3}$$



Consider a 3 g shrew with $V_{\text{blood}} = 0.1 V_{\text{body}}$



$\Rightarrow$ 3000 kg elephant with $V_{\text{blood}} = 10 V_{\text{body}}$





Geometric argument



“Optimal Form of Branching Supply and Collection Networks”

Peter Sheridan Dodds,

Phys. Rev. Lett., **104**, 048702, 2010. ^[12]


 Consider **one source** supplying **many sinks** in a **d -dim.** volume in a **D -dim.** ambient space.

 Assume **sinks** are **invariant**.

 Assume sink density **$\rho = \rho(V)$** .

 Assume some cap on flow speed of material.

 See network as a bundle of virtual vessels:



The PoCverse
Optimal Supply
Networks II
78 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Geometric argument

The PoCverse
Optimal Supply
Networks II
79 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

 **Q:** how does the number of sustainable sinks N_{sinks} scale with volume V for the most efficient network design?

 **Or:** what is the highest α for $N_{\text{sinks}} \propto V^\alpha$?



Geometric argument

The PoCverse
Optimal Supply
Networks II
80 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories


Geometric argument

Conclusion

References



Allometrically growing regions:



Have d length scales which scale as

$$L_i \propto V^{\gamma_i} \text{ where } \gamma_1 + \gamma_2 + \dots + \gamma_d = 1.$$



For **isometric** growth, $\gamma_i = 1/d$.



For **allometric** growth, we must have at least two of the $\{\gamma_i\}$ being different




Spherical cows and pancake cows:


Assume an isometrically Scaling family of cows:



Extremes of allometry:
The pancake cows—







Spherical cows and pancake cows:

 **Question:** How does the surface area S_{cow} of our two types of cows scale with cow volume V_{cow} ? [Insert assignment question ↗](#)

 **Question:** For general families of regions, how does surface area S scale with volume V ? [Insert assignment question ↗](#)



Geometric argument

The PoCserve
Optimal Supply
Networks II
85 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories


Geometric argument

Conclusion

References



Best and worst configurations (Banavar et al.)



Rather obviously:

$\min V_{\text{net}} \propto \sum \text{distances from source to sinks.}$



Minimal network volume:

Real supply networks are close to optimal:




Figure 1. (a) Commuter rail network in the Boston area. The arrow marks the assumed root of the network. (b) Star graph. (c) Minimum spanning tree. (d) The model of equation (3) applied to the same set of stations.

Gastner and Newman (2006): “Shape and efficiency in spatial distribution networks” [16]






“Rules for Biologically Inspired Adaptive Network Design”

Tero et al.,

Science, **327**, 439-442, 2010. ^[50]



Urban deslime in action:

<https://www.youtube.com/watch?v=GwKuFREOgmo>

The PoCverse
Optimal Supply
Networks II
87 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument


Conclusion

References



Minimal network volume:

We add one more element:




-  Vessel cross-sectional area may vary with distance from the source.
-  Flow rate increases as cross-sectional area decreases.
-  e.g., a collection network may have vessels tapering as they approach the central sink.
-  Find that vessel volume v must scale with vessel length ℓ to affect overall system scalings.



Minimal network volume:

Effecting scaling:



Consider vessel radius $r \propto (\ell + 1)^{-\epsilon}$, tapering from $r = r_{\text{max}}$ where $\epsilon \geq 0$.

Gives $v \propto \ell^{1-2\epsilon}$ if $\epsilon < 1/2$

Gives $v \propto 1 - \ell^{-(2\epsilon-1)} \rightarrow 1$ for large ℓ if $\epsilon > 1/2$

Previously, we looked at $\epsilon = 0$ only.



Minimal network volume:

For $0 \leq \epsilon < 1/2$, approximate network volume by integral over region:

$$\min V_{\text{net}} \propto \int_{\Omega_{d,D}(V)} \rho \|\vec{x}\|^{1-2\epsilon} d\vec{x}$$

Insert assignment question 

$$\propto \rho V^{1+\gamma_{\max}(1-2\epsilon)} \text{ where } \gamma_{\max} = \max_i \gamma_i.$$

For $\epsilon > 1/2$, find simply that

$$\min V_{\text{net}} \propto \rho V$$



So if supply lines can taper fast enough and without limit, minimum network volume can be made negligible.



For $0 \leq \epsilon < 1/2$:



$$\min V_{\text{net}} \propto \rho V^{1+\gamma_{\text{max}}(1-2\epsilon)}$$



If scaling is **isometric**, we have $\gamma_{\text{max}} = 1/d$:

$$\min V_{\text{net/iso}} \propto \rho V^{1+(1-2\epsilon)/d}$$



If scaling is **allometric**, we have $\gamma_{\text{max}} = \gamma_{\text{allo}} > 1/d$: and

$$\min V_{\text{net/allo}} \propto \rho V^{1+(1-2\epsilon)\gamma_{\text{allo}}}$$



Isometrically growing volumes **require less network volume** than allometrically growing volumes:

$$\frac{\min V_{\text{net/iso}}}{\min V_{\text{net/allo}}} \rightarrow 0 \text{ as } V \rightarrow \infty$$



For $\epsilon > 1/2$:



$$\min V_{\text{net}} \propto \rho V$$



Network volume scaling is now independent of overall shape scaling.

Limits to scaling



Can argue that ϵ must effectively be 0 for real networks over large enough scales.



Limit to how fast material can move, and how small material packages can be.



e.g., blood velocity and blood cell size.



Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

This
is a
really
clean
slide



Blood networks

The PoCverse
Optimal Supply
Networks II
94 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

 Velocity at capillaries and aorta approximately constant across body size^[52]: $\epsilon = 0$.

 **Material costly** $\Rightarrow$ expect lower optimal bound of $V_{\text{net}} \propto \rho V^{(d+1)/d}$ to be followed closely.

 For cardiovascular networks, $d = D = 3$.

 Blood volume scales linearly with body volume^[48], $V_{\text{net}} \propto V$.

 Sink density must $\therefore$ decrease as volume increases:

$$\rho \propto V^{-1/d}.$$

 Density of suppliable sinks **decreases** with organism size.



Blood networks

The PoCverse
Optimal Supply
Networks II
95 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

 Then P , the rate of overall energy use in Ω , can at most scale with volume as

$$P \propto \rho V \propto \rho M \propto M^{(d-1)/d}$$

 For $d = 3$ dimensional organisms, we have

$$P \propto M^{2/3}$$


 Including other constraints may raise scaling exponent to a higher, less efficient value.




 **Exciting bonus:** Scaling obtained by the supply network story and the surface-area law **only match** for isometrically growing shapes.

Insert assignment question 

The surface area—supply network mismatch for allometrically growing shapes:





Recall:

The PoCverse
Optimal Supply
Networks II
98 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

 The exponent $\alpha = 2/3$ works for all birds and mammals up to 10–30 kg

 For mammals $> 10\text{--}30$ kg, maybe we have a new scaling regime

 Economos: limb length break in scaling around 20 kg

 White and Seymour, 2005: unhappy with large herbivore measurements. Find $\alpha \simeq 0.686 \pm 0.014$



Prefactor:

Stefan-Boltzmann law: 



$$\frac{dE}{dt} = \sigma S T^4$$

where S is surface and T is temperature.



Very rough estimate of prefactor based on scaling of normal mammalian body temperature and surface area S :

$$B \simeq 10^5 M^{2/3} \text{erg/sec.}$$



Measured for $M \leq 10$ kg:

$$B = 2.57 \times 10^5 M^{2/3} \text{erg/sec.}$$



River networks

 View river networks as collection networks.

 Many sources and one sink.

 ϵ ?

 Assume ρ is constant over time and $\epsilon = 0$:

$$V_{\text{net}} \propto \rho V^{(d+1)/d} = \text{constant} \times V^{3/2}$$

 Network volume grows faster than basin 'volume' (really area).

 **It's all okay:**

Landscapes are $d=2$ surfaces living in $D=3$ dimensions.

 Streams can grow not just in width but in depth ...

 If $\epsilon > 0$, V_{net} will grow more slowly but $3/2$ appears to be confirmed from real data.



Hack's law

 Volume of water in river network can be calculated by adding up basin areas

 Flows sum in such a way that

$$V_{\text{net}} = \sum_{\text{all pixels}} a_{\text{pixel } i}$$

 Hack's law again:

$$\ell \sim a^h$$

 Can argue

$$V_{\text{net}} \propto V_{\text{basin}}^{1+h} = a_{\text{basin}}^{1+h}$$

where h is Hack's exponent.

 $\therefore$ minimal volume calculations gives

$$h = 1/2$$



Real data:



Banavar et al.'s approach ^[1] is okay because ρ really is constant.



The irony: shows optimal basins are isometric



Optimal Hack's law:
 $\ell \sim a^h$ with
 $h = 1/2$



(Zzzzz)





Figure 2 Allometric scaling in river networks. Double logarithmic plot of $C \propto \sum_{x \in A_x} A_x$ versus A for three river networks characterized by different climates, geology and geographic locations (Dry Fork, West Virginia, 586 km², digital terrain map (DTM) size 30 × 30 m²; Island Creek, Idaho, 260 km², DTM size 30 × 30 m²; Tirso, Italy, 2,024 km², DTM size 237 × 237 m²). The experimental points are obtained by binning total contributing areas, and computing the ensemble average of the sum of the inner areas for each sub-basin within the binned interval. The figure uses pixel units in which the smallest area element is assigned a unit value. Also plotted is the predicted scaling relationship with slope 3/2. The inset shows the raw data from the Tirso basin before any binning



Even better—prefactors match up:



The Cabal strikes back:

The PoCVerse
Optimal Supply
Networks II
104 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References

-  Banavar et al., 2010, PNAS:
“A general basis for quarter-power scaling in animals.” [2]
-  “It has been known for decades that the metabolic rate of animals scales with body mass with an exponent that is almost always < 1 , $> 2/3$, and often very close to $3/4$.”
-  Cough, cough, cough, hack, wheeze, cough.



Stories—Darth Quarter:

The PoCverse
Optimal Supply
Networks II
105 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Some people understand it's truly a disaster: ↗



Peter Sheridan Dodds, Theoretical Biology's Buzzkill

By Mark Changizi | February 9th 2010 03:24 PM | 1 comment | [Print](#) | [E-mail](#) | [Track Comments](#)

[RSS](#) [Share / Save](#) [f](#) [t](#) [v](#) [...](#) [Tweet](#) [Like](#)



Mark Changizi

Search This Blog

There is an apocryphal story about a graduate mathematics student at the University of Virginia studying the properties of certain mathematical objects. In his fifth year some killjoy bastard elsewhere published a paper proving that there are no such mathematical objects. He dropped out of the program, and I never did hear where he is today. He's probably making my cappuccino right now.

This week, a professor named Peter Sheridan Dodds published a new paper in *Physical Review Letters* further fleshing out a theory concerning why a $2/3$ power law may apply for metabolic rate. The $2/3$ law says that metabolic rate in animals rises as the $2/3$ power of body mass. It was in a 2001 *Journal of Theoretical Biology* paper that he first argued that perhaps a $2/3$ law applies, and that paper – along with others such as the one that just appeared – is what has put him in the Killjoy Hall of Fame. The University of Virginia's killjoy was a mere amateur.

Mark Changizi

MORE ARTICLES

- [The Ravenous Color-Blind: New Developments For Color-Deficients](#)
- [Don't Hold Your Breath Waiting For Artificial Brains](#)
- [Welcome To Humans, Version 3.0](#)

[All Articles](#)

ABOUT MARK

Mark Changizi is Director of Human Cognition at 2AI, and the author of *The Vision Revolution* (Benbella 2009) and *Harnessed: How...*

[View Mark's Profile](#)

The PoCVerse
Optimal Supply
Networks II
106 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



The unnecessary bafflement continues:

The PoCverse
Optimal Supply
Networks II
107 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References


“Testing the metabolic theory of ecology” [41]

C. Price, J. S. Weitz, V. Savage, J. Stegen, A. Clarke, D. Coomes, P. S. Dodds, R. Etienne, A. Kerkhoff, K. McCulloh, K. Niklas, H. Olff, and N. Swenson
Ecology Letters, **15**, 1465–1474, 2012.



Artisanal, handcrafted silliness:

“Critical truths about power laws”^[49]
Stumpf and Porter, Science, 2012



How good is your power law? The chart reflects the level of statistical support—as measured in (16, 21)—and our opinion about the mechanistic sophistication underlying hypothetical generative models for various reported power laws. Some relationships are identified by name; the others reflect the general characteristics of a wide range of reported

The PoCverse
Optimal Supply
Networks II
108 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



Conclusion

- Supply network story consistent with dimensional analysis.
- Isometrically growing regions can be more efficiently supplied than allometrically growing ones.
- Ambient and region dimensions matter ($D = d$ versus $D > d$).
- Deviations from optimal scaling suggest inefficiency (e.g., gravity for organisms, geological boundaries).
- Actual details of branching networks not that important.
- Exact nature of self-similarity varies.
- 2/3-scaling lives on, largely in hiding.
- 3/4-scaling? Jury ruled a mistrial.
- The truth will out. Maybe.



References I

- [1] J. R. Banavar, A. Maritan, and A. Rinaldo.
Size and form in efficient transportation networks.
Nature, 399:130–132, 1999. pdf ↗
- [2] J. R. Banavar, M. E. Moses, J. H. Brown, J. Damuth,
A. Rinaldo, R. M. Sibly, and A. Maritan.
A general basis for quarter-power scaling in animals.
Proc. Natl. Acad. Sci., 107:15816–15820, 2010. pdf ↗
- [3] P. Bennett and P. Harvey.
Active and resting metabolism in birds—allometry,
phylogeny and ecology.
J. Zool., 213:327–363, 1987. pdf ↗
- [4] K. L. Blaxter, editor.
Energy Metabolism; Proceedings of the 3rd symposium held
at Troon, Scotland, May 1964.
Academic Press, New York, 1965.



References II

- [5] J. J. Blum.
On the geometry of four-dimensions and the relationship
between metabolism and body mass.
[J. Theor. Biol.](#), 64:599–601, 1977. pdf ↗
- [6] S. Brody.
[Bioenergetics and Growth.](#)
Reinhold, New York, 1945.
reprint, . pdf ↗
- [7] J. H. Brown, G. B. West, and B. J. Enquist.
Yes, West, Brown and Enquist's model of allometric scaling
mathematically correct and biologically relevant?
[Functional Ecology](#), 19:735--738, 2005. pdf ↗



References III

- [8] A. B. Brummer, S. V. M., and B. J. Enquist.
A general model for metabolic scaling in self-similar
asymmetric networks.
[PLoS Comput Biol](#), 13, 2017. pdf 
- [9] E. Buckingham.
On physically similar systems: Illustrations of the use of
dimensional equations.
[Phys. Rev.](#), 4:345–376, 1914. pdf 
- [10] A. Clauset, C. R. Shalizi, and M. E. J. Newman.
Power-law distributions in empirical data.
[SIAM Review](#), 51:661–703, 2009. pdf 
- [11] M. H. DeGroot.
[Probability and Statistics](#).
Addison-Wesley, Reading, Massachusetts, 1975.



References IV

- [12] P. S. Dodds.
Optimal form of branching supply and collection networks.
[Phys. Rev. Lett.](#), 104(4):048702, 2010. pdf ↗
- [13] P. S. Dodds and D. H. Rothman.
Scaling, universality, and geomorphology.
[Annu. Rev. Earth Planet. Sci.](#), 28:571–610, 2000. pdf ↗
- [14] P. S. Dodds, D. H. Rothman, and J. S. Weitz.
Re-examination of the “3/4-law” of metabolism.
[Journal of Theoretical Biology](#), 209:9–27, 2001. pdf ↗
- [15] A. E. Economos.
Elastic and/or geometric similarity in mammalian design.
[Journal of Theoretical Biology](#), 103:167–172, 1983. pdf ↗



References V

- [16] M. T. Gastner and M. E. J. Newman.
Shape and efficiency in spatial distribution networks.
[J. Stat. Mech.: Theor. & Exp., 1:P01015, 2006. pdf](#) 
- [17] J. H. Gayford, D. J. Irschick, J. Martin, A. Chin, and J. L. Rummer.
The geometry of life: Testing the scaling of whole-organism surface area and volume using sharks.
[Royal Society Open Science, 12\(6\):242205, 2025. pdf](#) 
- [18] D. S. Glazier.
Beyond the '3/4-power law': variation in the intra- and interspecific scaling of metabolic rate in animals.
[Biol. Rev., 80:611–662, 2005. pdf](#) 



References VI

- [19] D. S. Glazier.

The $3/4$ -power law is not universal: Evolution of isometric, ontogenetic metabolic scaling in pelagic animals.

BioScience, 56:325–332, 2006. pdf ↗

- [20] J. T. Hack.

Studies of longitudinal stream profiles in Virginia and Maryland.

United States Geological Survey Professional Paper, 294-B:45–97, 1957. pdf ↗

- [21] A. Hemmingsen.

The relation of standard (basal) energy metabolism to total fresh weight of living organisms.

Rep. Steno Mem. Hosp., 4:1–58, 1950. pdf ↗



References VII

- [22] A. Hemmingsen.
Energy metabolism as related to body size and respiratory surfaces, and its evolution.
[Rep. Steno Mem. Hosp.](#), 9:1–110, 1960. pdf ↗
- [23] A. A. Heusner.
Size and power in mammals.
[Journal of Experimental Biology](#), 160:25–54, 1991. pdf ↗
- [24] M. R. Hirt, W. Jetz, B. C. Rall, and U. Brose.
A general scaling law reveals why the largest animals are not the fastest.
[Nature Ecology & Evolution](#), 1:1116, 2017. pdf ↗
- [25] N. Juster.
[The Phantom Tollbooth.](#)
Random House, 1961.



References VIII

- [26] M. Kleiber.
Body size and metabolism.
Hilgardia, 6:315–353, 1932. pdf ↗
- [27] M. Kleiber.
The Fire of Life. An Introduction to Animal Energetics.
Wiley, New York, 1961.
- [28] T. Kolokotronis, V. Savage, E. J. Deeds, and W. Fontana.
Curvature in metabolic scaling.
Nature, 464:753, 2010. pdf ↗
- [29] J. Kozłowski and M. Konarzewski.
Is West, Brown and Enquist’s model of allometric scaling
mathematically correct and biologically relevant?
Functional Ecology, 18:283–289, 2004. pdf ↗



References IX

- [30] M. LaBarbera.
Analyzing body size as a factor of ecology and evolution.
[Annu. Rev. Ecol. Syst.](#), 20:97–117, 1989. pdf ↗
- [31] N. Lane.
Power, Sex, Suicide: Mitochondria and the Meaning of Life.
Oxford University Press, Oxford, UK, 2005.
- [32] L. B. Leopold.
A View of the River.
Harvard University Press, Cambridge, MA, 1994.
- [33] T. McMahon.
Size and shape in biology.
[Science](#), 179:1201–1204, 1973. pdf ↗



References X

- [34] T. A. McMahon.
Allometry and biomechanics: Limb bones in adult ungulates.
[The American Naturalist](#), 109:547–563, 1975. pdf ↗
- [35] T. A. McMahon and J. T. Bonner.
On Size and Life.
Scientific American Library, New York, 1983.
- [36] N. Meyer-Vernet and J.-P. Rospars.
How fast do living organisms move: Maximum speeds from
bacteria to elephants and whales.
[American Journal of Physics](#), pages 719–722, 2015. pdf ↗
- [37] D. R. Montgomery and W. E. Dietrich.
Channel initiation and the problem of landscape scale.
[Science](#), 255:826–30, 1992. pdf ↗



References XI

- [38] C. D. Murray.
A relationship between circumference and weight in trees and
its bearing on branching angles.
J. Gen. Physiol., 10:725–729, 1927. [pdf](#) 
- [39] M. G. Newberry, E. D. B., and S. V. M.
Testing foundations of biological scaling theory using
automated measurements of vascular networks.
PLoS Comput Biol, 11:e1004455, 2015. [pdf](#) 
- [40] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P.
Flannery.
Numerical Recipes in C.
Cambridge University Press, second edition, 1992.



References XII

- [41] C. Price, J. S. Weitz, V. Savage, S. Stegen, A. Clarke, D. Coomes, P. S. Dodds, R. Etienne, A. Kerkhoff, K. McCulloh, K. Niklas, H. Olf, and N. Swenson. Testing the metabolic theory of ecology. [Ecology Letters](#), 15:1465–1474, 2012. pdf ↗
- [42] J. M. V. Rayner. Linear relations in biomechanics: the statistics of scaling functions. [J. Zool. Lond. \(A\)](#), 206:415–439, 1985. pdf ↗
- [43] M. Rubner. Ueber den einfluss der körpergrösse auf stoffund kraftwechsel. [Z. Biol.](#), 19:535–562, 1883. pdf ↗



References XIII

- [44] P. A. Samuelson.
A note on alternative regressions.
[Econometrica](#), 10:80–83, 1942. pdf ↗
- [45] Sarrus and Rameaux.
Rapport sur une mémoire adressé à l'Académie de Médecine.
[Bull. Acad. R. Méd. \(Paris\)](#), 3:1094–1100, 1838–39.
- [46] V. M. Savage, E. J. Deeds, and W. Fontana.
Sizing up allometric scaling theory.
[PLoS Computational Biology](#), 4:e1000171, 2008. pdf ↗
- [47] J. Speakman.
On Blum's four-dimensional geometric explanation for the
0.75 exponent in metabolic allometry.
[J. Theor. Biol.](#), 144(1):139–141, 1990. pdf ↗



References XIV

- [48] W. R. Stahl.
Scaling of respiratory variables in mammals.
[Journal of Applied Physiology](#), 22:453–460, 1967.
- [49] M. P. H. Stumpf and M. A. Porter.
Critical truths about power laws.
[Science](#), 335:665–666, 2012. [pdf](#) 
- [50] A. Tero, S. Takagi, T. Saigusa, K. Ito, D. P. Bebbler, M. D. Fricker, K. Yumiki, R. Kobayashi, and T. Nakagaki.
Rules for biologically inspired adaptive network design.
[Science](#), 327(5964):439–442, 2010. [pdf](#) 
- [51] D. L. Turcotte, J. D. Pelletier, and W. I. Newman.
Networks with side branching in biology.
[Journal of Theoretical Biology](#), 193:577–592, 1998. [pdf](#) 

The PoCVerse
Optimal Supply
Networks II
123 of 125

Metabolism and
Truthicide

Death by fractions

Measuring exponents

River networks

Earlier theories

Geometric argument

Conclusion

References



References XV

- [52] P. D. Weinberg and C. R. Ethier.
Twenty-fold difference in hemodynamic wall shear stress
between murine and human aortas.
[Journal of Biomechanics](#), 40(7):1594–1598, 2007. pdf ↗
- [53] G. B. West.
Scale: The Universal Laws of Growth, Innovation,
Sustainability, and the Pace of Life in Organisms, Cities,
Economies, and Companies.
Penguin Press, New York, 2017.
- [54] G. B. West, J. H. Brown, and B. J. Enquist.
A general model for the origin of allometric scaling laws in
biology.
[Science](#), 276:122–126, 1997. pdf ↗



References XVI

- [55] G. B. West, J. H. Brown, and J. Enquist.
The fourth dimension of life: Fractal geometry and allometric scaling of organisms.
[Science](#), 284:1677–1679, 1999. pdf ↗
- [56] G. B. West, J. H. Brown, and J. Enquist.
Nature.
[Nature](#), 400:664–667, 1999. pdf ↗
- [57] C. R. White and R. S. Seymour.
Allometric scaling of mammalian metabolism.
[J. Exp. Biol.](#), 208:1611–1619, 2005. pdf ↗

