

Biological Contagion

Principles of Complex Systems

Course 300, Fall, 2008

Prof. Peter Dodds

Department of Mathematics & Statistics
University of Vermont



Licensed under the *Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License*.

Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
References

Frame 1/58



Outline

Introduction

Simple disease spreading models
Background
Prediction

References

Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
References

Frame 2/58



Contagion

A confusion of contagions:

- ▶ Is Harry Potter some kind of virus?
- ▶ What about the Da Vinci Code?
- ▶ Does Sudoku spread like a disease?
- ▶ Religion?
- ▶ Democracy...?

Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
References

Frame 3/58



Contagion

Naturomorphisms

- ▶ “The feeling was contagious.”
- ▶ “The news spread like wildfire.”
- ▶ “Freedom is the most contagious virus known to man.”
—Hubert H. Humphrey, Johnson’s vice president
- ▶ “Nothing is so contagious as enthusiasm.”
—Samuel Taylor Coleridge

Biological
Contagion

Introduction
Simple disease
spreading models
Background
Prediction
References

Frame 4/58



Social contagion

Optimism according to Ambrose Bierce: (田)

The doctrine that everything is beautiful, including what is ugly, everything good, especially the bad, and everything right that is wrong. ... **It is hereditary, but fortunately not contagious.**

Social contagion

Eric Hoffer, 1902–1983

There is a grandeur in the uniformity of the mass. When a fashion, a dance, a song, a slogan or a joke sweeps like wildfire from one end of the continent to the other, and a hundred million people roar with laughter, sway their bodies in unison, hum one song or break forth in anger and denunciation, there is the overpowering feeling that in this country we have come nearer the brotherhood of man than ever before.

- ▶ Hoffer (田) was an interesting fellow...

The spread of fanaticism


Hoffer's acclaimed work:

"The True Believer:
Thoughts On The Nature Of Mass Movements" (1951) [3]

Quotes-aplenty:

- ▶ "We can be absolutely certain only about things we do not understand."
- ▶ "Mass movements can rise and spread without belief in a God, but never without belief in a devil."
- ▶ "Where freedom is real, equality is the passion of the masses. Where equality is real, freedom is the passion of a small minority."

Imitation




despair.com

"When people are free to do as they please, they usually imitate each other."

—Eric Hoffer
"The Passionate State of Mind" [4]

The collective...



despair.com

“Never Underestimate
the Power of Stupid
People in Large
Groups.”

Biological Contagion

Introduction

Simple disease
spreading models

Background
Prediction

References

Frame 9/58



Contagion

Definitions

- ▶ (1) The spreading of a quality or quantity between individuals in a population.
- ▶ (2) A disease itself: the plague, a blight, the dreaded lurgi, ...
- ▶ from Latin: *con* = 'together with' + *tangere* 'to touch.'
- ▶ Contagion has unpleasant overtones...
- ▶ Just **Spreading** might be a more neutral word
- ▶ But contagion is kind of exciting...

Biological Contagion

Introduction

Simple disease
spreading models

Background
Prediction

References

Frame 10/58



Examples of non-disease spreading:

Interesting infections:

- ▶ Spreading of buildings in the US. (田)
- ▶ Spreading of spreading (田).
- ▶ Viral get-out-the-vote video. (田)

Biological Contagion

Introduction

Simple disease
spreading models

Background
Prediction

References

Frame 11/58



Contagions

Two main classes of contagion

1. **Infectious diseases:** tuberculosis, HIV, ebola, SARS, influenza, ...
2. **Social contagion:** fashion, word usage, rumors, riots, religion, ...

Biological Contagion

Introduction

Simple disease
spreading models

Background
Prediction

References

Frame 12/58




Mathematical Epidemiology

The standard SIR model^[8]

- ▶ The basic model of disease contagion
- ▶ Three states:
 1. **S** = Susceptible
 2. **I** = Infective/Infectious
 3. **R** = Recovered or Removed or Refractory
- ▶ $S(t) + I(t) + R(t) = 1$
- ▶ Presumes random interactions (mass-action principle)
- ▶ Interactions are independent (no memory)
- ▶ Discrete and continuous time versions

Mathematical Epidemiology

Discrete time automata example:



Transition Probabilities:

- β for being infected given contact with infected
- r for recovery
- ρ for loss of immunity

Mathematical Epidemiology

Original models attributed to

- ▶ 1920's: Reed and Frost
- ▶ 1920's/1930's: Kermack and McKendrick^[5, 7, 6]
- ▶ Coupled differential equations with a mass-action principle

Independent Interaction models

Differential equations for continuous model

$$\frac{d}{dt} S = -\beta IS + \rho R$$

$$\frac{d}{dt} I = \beta IS - rI$$

$$\frac{d}{dt} R = rI - \rho R$$

β , r , and ρ are now **rates**.

Reproduction Number R_0 :

- ▶ R_0 = expected number of infected individuals resulting from a single initial infective
- ▶ Epidemic threshold: If $R_0 > 1$, 'epidemic' occurs.

Reproduction Number R_0

Discrete version:

- ▶ Set up: One Infective in a randomly mixing population of Susceptibles
- ▶ At time $t = 0$, single infective random bumps into a Susceptible
- ▶ Probability of transmission = β
- ▶ At time $t = 1$, single Infective remains infected with probability $1 - r$
- ▶ At time $t = k$, single Infective remains infected with probability $(1 - r)^k$

Reproduction Number R_0

Discrete version:

- ▶ Expected number infected by original Infective:

$$\begin{aligned} R_0 &= \beta + (1 - r)\beta + (1 - r)^2\beta + (1 - r)^3\beta + \dots \\ &= \beta \left(1 + (1 - r) + (1 - r)^2 + (1 - r)^3 + \dots \right) \\ &= \beta \frac{1}{1 - (1 - r)} = \beta/r \end{aligned}$$

For S_0 initial infectives ($1 - S_0 = R_0$ immune):

$$R_0 = S_0\beta/r$$

Independent Interaction models

For the continuous version

- ▶ Second equation:

$$\frac{d}{dt}I = \beta SI - rI$$

$$\frac{d}{dt}I = (\beta S - r)I$$


- ▶ Number of infectives grows initially if

$$\beta S(0) - r > 0 \Rightarrow \beta S(0) > r \Rightarrow \beta S(0)/r > 1$$

- ▶ Same story as for discrete model.

Independent Interaction models

Example of epidemic threshold:



- ▶ Continuous phase transition.
- ▶ Fine idea from a simple model.

Independent Interaction models

Many variants of the SIR model:

- ▶ **SIS**: susceptible-infective-susceptible
- ▶ **SIRS**: susceptible-infective-recovered-susceptible
- ▶ compartment models (age or gender partitions)
- ▶ more categories such as 'exposed' (**SEIRS**)
- ▶ recruitment (migration, birth)

Biological Contagion

Introduction

Simple disease spreading models

Background

Prediction

References

Frame 22/58

Disease spreading models

For novel diseases:

1. Can we predict the size of an epidemic?
2. How important is the reproduction number R_0 ?

Biological Contagion

Introduction

Simple disease spreading models

Background

Prediction

References

Frame 24/58

R_0 and variation in epidemic sizes

R_0 approximately same for all of the following:

- ▶ 1918-19 "Spanish Flu" ~ 500,000 deaths in US
- ▶ 1957-58 "Asian Flu" ~ 70,000 deaths in US
- ▶ 1968-69 "Hong Kong Flu" ~ 34,000 deaths in US
- ▶ 2003 "SARS Epidemic" ~ 800 deaths world-wide

Biological Contagion

Introduction

Simple disease spreading models

Background

Prediction

References

Frame 25/58

Size distributions

Size distributions are important elsewhere:

- ▶ earthquakes (Gutenberg-Richter law)
- ▶ city sizes, forest fires, war fatalities
- ▶ wealth distributions
- ▶ 'popularity' (books, music, websites, ideas)
- ▶ **Epidemics?**

Power laws distributions are common but not obligatory...

Biological Contagion

Introduction

Simple disease spreading models

Background

Prediction

References

Frame 26/58


Size distributions

Really, what about epidemics?

- ▶ Simply hasn't attracted much attention.
- ▶ Data not as clean as for other phenomena.

Feeling Ill in Iceland


Caseload recorded monthly for range of diseases in Iceland, 1888-1890



Treat outbreaks separated in time as 'novel' diseases.


Really not so good at all in Iceland

Epidemic size distributions $N(S)$ for Measles, Rubella, and Whooping Cough.



Spike near $S = 0$, relatively flat otherwise.

Measles & Pertussis



Insert plots:
Complementary cumulative frequency distributions:

$$N(\psi' > \psi) \propto \psi^{-\gamma+1}$$

Limited scaling with a possible break.

Power law distributions

Measured values of γ :

- ▶ measles: **1.40** (low Ψ) and **1.13** (high Ψ)
- ▶ pertussis: **1.39** (low Ψ) and **1.16** (high Ψ)


- ▶ Expect $2 \leq \gamma < 3$ (finite mean, infinite variance)
- ▶ When $\gamma < 1$, can't normalize
- ▶ Distribution is quite **flat**.

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 31/58

Resurgence—example of SARS



- ▶ Epidemic slows... then an infective moves to a new context.
- ▶ Epidemic discovers new 'pools' of susceptibles: **Resurgence**.
- ▶ **Importance of rare, stochastic events.**

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 32/58

The challenge

So... can a simple model produce


1. **broad epidemic distributions** and
2. **resurgence** ?

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 33/58

Size distributions



Simple models typically produce **bimodal** or **unimodal** size distributions.

- ▶ This **includes** network models: random, small-world, scale-free, ...
- ▶ Exceptions:
 1. Forest fire models
 2. Sophisticated metapopulation models

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 34/58

Burning through the population

Forest fire models:^[9]

- ▶ Rhodes & Anderson, 1996
- ▶ The physicist's approach:
"if it works for magnets, it'll work for people..."

A bit of a stretch:


1. Epidemics \equiv forest fires spreading on 3-d and 5-d lattices.
2. Claim Iceland and Faroe Islands exhibit power law distributions for outbreaks.
3. Original forest fire model not completely understood.

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 35/58

Size distributions



From Rhodes and Anderson, 1996.

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 36/58

Sophisticated metapopulation models

- ▶ Community based mixing: Longini (two scales).
- ▶ Eubank et al.'s EpiSims/TRANSIMS—city simulations.
- ▶ Spreading through countries—Airlines: Germann et al., Corlizza et al.
- ▶ Vital work but perhaps hard to generalize from...
- ▶ \Rightarrow Create a simple model involving multiscale travel
- ▶ Multiscale models suggested by others but not formalized (Bailey, Cliff and Haggett, Ferguson et al.)

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 37/58

Size distributions

- ▶ Very big question: **What is N ?**
- ▶ Should we model SARS in Hong Kong as spreading
- ▶ For simple models, we need to know the final size beforehand...


Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 38/58

Improving simple models

Contexts and Identities—Bipartite networks



- ▶ boards of directors
- ▶ movies
- ▶ transportation modes (subway)

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 39/58

Improving simple models

Idea for social networks: incorporate **identity**.

Identity is formed from attributes such as:

- ▶ Geographic location
- ▶ Type of employment
- ▶ Age
- ▶ Recreational activities

Groups are crucial...


- ▶ formed by people with at least one similar attribute
- ▶ Attributes \Leftrightarrow Contexts \Leftrightarrow Interactions \Leftrightarrow Networks. ^[11]

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 40/58

Infer interactions/network from identities




Distance makes sense in identity/context space.

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 41/58

Generalized context space



(Blau & Schwartz ^[1], Simmel ^[10], Breiger ^[2])

Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 42/58

A toy agent-based model

Geography—allow people to move between contexts:

- ▶ Locally: standard SIR model with random mixing
- ▶ discrete time simulation
- ▶ β = infection probability
- ▶ γ = recovery probability
- ▶ P = probability of travel
- ▶ **Movement distance:** $\Pr(d) \propto \exp(-d/\xi)$
- ▶ ξ = typical travel distance


Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 43/58

A toy agent-based model

Schematic:



Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 44/58

Model output

- ▶ Define P_0 = Expected number of infected individuals **leaving** initially infected context.
- ▶ Need $P_0 > 1$ for disease to spread (independent of R_0).
- ▶ Limit epidemic size by **restricting frequency of travel and/or range**


Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 45/58

Model output

Varying ξ :



- ▶ Transition in expected final size based on typical movement distance (**sensible**)


Biological Contagion

Introduction
Simple disease spreading models
Background
Prediction
References

Frame 46/58


Model output

Varying P_0 :



- ▶ Transition in expected final size based on typical number of infectives leaving first group (also sensible)
- ▶ Travel advisories: ξ has larger effect than P_0 .


Example model output: size distributions



- ▶ Flat distributions are possible for certain ξ and P .
- ▶ Different R_0 's may produce similar distributions
- ▶ Same epidemic sizes may arise from different R_0 's


Model output—resurgence

Standard model:



Model output—resurgence

Standard model with transport:



The upshot

Multiscale population structure

+
stochasticity

leads to

resurgence

+
broad epidemic size distributions

Conclusions

- ▶ For this model, epidemic size is highly unpredictable
- ▶ Model is more complicated than SIR but still simple
- ▶ We haven't even included normal social responses such as travel bans and self-quarantine.
- ▶ The reproduction number R_0 is not very useful.
- ▶ R_0 , however measured, is not informative about
 1. how likely the observed epidemic size was,
 2. and how likely future epidemics will be.
- ▶ Problem: R_0 summarises **one** epidemic after the fact and enfolds movement, everything.

Conclusions

- ▶ Disease spread highly sensitive to population structure
- ▶ Rare events may matter enormously (e.g., an infected individual taking an international flight)
- ▶ More support for controlling population movement (e.g., travel advisories, quarantine)

Conclusions

What to do:

- ▶ Need to separate movement from disease
- ▶ R_0 needs a friend or two.
- ▶ Need $R_0 > 1$ and $P_0 > 1$ and ξ sufficiently large for disease to have a chance of spreading

More wondering:

- ▶ Exactly how important are rare events in disease spreading?
- ▶ Again, what is N ?

Simple disease spreading models

Attempts to use beyond disease:

- ▶ Adoption of ideas/beliefs (Goffman & Newell, 1964)
- ▶ Spread of rumors (Daley & Kendall, 1965)
- ▶ Diffusion of innovations (Bass, 1969)
- ▶ Spread of fanatical behavior (Castillo-Chávez & Song, 2003)

References I

- P. M. Blau and J. E. Schwartz.
Crosscutting Social Circles.
Academic Press, Orlando, FL, 1984.
- R. L. Breiger.
The duality of persons and groups.
Social Forces, 53(2):181–190, 1974.
- E. Hoffer.
The True Believer: On The Nature Of Mass Movements.
Harper and Row, New York, 1951.
- E. Hoffer.
The Passionate State of Mind: And Other Aphorisms.
Buccaneer Books, 1954.

References II

- W. O. Kermack and A. G. McKendrick.
A contribution to the mathematical theory of epidemics.
Proc. R. Soc. Lond. A, 115:700–721, 1927. [pdf](#) (田)
- W. O. Kermack and A. G. McKendrick.
A contribution to the mathematical theory of epidemics. III. Further studies of the problem of endemicity.
Proc. R. Soc. Lond. A, 141(843):94–122, 1927.
[pdf](#) (田)
- W. O. Kermack and A. G. McKendrick.
Contributions to the mathematical theory of epidemics. II. The problem of endemicity.
Proc. R. Soc. Lond. A, 138(834):55–83, 1927. [pdf](#) (田)

References III

- J. D. Murray.
Mathematical Biology.
Springer, New York, Third edition, 2002.
- C. J. Rhodes and R. M. Anderson.
Power laws governing epidemics in isolated populations.
Nature, 381:600–602, 1996. [pdf](#) (田)
- G. Simmel.
The number of members as determining the sociological form of the group. I.
American Journal of Sociology, 8:1–46, 1902.
- D. J. Watts, P. S. Dodds, and M. E. J. Newman.
Identity and search in social networks.
Science, 296:1302–1305, 2002. [pdf](#) (田)