Scaling—a Plenitude of Power Laws

Principles of Complex Systems
Course CSYS/MATH 300, Fall, 2009

Prof. Peter Dodds

Dept. of Mathematics & Statistics
Center for Complex Systems :: Vermont Advanced Computing Center
University of Vermont

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Outline

Definitions

General observation:

Systems (complex or not)
that cross many spatial and temporal scales
often exhibit some form of scaling.

Outline

All about scaling:

- ▶ Definitions.
- ▶ Examples.
- ▶ How to measure your power-law relationship.
- ▶ Mechanisms giving rise to your power-laws.

Definitions

A power law relates two variables *x* and *y* as follows:

$$y = cx^{\alpha}$$

- ► α is the scaling exponent (or just exponent)
- ► (α can be any number in principle but we will find various restrictions.)
- ► *c* is the prefactor (which can be important!)

Definitions

- The prefactor *c* must balance dimensions.
- eg., length ℓ and volume *v* of common nails are related as:

$$\ell = c v^{1/4}$$

- Using $[\cdot]$ to indicate dimension, then

$$[c] = [l]/[V^{1/4}] = L/L^{3/4} = L^{1/4}.$$

Looking at data

▶ Power-law relationships are linear in log-log space:

$$y = cx^{\alpha}$$

$$\Rightarrow \log_b y = \alpha \log_b x + \log_b c$$

with slope equal to α, the scaling exponent.

▶ Much searching for straight lines on log-log or double-logarithmic plots.

▶ Good practice: Always, always, always use base 10.

▶ Talk only about orders of magnitude (powers of 10).

A beautiful, heart-warming example:

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 8/117

$\alpha \simeq 1.23$

gray matter: 'computing elements'

white matter: 'wiring'

from Zhang & Sejnowski, PNAS (2000) [26]

Why is $\alpha \simeq 1.23$?

Quantities (following Zhang and Sejnowski):

- G = Volume of gray matter (cortex/processors)
- W = Volume of white matter (wiring)
- T = Cortical thickness (wiring)
- S = Cortical surface area
- L = Average length of white matter fibers
- p = density of axons on white matter/cortex interface

A rough understanding:

- $G \sim ST$ (convolutions are okay)
- $W \sim \frac{1}{2} pSL$
- $G \sim L^3$ ← this is a little sketchy...
- Eliminate S and L to find $W \propto G^{4/3}/T$

Why is $\alpha \simeq 1.23$?

A rough understanding:

- We are here: $W \propto G^{4/3}/T$
- Observe weak scaling $T \propto G^{0.10 \pm 0.02}$.
- (Implies $S \propto G^{0.9} \rightarrow$ convolutions fill space.)
- $\Rightarrow W \propto G^{4/3}/T \propto G^{1.23 \pm 0.02}$

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Why is $\alpha \simeq 1.23$?

Figure showing:
$\log_{10} G = (0.955 \pm 0.002) \log_{10} V + (0.061 \pm 0.009)$
$r = 0.9998$

Gray Matter Volume G

White Matter Volume W

$\log_{10} W = (1.174 \pm 0.007) \log_{10} V - (1.40 \pm 0.03)$
$r = 0.9988$

(axes: Volume in mm^3 vs Total Volume $V = G + W$ (mm^3))

Trickiness:

- With $V = G + W$, some power laws must be approximations.
- Measuring exponents is a hairy business...

Good scaling:

General rules of thumb:

- *High quality:* scaling persists over three or more orders of magnitude for each variable.

- *Medium quality:* scaling persists over three or more orders of magnitude for only one variable and at least one for the other.

- *Very dubious:* scaling 'persists' over less than an order of magnitude for both variables.

Unconvincing scaling:

Average walking speed as a function of city population:

$\beta=0.093$ $R^2=0.80$

Two problems:

1. use of natural log, and
2. minute varation in dependent variable.

from Bettencourt et al. (2007) [3]; otherwise very interesting!

Definitions

Power laws are the signature of
scale invariance:

Scale invariant 'objects'
look the 'same'
when they are appropriately
rescaled.

- ▶ Objects = geometric shapes, time series, functions, relationships, distributions,...
- ▶ 'Same' might be 'statistically the same'
- ▶ To rescale means to change the units of measurement for the relevant variables

Scale invariance

Our friend $y = cx^\alpha$:

- If we rescale x as $x = rx'$ and y as $y = r^\alpha y'$,
- then

$$r^\alpha y' = c(rx')^\alpha$$

-

$$\Rightarrow y' = cr^\alpha x'^\alpha r^{-\alpha}$$

-

$$\Rightarrow y' = cx'^\alpha$$

Scale invariance

Compare with $y = ce^{-\lambda x}$:

▶ If we rescale x as $x = rx'$, then

$$y = ce^{-\lambda r x'}$$

▶ Original form cannot be recovered.
▶ $\Rightarrow$ scale matters for the exponential.

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Scale invariance

More on $y = ce^{-\lambda x}$:

- Say $x_0 = 1/\lambda$ is the characteristic scale.
- For $x \gg x_0$, y is small,
  while for $x \ll x_0$, y is large.
- $\Rightarrow$ More on this later with size distributions.

Definitions

Allometry (⊞):

[refers to] differential growth rates of the parts of a living organism's body part or process.

Definitions:

Isometry:
dimensions scale
linearly with each
other.

Allometry:
dimensions scale
nonlinearly.

Definitions

Isometry versus Allometry:

- Isometry = 'same measure'
- Allometry = 'other measure'

Confusingly, we use allometric scaling to refer to both:

1. nonlinear scaling (e.g., $x \propto y^{1/3}$)
2. and the relative scaling of different measures (e.g., resting heart rate as a function of body size)

A wonderful treatise on scaling:

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

ON SIZE AND LIFE

THOMAS A. McMAHON AND JOHN TYLER BONNER

McMahon and
Bonner, 1983[18]

For the following slide:

Scaling

Scaling-at-large

Allometry

Definitions

Examples

History: Metabolism

Measuring exponents

History: River networks

Earlier theories

Geometric argument

Blood networks

River networks

Conclusion

References

The biggest living things (*left*). All the organisms are drawn to the same scale. *1*, The largest flying bird (albatross); *2*, the largest known animal (the blue whale), *3*, the largest extinct land mammal (*Baluchitherium*) with a human figure shown for scale; *4*, the tallest living land animal (giraffe); *5*, *Tyrannosaurus*; *6*, *Diplodocus*; *7*, one of the largest flying reptiles (*Pteranodon*); *8*, the largest extinct snake; *9*, the length of the largest tapeworm found in man; *10*, the largest living reptile (West African crocodile); *11*, the largest extinct lizard; *12*, the largest extinct bird (*Aepyornis*); *13*, the largest jellyfish (*Cyanea*); *14*, the largest living lizard (Komodo dragon); *15*, sheep; *16*, the largest bivalve mollusc (*Tridacna*); *17*; the largest fish (whale shark); *18*, horse; *19*, the largest crustacean (Japanese spider crab); *20*, the largest sea scorpion (Eurypterid); *21*, large tarpon; *22*, the largest lobster; *23*, the largest mollusc (deep-water squid, *Architeuthis*); *24*, ostrich; *25*, the lower 105 feet of the largest organism (giant sequoia), with a 100-foot larch superposed.

p. 2, McMahon and Bonner[18]

The many scales of life:

p. 2, McMahon and Bonner [18]

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

For the following slide:

Medium-sized creatures (*above*). *1*, Dog; *2*, common herring; *3*, the largest egg (*Aepyornis*); *4*, song thrush with egg; *5*, the smallest bird (hummingbird) with egg; *6*, queen bee; *7*, common cockroach; *8*, the largest stick insect; *9*, the largest polyp (*Branchiocerianthus*); *10*, the smallest mammal (flying shrew); *11*, the smallest vertebrate (a tropical frog); *12*, the largest frog (goliath frog); *13*, common grass frog; *14*, house mouse; *15*, the largest land snail (*Achatina*) with egg; *16*, common snail; *17*, the largest beetle (goliath beetle); *18*, human hand; *19*, the largest starfish (*Luidia*); *20*, the largest free-moving protozoan (an extinct nummulite).

p. 2, McMahon and Bonner[18]

The many scales of life:

p. 3, McMahon and Bonner[18]

For the following slide:

Scaling

Scaling-at-large

Allometry

Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Small, "naked-eye" creatures (*lower left*).
1, One of the smallest fishes (*Trimmatom nanus*); *2*, common brown hydra, expanded; *3*, housefly; *4*, medium-sized ant; *5*, the smallest vertebrate (a tropical frog, the same as the one numbered *11* in the figure *above*); *6*, flea (*Xenopsylla cheopis*); *7*, the smallest land snail; *8*, common water flea (*Daphnia*).

The smallest "naked-eye" creatures and some large microscopic animals and cells (*below right*). *1*, *Vorticella*, a ciliate; *2*, the largest ciliate protozoan (*Bursaria*); *3*, the smallest many-celled animal (a rotifer); *4*, smallest flying insect (*Elaphis*); *5*, another ciliate (*Paramecium*); *6*, cheese mite; *7*, human sperm; *8*, human ovum; *9*, dysentery amoeba; *10*, human liver cell; *11*, the foreleg of the flea (numbered *6* in the figure to the *left*).

p. 2, McMahon and Bonner[18]

The many scales of life:

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 29/117

p. 3, McMahon and Bonner[18]

Size range and cell differentiation:

Non-uniform growth:

years
0 • 42   0 • 75   2 • 75   6 • 75   12 • 75   25 • 75

p. 32, McMahon and Bonner [18]

Non-uniform growth—arm length versus height:

Good example of a break in scaling:

A crossover in scaling occurs around a height of 1 metre.

p. 32, McMahon and Bonner [18]

Weightlifting: $M_{\text{worldrecord}} \propto M_{\text{lifter}}^{2/3}$

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Idea: Power $\sim$ cross-sectional area of isometric lifters.

p. 53, McMahon and Bonner[18]

Titanothere horns: $L_{\mathrm{horn}} \sim L_{\mathrm{skull}}^4$

p. 36, McMahon and Bonner[18]

The allometry of nails:

- ▶ Diameter $\propto$ Mass$^{3/8}$
- ▶ Length $\propto$ Mass$^{1/4}$
- ▶ Diameter $\propto$ Length$^{2/3}$

p. 58–59, McMahon and Bonner[18]

The allometry of nails:

A buckling instability?:

- ▶ Physics/Engineering result: Columns buckle under a load which depends on d^4/ℓ^2.
- ▶ To drive nails in, resistive force $\propto$ nail circumference $= \pi d$.
- ▶ Match forces independent of nail size: $d^4/\ell^2 \propto d$.
- ▶ Leads to $d \propto \ell^{2/3}$.

Rowing: Speed $\propto$ (number of rowers)$^{1/9}$

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 37/117

Shell dimensions and performances.

No. of oarsmen	Modifying description	Length, l (m)	Beam, b (m)	l/b	Boat mass per oarsman (kg)	Time for 2000 m (min)			
						I	II	III	IV
8	Heavyweight	18.28	0.610	30.0	14.7	5.87	5.92	5.82	5.73
8	Lightweight	18.28	0.598	30.6	14.7				
4	With coxswain	12.80	0.574	22.3	18.1				
4	Without coxswain	11.75	0.574	21.0	18.1	6.33	6.42	6.48	6.13
2	Double scull	9.76	0.381	25.6	13.6				
2	Pair-oared shell	9.76	0.356	27.4	13.6	6.87	6.92	6.95	6.77
1	Single scull	7.93	0.293	27.0	16.3	7.16	7.25	7.28	7.17

Scaling in Cities:

- ▶ "Growth, innovation, scaling, and the pace of life in cities"
  Bettencourt et al., PNAS, 2007. [3]
- ▶ Quantified levels of
  - ▶ Infrastructure
  - ▶ Wealth
  - ▶ Crime levels
  - ▶ Disease
  - ▶ Energy consumption

  as a function of city size *N* (population).

Scaling in Cities:

Table 1. Scaling exponents for urban indicators vs. city size

Y	β	95% CI	Adj-R^2	Observations	Country–year
New patents	1.27	[1.25,1.29]	0.72	331	U.S. 2001
Inventors	1.25	[1.22,1.27]	0.76	331	U.S. 2001
Private R&D employment	1.34	[1.29,1.39]	0.92	266	U.S. 2002
"Supercreative" employment	1.15	[1.11,1.18]	0.89	287	U.S. 2003
R&D establishments	1.19	[1.14,1.22]	0.77	287	U.S. 1997
R&D employment	1.26	[1.18,1.43]	0.93	295	China 2002
Total wages	1.12	[1.09,1.13]	0.96	361	U.S. 2002
Total bank deposits	1.08	[1.03,1.11]	0.91	267	U.S. 1996
GDP	1.15	[1.06,1.23]	0.96	295	China 2002
GDP	1.26	[1.09,1.46]	0.64	196	EU 1999–2003
GDP	1.13	[1.03,1.23]	0.94	37	Germany 2003
Total electrical consumption	1.07	[1.03,1.11]	0.88	392	Germany 2002
New AIDS cases	1.23	[1.18,1.29]	0.76	93	U.S. 2002–2003
Serious crimes	1.16	[1.11, 1.18]	0.89	287	U.S. 2003
Total housing	1.00	[0.99,1.01]	0.99	316	U.S. 1990
Total employment	1.01	[0.99,1.02]	0.98	331	U.S. 2001
Household electrical consumption	1.00	[0.94,1.06]	0.88	377	Germany 2002
Household electrical consumption	1.05	[0.89,1.22]	0.91	295	China 2002
Household water consumption	1.01	[0.89,1.11]	0.96	295	China 2002
Gasoline stations	0.77	[0.74,0.81]	0.93	318	U.S. 2001
Gasoline sales	0.79	[0.73,0.80]	0.94	318	U.S. 2001
Length of electrical cables	0.87	[0.82,0.92]	0.75	380	Germany 2002
Road surface	0.83	[0.74,0.92]	0.87	29	Germany 2002

Data sources are shown in *SI Text*. CI, confidence interval; Adj-R^2, adjusted R^2; GDP, gross domestic product.

Scaling in Cities:

Intriguing findings:

- ▶ Global supply costs scale sublinearly with N ($\beta < 1$).
    - ▶ Returns to scale for infrastructure.
- ▶ Total individual costs scale linearly with N ($\beta = 1$)
    - ▶ Individuals consume similar amounts independent of city size.
- ▶ Social quantities scale superlinearly with N ($\beta > 1$)
    - ▶ Creativity (# patents), wealth, disease, crime, ...

Density doesn't seem to matter...

- ▶ Surprising given that across the world, we observe two orders of magnitude variation in area covered by agglomerations of fixed populations.

Ecology—Species-area law: $N_{species} \propto A^\beta$

Allegedly (data is messy):

- On islands: $\beta \approx 1/4$.
- On continuous land: $\beta \approx 1/8$.

A focus:

- How much energy do organisms need to live?
- And how does this scale with organismal size?

Animal power

Fundamental biological and ecological constraint:

$$\boxed{P = c\,M^{\alpha}}$$

$P =$ basal metabolic rate

$M =$ organismal body mass

$$P = c M^{\alpha}$$

Prefactor c depends on body plan and body temperature:

Birds	39–41 $^{\circ}C$
Eutherian Mammals	36–38 $^{\circ}C$
Marsupials	34–36 $^{\circ}C$
Monotremes	30–31 $^{\circ}C$

What one might expect:

$\alpha = 2/3$ because ...

- ▶ Dimensional analysis suggests an energy balance surface law:

$$P \propto S \propto V^{2/3} \propto M^{2/3}$$

- ▶ Lognormal fluctuations:
  Gaussian fluctuations in $\log P$ around $\log cM^{\alpha}$.

- ▶ Stefan-Boltzmann relation for radiated energy:

$$\frac{\mathrm{d}E}{\mathrm{d}t} = \sigma \varepsilon S T^4$$

The prevailing belief of the church of quarterology

$$\boxed{\alpha = 3/4}$$

$$P \propto M^{3/4}$$

Huh?

Related putative scalings:

- number of capillaries $\propto M^{3/4}$
- time to reproductive maturity $\propto M^{1/4}$
- heart rate $\propto M^{-1/4}$
- cross-sectional area of aorta $\propto M^{3/4}$
- population density $\propto M^{-3/4}$

The great 'law' of heartbeats:

Assuming:

- Average lifespan $\propto M^{\beta}$
- Average heart rate $\propto M^{-\beta}$
- Irrelevant but perhaps $\beta = 1/4$.

Then:

- Average number of heart beats in a lifespan
  $\simeq$ (Average lifespan) $\times$ (Average heart rate)
  $$\propto M^{\beta-\beta}$$
  $$\propto M^0$$
- Number of heartbeats per life time is independent of organism size!
- ≈ 1.5 billion....

History

1840's: Sarrus and Rameaux [22] first suggested $\alpha = 2/3$.

History

1883: Rubner [21] found $\alpha \simeq 2/3$.

History

1930's: Brody, Benedict study mammals. [6]
Found $\alpha \simeq 0.73$ (standard).

History

1932: Kleiber analyzed 13 mammals. [15]
Found $\alpha = 0.76$ and suggested $\alpha = 3/4$.

History

1950/1960: Hemmingsen [12, 13]
Extension to unicellular organisms.
$\alpha = 3/4$ assumed true.

History

1964: Troon, Scotland: [4]
3rd symposium on energy metabolism.
$\alpha = 3/4$ made official . . .   . . . 29 to zip.

Today

▶ 3/4 is held by many to be the one true exponent.

In the Beat of a Heart: Life, Energy, and the Unity of Nature—by John Whitfield

▶ But—much controversy...

▶ See 'Re-examination of the "3/4-law" of metabolism' Dodds, Rothman, and Weitz [9]

Some data on metabolic rates

$B = 0.026\,M^{0.668}$

[10-Dec-2001 peter dodds]

$\log_{10} M$

$\log_{10} B$

(source: brunehaldeworkbiologyallometryhesunerfigureofigheusner91.ps)

- ▶ Heusner's data (1991) [14]
- ▶ 391 Mammals
- ▶ blue line: 2/3
- ▶ red line: 3/4.
- ▶ ($B = P$)

Some data on metabolic rates

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

$B = 0.041 \, M^{0.664}$

- Bennett and Harvey's data (1987) [2]
- 398 birds
- blue line: 2/3
- red line: 3/4.
- ($B = P$)

Passerine vs. non-passerine...

Linear regression

Important:

- ► Ordinary Least Squares (OLS) Linear regression is only appropriate for analyzing a dataset $\{(x_i, y_i)\}$ when we know the x_i are measured without error.
- ► Here we assume that measurements of mass M have less error than measurements of metabolic rate B.
- ► Linear regression assumes Gaussian errors.

Measuring exponents

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

More on regression:

If (a) we don't know what the errors of either variable are,

or (b) no variable can be considered independent,

then we need to use
Standardized Major Axis Linear Regression.

(aka Reduced Major Axis = RMA.)

Measuring exponents

For Standardized Major Axis Linear Regression:

$$\text{slope}_{\text{SMA}} = \frac{\text{standard deviation of } y \text{ data}}{\text{standard deviation of } x \text{ data}}$$

Very simple!

Measuring exponents

Relationship to ordinary least squares regression is simple:

$$\begin{aligned}\text{slope}_{\text{SMA}} &= r^{-1} \times \text{slope}_{\text{OLS } y \text{ on } x} \\ &= r \times \text{slope}_{\text{OLS } x \text{ on } y}\end{aligned}$$

where r = standard correlation coefficient:

$$r = \frac{\sum_{i=1}^{n}(x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^{n}(x_i - \bar{x})^2}\sqrt{\sum_{i=1}^{n}(y_i - \bar{y})^2}}$$

Heusner's data, 1991 (391 Mammals)

range of M	N	$\hat{\alpha}$
≤ 0.1 kg	167	0.678 ± 0.038
≤ 1 kg	276	0.662 ± 0.032
≤ 10 kg	357	0.668 ± 0.019
≤ 25 kg	366	0.669 ± 0.018
≤ 35 kg	371	0.675 ± 0.018
≤ 350 kg	389	0.706 ± 0.016
≤ 3670 kg	391	0.710 ± 0.021

Bennett and Harvey, 1987 (398 birds)

$M_{\max}$	N	$\hat{\alpha}$
≤ 0.032	162	0.636 ± 0.103
≤ 0.1	236	0.602 ± 0.060
≤ 0.32	290	0.607 ± 0.039
≤ 1	334	0.652 ± 0.030
≤ 3.2	371	0.655 ± 0.023
≤ 10	391	0.664 ± 0.020
≤ 32	396	0.665 ± 0.019
≤ 100	398	0.664 ± 0.019

Hypothesis testing

Test to see if α' is consistent with our data $\{(M_i, B_i)\}$:

$$H_0 : \alpha = \alpha' \text{ and } H_1 : \alpha \neq \alpha'.$$

- Assume each $\mathbf{B}_i$ (now a random variable) is normally distributed about $\alpha' \log_{10} M_i + \log_{10} c$.
- Follows that the measured α for one realization obeys a t distribution with $N - 2$ degrees of freedom.
- Calculate a p-value: probability that the measured α is as least as different to our hypothesized α' as we observe.
- (see, for example, DeGroot and Scherish, "Probability and Statistics"[7])

Revisiting the past—mammals

Full mass range:

	N	$\hat{\alpha}$	$p_{2/3}$	$p_{3/4}$
Kleiber	13	0.738	$< 10^{-6}$	0.11
Brody	35	0.718	$< 10^{-4}$	$< 10^{-2}$
Heusner	391	0.710	$< 10^{-6}$	$< 10^{-5}$
Bennett and Harvey	398	0.664	0.69	$< 10^{-15}$

Revisiting the past—mammals

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

$M \leq 10$ kg:

	N	$\hat{\alpha}$	$p_{2/3}$	$p_{3/4}$
Kleiber	5	0.667	0.99	0.088
Brody	26	0.709	$< 10^{-3}$	$< 10^{-3}$
Heusner	357	0.668	0.91	$< 10^{-15}$

$M \geq 10$ kg:

	N	$\hat{\alpha}$	$p_{2/3}$	$p_{3/4}$
Kleiber	8	0.754	$< 10^{-4}$	0.66
Brody	9	0.760	$< 10^{-3}$	0.56
Heusner	34	0.877	$< 10^{-12}$	$< 10^{-7}$

Fluctuations—Kolmogorov-Smirnov test

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 69/117

[07–Nov–1999 peter dodds]

20 bins

$P(\log_{10} B/M^{2/3})$

$\log_{10} B/M^{2/3}$

$$P(B\,|\,M) = 1/M^{2/3} f(B/M^{2/3})$$

Analysis of residuals

1. Presume an exponent of your choice: 2/3 or 3/4.
2. Fit the prefactor ($\log_{10} c$) and then examine the residuals:

$$r_i = \log_{10} B_i - (\alpha' \log_{10} M_i - \log_{10} c).$$

3. H_0: residuals are uncorrelated
   H_1: residuals are correlated.
4. Measure the correlations in the residuals and compute a *p*-value.

Analysis of residuals

We use the spiffing Spearman Rank-Order Correlation Cofficient.

Basic idea:

- ▶ Given $\{(x_i, y_i)\}$, rank the $\{x_i\}$ and $\{y_i\}$ separately from smallest to largest. Call these ranks R_i and S_i.
- ▶ Now calculate correlation coefficient for ranks, r_s:
- ▶
$$r_s = \frac{\sum_{i=1}^{n}(R_i - \bar{R})(S_i - \bar{S})}{\sqrt{\sum_{i=1}^{n}(R_i - \bar{R})^2}\sqrt{\sum_{i=1}^{n}(S_i - \bar{S})^2}}$$
- ▶ Perfect correlation: x_i's and y_i's both increase monotonically.

Analysis of residuals

We assume all rank orderings are equally likely:

- r_s is distributed according to a Student's distribution with $N - 2$ degrees of freedom.
- Excellent feature: Non-parametric—real distribution of x's and y's doesn't matter.
- Bonus: works for non-linear monotonic relationships as well.
- See "Numerical Recipes in C/Fortran" which contains many good things. [20]

Analysis of residuals—mammals

(a) $M < 3.2$ kg, (b) $M < 10$ kg, (c) $M < 32$ kg, (d) all mammals.

Analysis of residuals—birds

(a) $M < .1$ kg, (b) $M < 1$ kg, (c) $M < 10$ kg, (d) all birds.

Basic basin quantities: a, l, $L_\parallel$, $L_\perp$:

- a = drainage basin area
- ℓ = length of longest (main) stream
- $L = L_\parallel$ = longitudinal length of basin

River networks

- 1957: J. T. Hack[11]
  "Studies of Longitudinal Stream Profiles in Virginia and Maryland"

$$\ell \sim a^h$$

$$h \sim 0.6$$

- Anomalous scaling: we would expect $h = 1/2$...
- Subsequent studies: $0.5 \lesssim h \lesssim 0.6$
- Another quest to find universality/god...
- A catch: studies done on small scales.

Large-scale networks

(1992) Montgomery and Dietrich [19]:

- ▶ Composite data set: includes everything from unchanneled valleys up to world's largest rivers.
- ▶ Estimated fit:

$$L \simeq 1.78 a^{0.49}$$

- ▶ Mixture of basin and main stream lengths.

World's largest rivers only:

- Data from Leopold (1994) [16, 8]
- Estimate of Hack exponent: $h = 0.50 \pm 0.06$

Earlier theories

Building on the surface area idea...

- Blum (1977) [5] speculates on four-dimensional biology:

$$P \propto M^{(d-1)/d}$$

- $d = 3$ gives $\alpha = 2/3$
- $d = 4$ gives $\alpha = 3/4$
- So we need another dimension...
- Obviously, a bit silly.

Earlier theories

Building on the surface area idea:

► McMahon (70's, 80's): Elastic Similarity [17, 18]

► Idea is that organismal shapes scale allometrically with 1/4 powers (like nails and trees...)

► Appears to be true for ungulate legs.

► Metabolism and shape never properly connected.

Nutrient delivering networks:

- 1960's: Rashevsky considers blood networks and finds a 2/3 scaling.
- 1997: West *et al.* [25] use a network story to find 3/4 scaling.

A Mammal

B Plant

C Model

$k = 0 \quad 1 \quad 2 \quad 3 \quad 4 \; ..... \; N$

D Parameters

Δp_k

l_k

$\bar{u}_k \longrightarrow$

r_k

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Nutrient delivering networks:

West et al.'s assumptions:

- hierarchical network
- capillaries (delivery units) invariant
- network impedance is minimized via evolution

Claims:

- $P \propto M^{3/4}$
- networks are fractal
- quarter powers everywhere

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Impedance measures:

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Poiseuille flow (outer branches):

$$Z = \frac{8\mu}{\pi} \sum_{k=0}^{N} \frac{\ell_k}{r_k^4 N_k}$$

Pulsatile flow (main branches):

$$Z \propto \sum_{k=0}^{N} \frac{h_k^{1/2}}{r_k^{5/2} N_k}$$

Not so fast . . .

Actually, model shows:

- $P \propto M^{3/4}$ does not follow for pulsatile flow
- networks are not necessarily fractal.

Do find:

- Murray's cube law (1927) for outer branches:

$$r_0^3 = r_1^3 + r_2^3$$

- Impedance is distributed evenly.
- Can still assume networks are fractal.

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Connecting network structure to α

1. Ratios of network parameters:

$$R_n = \frac{n_{k+1}}{n_k},\ R_\ell = \frac{\ell_{k+1}}{\ell_k},\ R_r = \frac{r_{k+1}}{r_k}$$

2. Number of capillaries $\propto P \propto M^\alpha$.

$$\Rightarrow \boxed{\alpha = -\frac{\ln R_n}{\ln R_r^2 R_\ell}}$$

(also problematic due to prefactor issues)

Soldiering on, assert:

- area-preservingness: $R_r = R_n^{-1/2}$
- space-fillingness: $R_\ell = R_n^{-1/3}$
-

$$\Rightarrow \alpha = 3/4$$

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 87/117

Data from real networks

Network	R_n	R_r^{-1}	R_ℓ^{-1}	$-\frac{\ln R_r}{\ln R_n}$	$-\frac{\ln R_\ell}{\ln R_n}$	α
West *et al.*	–	–	–	1/2	1/3	3/4
rat (PAT)	2.76	1.58	1.60	0.45	0.46	0.73
cat (PAT) (Turcotte *et al.*[24])	3.67	1.71	1.78	0.41	0.44	0.79
dog (PAT)	3.69	1.67	1.52	0.39	0.32	0.90
pig (LCX)	3.57	1.89	2.20	0.50	0.62	0.62
pig (RCA)	3.50	1.81	2.12	0.47	0.60	0.65
pig (LAD)	3.51	1.84	2.02	0.49	0.56	0.65
human (PAT)	3.03	1.60	1.49	0.42	0.36	0.83
human (PAT)	3.36	1.56	1.49	0.37	0.33	0.94

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Simple supply networks

- ▶ Banavar et al., Nature, (1999)[1]
- ▶ Flow rate argument
- ▶ Ignore impedance
- ▶ Very general attempt to find most efficient transportation networks

Simple supply networks

▶ Banavar *et al.* find 'most efficient' networks with

$$P \propto M^{d/(d+1)}$$

▶ ... but also find

$$V_{\text{network}} \propto M^{(d+1)/d}$$

▶ $d = 3$:

$$V_{\text{blood}} \propto M^{4/3}$$

▶ Consider a 3 g shrew with $V_{\text{blood}} = 0.1\, V_{\text{body}}$

▶ $\Rightarrow$ 3000 kg elephant with $V_{\text{blood}} = 10\, V_{\text{body}}$

▶ Such a pachyderm would be rather miserable.

Geometric argument

- ▶ Consider one source supplying many sinks in a *d*-dim. volume in a *D*-dim. ambient space.
- ▶ Assume sinks are invariant.
- ▶ Assume $\rho = \rho(V)$.
- ▶ Assume some cap on flow speed of material.
- ▶ See network as a bundle of virtual vessels:

- ▶ Q: how does the number of sustainable sinks N_{sinks} scale with volume *V* for the most efficient network design?
- ▶ Or: what is the highest α for $N_{\text{sinks}} \propto V^\alpha$?

Geometric argument

- Allometrically growing regions:

- Have d length scales which scale as

$$L_i \propto V^{\gamma_i} \text{ where } \gamma_1 + \gamma_2 + \ldots + \gamma_d = 1.$$

- For isometric growth, $\gamma_i = 1/d$.
- For allometric growth, we must have at least two of the $\{\gamma_i\}$ being different

Geometric argument

▶ Best and worst configurations (Banavar et al.)

▶ Rather obviously:
min $V_{\mathrm{net}} \propto \sum$ distances from source to sinks.

Minimal network volume:

Real supply networks are close to optimal:

Figure 1. (a) Commuter rail network in the Boston area. The arrow marks the assumed root of the network. (b) Star graph. (c) Minimum spanning tree. (d) The model of equation (3) applied to the same set of stations.

(2006) Gastner and Newman [10]: "Shape and efficiency in spatial distribution networks"

Minimal network volume:

Approximate network volume by integral over region:

$$\min V_{\text{net}} \propto \int_{\Omega_{d,D}(V)} \rho \, ||\vec{x}|| \, d\vec{x}$$

$$\rightarrow \rho V^{1+\gamma_{\max}} \int_{\Omega_{d,D}(c)} (c_1^2 u_1^2 + \ldots + c_k^2 u_k^2)^{1/2} d\vec{u}$$

$$\propto \rho V^{1+\gamma_{\max}}$$

Geometric argument

- General result:

$$\min V_{\text{net}} \propto \rho V^{1+\gamma_{\max}}$$

- If scaling is isometric, we have $\gamma_{\max} = 1/d$:

$$\min V_{\text{net/iso}} \propto \rho V^{1+1/d} = \rho V^{(d+1)/d}$$

- If scaling is allometric, we have $\gamma_{\max} = \gamma_{\text{allo}} > 1/d$: and

$$\min V_{\text{net/allo}} \propto \rho V^{1+\gamma_{\text{allo}}}$$

- Isometrically growing volumes require less network volume than allometrically growing volumes:

$$\frac{\min V_{\text{net/iso}}}{\min V_{\text{net/allo}}} \to 0 \text{ as } V \to \infty$$

Blood networks

- ▶ Material costly $\Rightarrow$ expect lower optimal bound of $V_{\mathrm{net}} \propto \rho V^{(d+1)/d}$ to be followed closely.

- ▶ For cardiovascular networks, $d = D = 3$.

- ▶ Blood volume scales linearly with body volume [23], $V_{\mathrm{net}} \propto V$.

- ▶ Sink density must $\therefore$ decrease as volume increases:

$$\rho \propto V^{-1/d}.$$

- ▶ Density of suppliable sinks decreases with organism size.

Blood networks

▶ Then P, the rate of overall energy use in Ω, can at most scale with volume as

$$P \propto \rho V \propto \rho M \propto M^{(d-1)/d}$$

▶ For $d = 3$ dimensional organisms, we have

$$\boxed{P \propto M^{2/3}}$$

Recap:

- ▶ The exponent $\alpha = 2/3$ works for all birds and mammals up to 10–30 kg
- ▶ For mammals > 10–30 kg, maybe we have a new scaling regime
- ▶ Economos: limb length break in scaling around 20 kg
- ▶ White and Seymour, 2005: unhappy with large herbivore measurements. Find $\alpha \simeq 0.686 \pm 0.014$

Prefactor:

Stefan-Boltzmann law:

►
$$\frac{\mathrm{d}E}{\mathrm{d}t} = \sigma S T^4$$

where S is surface and T is temperature.

► Very rough estimate of prefactor based on scaling of normal mammalian body temperature and surface area S:

$$B \simeq 10^5 M^{2/3} \text{erg/sec.}$$

► Measured for $M \leq 10$ kg:

$$B = 2.57 \times 10^5 M^{2/3} \text{erg/sec.}$$

Scaling

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 102/117

River networks

- View river networks as collection networks.
- Many sources and one sink.
- Assume ρ is constant over time:

$$V_{\text{net}} \propto \rho V^{(d+1)/d} = \text{constant} \times V^{3/2}$$

- Network volume grows faster than basin 'volume' (really area).
- It's all okay:
  Landscapes are d=2 surfaces living in D=3 dimension.
- Streams can grow not just in width but in depth...

Hack's law

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

- ▶ Volume of water in river network can be calculated by adding up basin areas
- ▶ Flows sum in such a way that

$$V_{\text{net}} = \sum_{\text{all pixels}} a_{\text{pixel } i}$$

- ▶ Hack's law again:

$$\ell \sim a^h$$

- ▶ Can argue

$$V_{\text{net}} \propto V_{\text{basin}}^{1+h} = a_{\text{basin}}^{1+h}$$

  where h is Hack's exponent.

- ▶ ∴ minimal volume calculations gives

$$\boxed{h = 1/2}$$

Real data:

- Banavar et al.'s approach[1] is okay because ρ really is constant.
- The irony: shows optimal basins are isometric
- Optimal Hack's law: $\ell \sim a^h$ with $h = 1/2$
- (Zzzzz)

From Banavar et al. (1999)[1]

Even better—prefactors match up:

Conclusion

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

▶ Supply network story consistent with dimensional analysis.

▶ Isometrically growing regions can be more efficiently supplied than allometrically growing ones.

▶ Ambient and region dimensions matter ($D = d$ versus $D > d$).

▶ Deviations from optimal scaling suggest inefficiency (e.g., gravity for organisms, geological boundaries).

▶ Actual details of branching networks not that important.

▶ Exact nature of self-similarity varies.

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

References I

📄 J. R. Banavar, A. Maritan, and A. Rinaldo.
Size and form in efficient transportation networks.
Nature, 399:130–132, 1999. pdf (⊞)

📄 P. Bennett and P. Harvey.
Active and resting metabolism in birds—allometry, phylogeny and ecology.
J. Zool., 213:327–363, 1987.

📄 L. M. A. Bettencourt, J. Lobo, D. Helbing, Kühnhert, and G. B. West.
Growth, innovation, scaling, and the pace of life in cities.
Proc. Natl. Acad. Sci., 104(17):7301–7306, 2007. pdf (⊞)

References II

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 111/117

📄 K. L. Blaxter, editor.
Energy Metabolism; Proceedings of the 3rd symposium held at Troon, Scotland, May 1964.
Academic Press, New York, 1965.

📄 J. J. Blum.
On the geometry of four-dimensions and the relationship between metabolism and body mass.
J. Theor. Biol., 64:599–601, 1977.

📄 S. Brody.
Bioenergetics and Growth.
Reinhold, New York, 1945.
reprint, .

📄 M. H. DeGroot.
Probability and Statistics.
Addison-Wesley, Reading, Massachusetts, 1975.

References III

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

Frame 112/117

📄 P. S. Dodds and D. H. Rothman.
Scaling, universality, and geomorphology.
Annu. Rev. Earth Planet. Sci., 28:571–610, 2000.
pdf (⊞)

📄 P. S. Dodds, D. H. Rothman, and J. S. Weitz.
Re-examination of the "3/4-law" of metabolism.
Journal of Theoretical Biology, 209(1):9–27, March 2001.
. pdf (⊞)

📄 M. T. Gastner and M. E. J. Newman.
Shape and efficiency in spatial distribution networks.
J. Stat. Mech.: Theor. & Exp., 1:01015–, 2006.
pdf (⊞)

References IV

Scaling

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

📄 J. T. Hack.
Studies of longitudinal stream profiles in Virginia and Maryland.
United States Geological Survey Professional Paper, 294-B:45–97, 1957.

📄 A. Hemmingsen.
The relation of standard (basal) energy metabolism to total fresh weight of living organisms.
Rep. Steno Mem. Hosp., 4:1–58, 1950.

📄 A. Hemmingsen.
Energy metabolism as related to body size and respiratory surfaces, and its evolution.
Rep. Steno Mem. Hosp., 9:1–110, 1960.

References V

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

📄 A. A. Heusner.
Size and power in mammals.
Journal of Experimental Biology, 160:25–54, 1991.

📄 M. Kleiber.
Body size and metabolism.
Hilgardia, 6:315–353, 1932.

📄 L. B. Leopold.
A View of the River.
Harvard University Press, Cambridge, MA, 1994.

📄 T. McMahon.
Size and shape in biology.
Science, 179:1201–1204, 1973. pdf (⊞)

📄 T. A. McMahon and J. T. Bonner.
On Size and Life.
Scientific American Library, New York, 1983.

References VI

Scaling-at-large

Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

📄 D. R. Montgomery and W. E. Dietrich.
Channel initiation and the problem of landscape scale.
Science, 255:826–30, 1992. pdf (⊞)

📄 W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery.
Numerical Recipes in C.
Cambridge University Press, second edition, 1992.

📄 M. Rubner.
Ueber den einfluss der körpergrösse auf stoffund kraftwechsel.
Z. Biol., 19:535–562, 1883.

References VII

Scaling-at-large
Allometry
Definitions
Examples
History: Metabolism
Measuring exponents
History: River networks
Earlier theories
Geometric argument
Blood networks
River networks
Conclusion

References

📄 Sarrus and Rameaux.
Rapport sur une mémoire adressé à l'Académie de Médecine.
Bull. Acad. R. Méd. (Paris), 3:1094–1100, 1838–39.

📄 W. R. Stahl.
Scaling of respiratory variables in mammals.
Journal of Applied Physiology, 22:453–460, 1967.

📄 D. L. Turcotte, J. D. Pelletier, and W. I. Newman.
Networks with side branching in biology.
Journal of Theoretical Biology, 193:577–592, 1998.

📄 G. B. West, J. H. Brown, and B. J. Enquist.
A general model for the origin of allometric scaling laws in biology.
Science, 276:122–126, 1997. pdf (⊞)

References VIII

📄 K. Zhang and T. J. Sejnowski.
A universal scaling law between gray matter and
white matter of cerebral cortex.
Proceedings of the National Academy of Sciences,
97:5621–5626, May 2000. pdf (⊞)