

**Principles of Complex Systems, CSYS/MATH 300**  
**University of Vermont, Fall 2011**  
**Assignment 4**

**Dispersed:** Thursday, October 7, 2011.

**Due:** By start of lecture, 11:30 am, Thursday, October 14, 2011.

*Some useful reminders:*

**Instructor:** Peter Dodds

**Office:** Farrell Hall, second floor, Trinity Campus

**E-mail:** peter.dodds@uvm.edu

**Office hours:** 12:50 pm to 3:50 pm, Wednesday

**Course website:** <http://www.uvm.edu/~pdodds/teaching/courses/2011-08UVM-300>

---

All parts are worth 3 points unless marked otherwise. Please show all your working clearly and list the names of others with whom you collaborated.

Graduate students are requested to use  $\LaTeX$  (or related  $\TeX$  variant).

---

1. (3+3 points) *Simon's model I:*

For Herbert Simon's model of what we've called Random Competitive Replication, we found in class that the normalized number of groups in the long time limit,  $n_k$ , satisfies the following difference equation:

$$\frac{n_k}{n_{k-1}} = \frac{(k-1)(1-\rho)}{1+(1-\rho)k} \quad (1)$$

where  $k \geq 2$ . The model parameter  $\rho$  is the probability that a newly arriving node forms a group of its own (or is a novel word, starts a new city, has a unique flavor, etc.). For  $k = 1$ , we have instead

$$n_1 = \rho - (1-\rho)n_1 \quad (2)$$

which directly gives us  $n_1$  in terms of  $\rho$ .

- (a) Derive the exact solution for  $n_k$  in terms of gamma functions and ultimately the beta function.
- (b) From this exact form, determine the large  $k$  behavior for  $n_k$  ( $\sim k^{-\gamma}$ ) and identify the exponent  $\gamma$  in terms of  $\rho$ .

2. (3+3 points) *Simon's model II:*

- (a) A missing piece from the lectures: Obtain  $\gamma$  in terms of  $\rho$  by expanding Eq. 1 in terms of  $1/k$ . In the end, you will need to express  $n_k/n_{k-1}$  as  $(1 - 1/k)^\theta$ ; from here, you will be able to identify  $\gamma$ . Taylor expansions and Procrustean truncations will be in order.

This (dirty) method avoids finding the exact form for  $n_k$ .

- (b) What happens to  $\gamma$  in the limits  $\rho \rightarrow 0$  and  $\rho \rightarrow 1$ ? Explain in a sentence or two what's going on in these cases and how the specific limiting value of  $\gamma$  makes sense.

3. (6 + 3 + 3 points)

In Simon's original model, the expected total number of distinct groups at time  $t$  is  $\rho t$ . Recall that each group is made up of elements of a particular flavor.

In class, we derived the fraction of groups containing only 1 element, finding

$$n_1^{(g)} = \frac{N_1(t)}{\rho t} = \frac{1}{2 - \rho}$$

- (a) (3 + 3 points)

Find the form of  $n_2^{(g)}$  and  $n_3^{(g)}$ , the fraction of groups that are of size 2 and size 3.

- (b) Using data for James Joyce's Ulysses (see below), first show that Simon's estimate for the innovation rate  $\rho_{\text{est}} \simeq 0.115$  is reasonably accurate for the version of the text's word counts given below. You should find a slightly higher number.

- (c) Now compare the theoretical estimates for  $n_1^{(g)}$ ,  $n_2^{(g)}$ , and  $n_3^{(g)}$ , with empirical values you obtain for Ulysses.

The data (links are clickable):

- Matlab file (sortedcounts = word frequency  $f$  in descending order, sortedwords = ranked words):  
<http://www.uvm.edu/~pdodds/teaching/courses/2011-08UVM-300/docs/ulysses.mat>
- Colon-separated text file (first column = word, second column = word frequency  $f$ ): <http://www.uvm.edu/~pdodds/teaching/courses/2011-08UVM-300/docs/ulysses.txt>

Data taken from <http://www.doc.ic.ac.uk/~rac101/concord/texts/ulysses/>. Note that some matching words with differing capitalization are recorded as separate words.

## References

- [1] M. Abramowitz and I. A. Stegun, editors. *Handbook of Mathematical Functions*. Dover Publications, New York, 1974.
- [2] I. Gradshteyn and I. Ryzhik. *Table of Integrals, Series, and Products*. Academic Press, San Diego, fifth edition, 1994.