Power Law Size Distributions

Principles of Complex Systems CSYS/MATH 300, Fall, 2011

Prof. Peter Dodds

Department of Mathematics & Statistics Center for Complex Systems Vermont Advanced Computing Center University of Vermont

















Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

Poforonoos







Outline

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

References

Power Law Size Distributions

efinition

Examples

Wild vs. Mild

CCDFs

Zipf's law $Zipf \Leftrightarrow CCDF$





Examples
Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF

$$P(\text{size} = x) \sim c x^{-\gamma}$$

where
$$0 < x_{\min} < x < x_{\max}$$

and $\gamma > 1$

- Exciting class exercise: sketch this function.
- $\triangleright x_{\min} = \text{lower cutoff}$
- $> x_{\text{max}} = \text{upper cutoff}$
- ► Negative linear relationship in log-log space:

$$\log_{10} P(x) = \log_{10} c - \gamma \log_{10} x$$

▶ We use base 10 because we are good people.







20 3 of 36

Examples
Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF

$$P(\text{size} = x) \sim c x^{-\gamma}$$

where
$$0 < x_{min} < x < x_{max}$$

and $\gamma > 1$

- Exciting class exercise: sketch this function.
- $\rightarrow x_{\min} = lower cutoff$
- $\triangleright x_{\max} = \text{upper cutoff}$
- ► Negative linear relationship in log-log space:

$$\log_{10} P(x) = \log_{10} c - \gamma \log_{10} x$$

▶ We use base 10 because we are good people.





Examples
Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF

The sizes of many systems' elements appear to obey an inverse power-law size distribution:

$$P(\mathsf{size} = x) \sim c \, x^{-\gamma}$$
 where $0 < x_{\min} < x < x_{\max}$ and $\gamma > 1$

- Exciting class exercise: sketch this function.
- $\rightarrow x_{\min} = lower cutoff$
- $\rightarrow x_{\text{max}} = \text{upper cutoff}$
- ▶ Negative linear relationship in log-log space:

$$\log_{10} P(x) = \log_{10} c - \gamma \log_{10} x$$

▶ We use base 10 because we are good people.





20 3 of 36

Examples
Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF

The sizes of many systems' elements appear to obey an inverse power-law size distribution:

$$P(\text{size} = x) \sim c \, x^{-\gamma}$$
 where $0 < x_{\min} < x < x_{\max}$ and $\gamma > 1$

- Exciting class exercise: sketch this function.
- $\rightarrow x_{\min} = lower cutoff$
- $\rightarrow x_{\text{max}} = \text{upper cutoff}$
- Negative linear relationship in log-log space:

$$\log_{10} P(x) = \log_{10} c - \frac{\gamma}{\gamma} \log_{10} x$$

▶ We use base 10 because we are good people.





20 € 3 of 36

Examples
Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF

The sizes of many systems' elements appear to obey an inverse power-law size distribution:

$$P(\text{size} = x) \sim c x^{-\gamma}$$

where
$$0 < x_{\min} < x < x_{\max}$$
 and $\gamma > 1$

- Exciting class exercise: sketch this function.
- $\rightarrow x_{\min} = lower cutoff$
- $\rightarrow x_{\text{max}} = \text{upper cutoff}$
- Negative linear relationship in log-log space:

$$\log_{10} P(x) = \log_{10} c - \frac{\gamma}{\gamma} \log_{10} x$$

▶ We use base 10 because we are good people.







$$P(x) \sim c x^{-\gamma}$$
 for x large.

- Still use term 'power law distribution.'
- ▶ Other terms:
 - Fat-tailed distributions.
 - Heavy-tailed distributions.

Beware:

Inverse power laws aren't the only ones: lognormals (⊞), Weibull distributions (⊞), ...

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







$$P(x) \sim c x^{-\gamma}$$
 for x large.

- Still use term 'power law distribution.'
- ▶ Other terms:
 - Fat-tailed distributions.
 - Heavy-tailed distributions.

Beware:

Inverse power laws aren't the only ones: lognormals (⊞), Weibull distributions (⊞), . . .

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







$$P(x) \sim c x^{-\gamma}$$
 for x large.

- Still use term 'power law distribution.'
- Other terms:
 - Fat-tailed distributions.
 - Heavy-tailed distributions.

Beware

Inverse power laws aren't the only ones: lognormals (⊞), Weibull distributions (⊞), . . .

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $Zipf \Leftrightarrow CCDF$







$$P(x) \sim c x^{-\gamma}$$
 for x large.

- Still use term 'power law distribution.'
- Other terms:
 - Fat-tailed distributions.
 - Heavy-tailed distributions.

Beware:

Inverse power laws aren't the only ones: lognormals (⊞), Weibull distributions (⊞), ...

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References

Many systems have discrete sizes *k*:

- Word frequency
- ▶ Node degree in networks: # friends, # hyperlinks, etc.
- ▶ # citations for articles, court decisions, etc.

$$P(k) \sim c \, k^{-\gamma}$$
 where $k_{\min} \leq k \leq k_{\max}$

- ▶ Obvious fail for k = 0.
- ► Again, typically a description of distribution's tail.







Zipf's law

Zipf ⇔ CCDF

References

Many systems have discrete sizes *k*:

- Word frequency
- ▶ Node degree in networks: # friends, # hyperlinks, etc.
- ▶ # citations for articles, court decisions, etc.

$$P(k) \sim c \, k^{-\gamma}$$
 where $k_{\min} \leq k \leq k_{\max}$

- ▶ Obvious fail for k = 0.
- Again, typically a description of distribution's tail.







Brown Corpus (\boxplus) ($\sim 10^6$ words):

		_	
rank	word	% q	
1.	the	6.8872	
2.	of	3.5839	
3.	and	2.8401	
4.	to	2.5744	
5.	а	2.2996	
6.	in	2.1010	
7.	that	1.0428	
8.	is	0.9943	
9.	was	0.9661	
10.	he	0.9392	
11.	for	0.9340	
12.	it	0.8623	
13.	with	0.7176	
14.	as	0.7137	
15.	his	0.6886	

 	٥).	
rank	word	% q
1945.	apply	0.0055
1946.	vital	0.0055
1947.	September	0.0055
1948.	review	0.0055
1949.	wage	0.0055
1950.	motor	0.0055
1951.	fifteen	0.0055
1952.	regarded	0.0055
1953.	draw	0.0055
1954.	wheel	0.0055
1955.	organized	0.0055
1956.	vision	0.0055
1957.	wild	0.0055
1958.	Palmer	0.0055
1959.	intensity	0.0055

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







Power Law Size Distributions

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF









First—a Gaussian example:

$$P(x)dx = \frac{1}{\sqrt{2\pi}\sigma}e^{-(x-\mu)^2/2\sigma}dx$$






mean $\mu = 10$, variance $\sigma^2 = 1$.

The statistics of surprise—words:

Raw 'probability' (binned):



Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law


 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





The statistics of surprise—words:

Raw 'probability' (binned):



Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law


Zipf ⇔ CCDF






The statistics of surprise—words:

'Exceedance probability':





Power Law Size Distributions

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







Wild vs. Mild

CCDFs Zipf's law

Zipf ⇔ CCDF

How many words Test do you know? Vour Voca

- Test capitalizes on word frequency following a heavily skewed frequency distribution with a decaying power law tail.
- ► Let's do it collectively... (⊞)



Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF

Poforonooo

 $\mathbf{v}^{-\gamma}$


Test How many words do you know?

Your Vocab

- Test capitalizes on word frequency following a heavily skewed frequency distribution with a decaying power law tail.
- ▶ Let's do it collectively... (⊞)



Gutenberg-Richter law (⊞)



- Log-log plot
- ▶ Base 10
- ▶ Slope = -1

 $N(M > m) \propto m^{-1}$

 From both the very awkwardly similar Christensen et al. and Bak et al.:
 "Unified scaling law for earthquakes" [3, 1] Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

 $\mathbf{x}^{-\gamma}$





What is perhaps most surprising about the Japan earthquake is how misleading history can be. In the past 300 years, no earthquake nearly that large—nothing larger than magnitude eight—had struck in the Japan subduction zone. That, in turn, led to assumptions about how large a tsunami might strike the coast.

"It did them a giant disservice," said Dr. Stein of the geological survey. That is not the first time that the earthquake potential of a fault has been underestimated. Most geophysicists did not think the Sumatra fault could generate a magnitude 9.1 earthquake, . . .

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







What is perhaps most surprising about the Japan earthquake is how misleading history can be. In the past 300 years, no earthquake nearly that large—nothing larger than magnitude eight—had struck in the Japan subduction zone. That, in turn, led to assumptions about how large a tsunami might strike the coast.

"It did them a giant disservice," said Dr. Stein of the geological survey. That is not the first time that the earthquake potential of a fault has been underestimated. Most geophysicists did not think the Sumatra fault could generate a magnitude 9.1 earthquake, . . .

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







Two things we have poor cognitive understanding of:

- Probability
 - ► Ex. The Monty Hall Problem (⊞)
 - ► Ex. Son born on Tuesday (⊞).
- 2. Logarithmic scales.

On counting and logarithms:



- Listen to Radiolab's "Numbers." (⊞).
- ▶ Later: Benford's Law (⊞).

Definition

Examples

Wild vs. Mild


CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







The distributions of twelve quantities reputed to follow power laws. (l) Populations of US cities in the vear occurrence of family Magnitude

Power Law Size Distributions

Examples

Wild vs. Mild Zipf's law

Zipf ⇔ CCDF







Zipf's law

Zipf ⇔ CCDF

Examples:

- Earthquake magnitude (Gutenberg-Richter) law (\boxplus)): [1] $P(M) \propto M^{-2}$
- ▶ Number of war deaths: [9] $P(d) \propto d^{-1.8}$
- Sizes of forest fires [4]
- ► Sizes of cities: [10] $P(n) \propto n^{-2.1}$
- Number of links to and from websites [2]
- See in part Simon [10] and M.E.J. Newman [6] "Power
- Note: Exponents range in error





Examples:

- Earthquake magnitude (Gutenberg-Richter) law (\boxplus)):^[1] $P(M) \propto M^{-2}$
- ▶ Number of war deaths: [9] $P(d) \propto d^{-1.8}$
- Sizes of forest fires [4]
- ► Sizes of cities: [10] $P(n) \propto n^{-2.1}$
- Number of links to and from websites [2]
- See in part Simon [10] and M.E.J. Newman [6] "Power laws, Pareto distributions and Zipf's law" for more.
- Note: Exponents range in error

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF





Examples:

- Number of citations to papers: [7, 8] $P(k) \propto k^{-3}$.
- ▶ Individual wealth (maybe): $P(W) \propto W^{-2}$.
- ▶ Distributions of tree trunk diameters: $P(d) \propto d^{-2}$.
- The gravitational force at a random point in the universe: [5] $P(F) \propto F^{-5/2}$.
- ▶ Diameter of moon craters: [6] $P(d) \propto d^{-3}$.
- ▶ Word frequency: [10] e.g., $P(k) \propto k^{-2.2}$ (variable)





Examples
Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

Gaussians versus power-law distributions:

- ► Mediocristan versus Extremistan
- Mild versus Wild (Mandelbrot)
- Example: Height versus wealth.

BLACK SWAN



 See "The Black Swan" by Nassim Taleb. [11]


 $\mathbf{x}^{-\gamma}$

Nassim Nicholas Taleb



Turkeys...

FIGURE 1: ONE THOUSAND AND ONE DAYS OF HISTORY



A turkey before and after Thanksgiving. The history of a process over a thousand days tells you nothing about what is to happen next. This naïve projection of the future from the past can be applied to anything.

From "The Black Swan" [11]

Power Law Size Distributions

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References





2 0 0 18 of 36

Zipf ⇔ CCDF

References

$\mathbf{x}^{-\gamma}$

Mediocristan/Extremistan

- Most typical member is mediocre/Most typical is either giant or tiny
- Winners get a small segment/Winner take almost all effects
- When you observe for a while, you know what's going on/It takes a very long time to figure out what's going or
- ► Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- Tyranny of the collective/Tyranny of the rare and accidental



Mediocristan/Extremistan

- Most typical member is mediocre/Most typical is either giant or tiny
- ▶ When you observe for a while, you know what's going
- Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- Tyranny of the collective/Tyranny of the rare and

Examples

Wild vs. Mild CCDFs

Zipf's law

Zipf ⇔ CCDF







Zipf ⇔ CCDF

References

Mediocristan/Extremistan

- Most typical member is mediocre/Most typical is either giant or tiny
- Winners get a small segment/Winner take almost all effects
- When you observe for a while, you know what's going on/It takes a very long time to figure out what's going or
- Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- Tyranny of the collective/Tyranny of the rare and accidental







Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References

$\mathbf{x}^{-\gamma}$

Mediocristan/Extremistan

- Most typical member is mediocre/Most typical is either giant or tiny
- Winners get a small segment/Winner take almost all effects
- When you observe for a while, you know what's going on/It takes a very long time to figure out what's going on
- ► Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- Tyranny of the collective/Tyranny of the rare and accidental





- Most typical member is mediocre/Most typical is either giant or tiny
- Winners get a small segment/Winner take almost all effects
- When you observe for a while, you know what's going on/It takes a very long time to figure out what's going on
- Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- ► Tyranny of the collective/Tyranny of the rare and accidental

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF







Zipf's law

Zipf ⇔ CCDF

References

Mediocristan/Extremistan

- Most typical member is mediocre/Most typical is either giant or tiny
- Winners get a small segment/Winner take almost all effects
- When you observe for a while, you know what's going on/It takes a very long time to figure out what's going on
- Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- Tyranny of the collective/Tyranny of the rare and accidental







Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References

Mediocristan/Extremistan

- Most typical member is mediocre/Most typical is either giant or tiny
- Winners get a small segment/Winner take almost all effects
- When you observe for a while, you know what's going on/It takes a very long time to figure out what's going on
- Prediction is easy/Prediction is hard
- History crawls/History makes jumps
- Tyranny of the collective/Tyranny of the rare and accidental





Size distributions





Power law size distributions are sometimes called Pareto distributions (⊞) after Italian scholar Vilfredo Pareto. (⊞)

Wild vs. Mild CCDFs Zipf's law

Examples

Zipf ⇔ CCDF
References

Pareto noted wealth in Italy was distributed unevenly (80–20 rule; misleading).
 Term used especially by







Size distributions





Power law size distributions are sometimes called Pareto distributions (⊞) after Italian scholar Vilfredo Pareto. (⊞)

- Pareto noted wealth in Italy was distributed unevenly (80–20 rule; misleading).
- ► Term used especially by practitioners of the Disma Science (⊞).

Definition

Examples

Wild vs. Mild CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





Size distributions





Power law size distributions are sometimes called Pareto distributions (⊞) after Italian scholar Vilfredo Pareto. (⊞)

- Pareto noted wealth in Italy was distributed unevenly (80-20 rule; misleading).
- Term used especially by practitioners of the Dismal Science (\boxplus) .

Examples

Wild vs. Mild CCDFs

Zipf's law

Zipf ⇔ CCDF







Exhibit A:

• Given $P(x) = cx^{-\gamma}$ with $0 < x_{min} < x < x_{max}$, the mean is $(\gamma \neq 2)$:

$$\langle \mathbf{x} \rangle = \frac{\mathbf{c}}{\mathbf{2} - \gamma} \left(\mathbf{x}_{\text{max}}^{2 - \gamma} - \mathbf{x}_{\text{min}}^{2 - \gamma} \right).$$

- ▶ Mean 'blows up' with upper cutoff if γ < 2.
- ▶ Mean depends on lower cutoff if $\gamma > 2$.
- $ightharpoonup \gamma < 2$: Typical sample is large.
- $\sim \gamma > 2$: Typical sample is small.

Insert question from assignment 1 (⊞)

Examples

Wild vs. Mild

CCDFs Zipf's law

Zipf ⇔ CCDF







Wild vs. Mild

CCDFs

Zipf's law
Zipf ⇔ CCDF

Exhibit A:

• Given $P(x) = cx^{-\gamma}$ with $0 < x_{min} < x < x_{max}$, the mean is $(\gamma \neq 2)$:

$$\langle {\it x}
angle = rac{c}{2-\gamma} \left({\it x}_{
m max}^{2-\gamma} - {\it x}_{
m min}^{2-\gamma}
ight).$$

- ▶ Mean 'blows up' with upper cutoff if γ < 2.
- ▶ Mean depends on lower cutoff if γ > 2.
- $ightharpoonup \gamma <$ 2: Typical sample is large.
- $ightharpoonup \gamma > 2$: Typical sample is small.

Insert question from assignment 1 (⊞)





Examples

CCDFs

Zipf's law Zipf ⇔ CCDF

Wild vs. Mild

Exhibit A:

• Given $P(x) = cx^{-\gamma}$ with $0 < x_{min} < x < x_{max}$, the mean is $(\gamma \neq 2)$:

$$\langle \mathbf{x} \rangle = \frac{\mathbf{c}}{\mathbf{2} - \gamma} \left(\mathbf{x}_{\text{max}}^{2 - \gamma} - \mathbf{x}_{\text{min}}^{2 - \gamma} \right).$$

- ▶ Mean 'blows up' with upper cutoff if γ < 2.
- ▶ Mean depends on lower cutoff if γ > 2.
- $ightharpoonup \gamma < 2$: Typical sample is large.
- $\sim \gamma > 2$: Typical sample is small.

Insert question from assignment 1 (⊞)





21 of 36

Exhibit A:

• Given $P(x) = cx^{-\gamma}$ with $0 < x_{min} < x < x_{max}$, the mean is $(\gamma \neq 2)$:

$$\langle x
angle = rac{c}{2-\gamma} \left(x_{
m max}^{2-\gamma} - x_{
m min}^{2-\gamma}
ight).$$

- ▶ Mean 'blows up' with upper cutoff if γ < 2.
- ▶ Mean depends on lower cutoff if γ > 2.
- $ightharpoonup \gamma < 2$: Typical sample is large.
- $ightharpoonup \gamma > 2$: Typical sample is small.

Insert question from assignment 1 (⊞)

Examples

Wild vs. Mild CCDFs

Zipf's law

Zipf ⇔ CCDF





Exhibit A:

• Given $P(x) = cx^{-\gamma}$ with $0 < x_{min} < x < x_{max}$, the mean is $(\gamma \neq 2)$:

$$\langle x
angle = rac{c}{2-\gamma} \left(x_{
m max}^{2-\gamma} - x_{
m min}^{2-\gamma}
ight).$$

- ▶ Mean 'blows up' with upper cutoff if γ < 2.
- ▶ Mean depends on lower cutoff if $\gamma > 2$.
- $ightharpoonup \gamma < 2$: Typical sample is large.
- $ightharpoonup \gamma > 2$: Typical sample is small.

Insert question from assignment 1 (⊞)

Wild vs. Mild CCDFs

Zipf's law

Zipf ⇔ CCDF





- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- ► Compare to a Gaussian, exponential, etc.

- mean is finite (depends on lower cutoff)
- σ^2 = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite
- ▶ If γ > 3, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF





- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- Compare to a Gaussian, exponential, etc.

- mean is finite (depends on lower cutoff).
- σ^2 = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- ▶ If γ > 3, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Lipi S law

Zipf ⇔ CCDF





- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- ► Compare to a Gaussian, exponential, etc.

- mean is finite (depends on lower cutoff)
- $ightharpoonup \sigma^2$ = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- If $\gamma >$ 3, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

.....





Moments:

- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- ► Compare to a Gaussian, exponential, etc.

For many real size distributions: $2 < \gamma < 3$

- mean is finite (depends on lower cutoff)
- σ^2 = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- If $\gamma > 3$, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF





Moments:

- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- ► Compare to a Gaussian, exponential, etc.

For many real size distributions: $2 < \gamma < 3$

- mean is finite (depends on lower cutoff)
- σ^2 = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- If $\gamma > 3$, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- ▶ Compare to a Gaussian, exponential, etc.

- mean is finite (depends on lower cutoff)
- $ightharpoonup \sigma^2$ = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- If $\gamma >$ 3, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF





- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- Compare to a Gaussian, exponential, etc.

- mean is finite (depends on lower cutoff)
- $ightharpoonup \sigma^2$ = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- If $\gamma > 3$, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF





Moments:

- All moments depend only on cutoffs.
- ▶ No internal scale that dominates/matters.
- ► Compare to a Gaussian, exponential, etc.

For many real size distributions: $2 < \gamma < 3$

- mean is finite (depends on lower cutoff)
- σ^2 = variance is 'infinite' (depends on upper cutoff)
- Width of distribution is 'infinite'
- ▶ If γ > 3, distribution is less terrifying and may be easily confused with other kinds of distributions.

Insert question from assignment 1 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





- ► Variance is nice analytically...
- ► Another measure of distribution width:

Mean average deviation (MAD) = $\langle |x - \langle x \rangle| \rangle$

▶ For a pure power law with 2 < γ < 3:

$$\langle |x - \langle x \rangle| \rangle$$
 is finite.

- But MAD is mildly unpleasant analytically...
- ▶ We still speak of infinite 'width' if γ < 3.

Insert question from assignment 2 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF







Standard deviation is a mathematical convenience:

- ► Variance is nice analytically...
- ► Another measure of distribution width:

Mean average deviation (MAD) =
$$\langle |x - \langle x \rangle| \rangle$$

▶ For a pure power law with 2 < γ < 3:

$$\langle |x - \langle x \rangle| \rangle$$
 is finite.

- But MAD is mildly unpleasant analytically...
- ▶ We still speak of infinite 'width' if γ < 3.

Insert question from assignment 2 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF





Standard deviation is a mathematical convenience:

- Variance is nice analytically...
- ► Another measure of distribution width:

Mean average deviation (MAD) =
$$\langle |x - \langle x \rangle| \rangle$$

▶ For a pure power law with 2 < γ < 3:

$$\langle |x - \langle x \rangle| \rangle$$
 is finite.

- ▶ But MAD is mildly unpleasant analytically...
- ▶ We still speak of infinite 'width' if γ < 3.

Insert question from assignment 2 (⊞)



Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





Standard deviation is a mathematical convenience:

- Variance is nice analytically...
- ► Another measure of distribution width:

Mean average deviation (MAD) =
$$\langle |x - \langle x \rangle| \rangle$$

▶ For a pure power law with 2 < γ < 3:

$$\langle |x - \langle x \rangle| \rangle$$
 is finite.

- But MAD is mildly unpleasant analytically...
- ▶ We still speak of infinite 'width' if γ < 3.

Insert question from assignment 2 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





- Variance is nice analytically...
- ▶ Another measure of distribution width:

Mean average deviation (MAD) = $\langle |x - \langle x \rangle| \rangle$

▶ For a pure power law with 2 < γ < 3:

$$\langle |x - \langle x \rangle| \rangle$$
 is finite.

- But MAD is mildly unpleasant analytically...
- ▶ We still speak of infinite 'width' if γ < 3.

Insert question from assignment 2 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF





- ▶ Variance is nice analytically...
- ▶ Another measure of distribution width:

Mean average deviation (MAD) = $\langle |x - \langle x \rangle| \rangle$

▶ For a pure power law with 2 < γ < 3:

$$\langle |x - \langle x \rangle| \rangle$$
 is finite.

- But MAD is mildly unpleasant analytically...
- ▶ We still speak of infinite 'width' if γ < 3.

Insert question from assignment 2 (⊞)

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF





Given $P(x) \sim cx^{-\gamma}$:

► We can show that after *n* samples, we expect the largest sample to be

$$x_1 \gtrsim c' n^{1/(\gamma-1)}$$

- ➤ Sampling from a finite-variance distribution gives a much slower growth with *n*.
- e.g., for $P(x) = \lambda e^{-\lambda x}$, we find

$$x_1 \gtrsim \frac{1}{\lambda} \ln n$$
.

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References



Insert question from assignment 2 (⊞)



Given $P(x) \sim cx^{-\gamma}$:

▶ We can show that after *n* samples, we expect the largest sample to be

$$x_1 \gtrsim c' n^{1/(\gamma-1)}$$

- ➤ Sampling from a finite-variance distribution gives a much slower growth with *n*.
- e.g., for $P(x) = \lambda e^{-\lambda x}$, we find

$$x_1 \gtrsim \frac{1}{\lambda} \ln n$$
.

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References



Insert question from assignment 2 (⊞)



Given $P(x) \sim cx^{-\gamma}$:

► We can show that after *n* samples, we expect the largest sample to be

$$x_1 \gtrsim c' n^{1/(\gamma-1)}$$

- ➤ Sampling from a finite-variance distribution gives a much slower growth with *n*.
- e.g., for $P(x) = \lambda e^{-\lambda x}$, we find

$$x_1 \gtrsim \frac{1}{\lambda} \ln n$$
.

Definition

Examples

Wild vs. Mild

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References



Insert question from assignment 2 (⊞)



$$P_{\geq}(x) = P(x' \geq x) = 1 - P(x' < x)$$

$$= \int_{x'=x}^{\infty} P(x') \mathrm{d}x$$

$$\propto \int_{y'=y}^{\infty} (x')^{-\gamma} \mathrm{d}x'$$

$$= \frac{1}{-\gamma+1} (x')^{-\gamma+1} \Big|_{x'=x}^{\infty}$$

$$\propto x^{-\gamma+1}$$

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







$$P_{\geq}(x) = P(x' \geq x) = 1 - P(x' < x)$$

$$= \int_{x'=x}^{\infty} P(x') \mathrm{d}x'$$

$$\propto \int_{y'-y}^{\infty} (x')^{-\gamma} dx'$$

)

$$= \frac{1}{-\gamma+1} (x')^{-\gamma+1} \Big|_{x'=x}^{\infty}$$

$$\propto x^{-\gamma+1}$$

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





$$P_{\geq}(x) = P(x' \geq x) = 1 - P(x' < x)$$

$$=\int_{x'=x}^{\infty}P(x')\mathrm{d}x'$$

$$\propto \int_{x'=x}^{\infty} (x')^{-\gamma} \mathrm{d}x'$$

$$= \frac{1}{-\gamma+1} (x')^{-\gamma+1} \Big|_{x'=1}^{\infty}$$

$$\propto x^{-\gamma+1}$$

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







$$P_{\geq}(x) = P(x' \geq x) = 1 - P(x' < x)$$

$$= \int_{x'=x}^{\infty} P(x') \mathrm{d}x'$$

$$\propto \int_{x'=x}^{\infty} (x')^{-\gamma} \mathrm{d}x'$$

$$= \frac{1}{-\gamma+1} (x')^{-\gamma+1} \Big|_{x'=1}^{\infty}$$

$$\propto x^{-\gamma+1}$$

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







CCDF:

$$P_{\geq}(x) = P(x' \geq x) = 1 - P(x' < x)$$

$$= \int_{x'=x}^{\infty} P(x') \mathrm{d}x'$$

•

$$\propto \int_{x'=x}^{\infty} (x')^{-\gamma} \mathrm{d}x'$$

$$= \frac{1}{-\gamma+1} (x')^{-\gamma+1} \Big|_{x'=x}^{\infty}$$

$$\propto x^{-\gamma+1}$$

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







CCDF:

$$P_{\geq}(x) = P(x' \geq x) = 1 - P(x' < x)$$

$$= \int_{x'=x}^{\infty} P(x') \mathrm{d}x'$$

$$\propto \int_{x'=x}^{\infty} (x')^{-\gamma} \mathrm{d}x'$$

1

$$= \left. \frac{1}{-\gamma + 1} (x')^{-\gamma + 1} \right|_{x' = x}^{\infty}$$

ì

$$\propto x^{-\gamma+1}$$

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

__~







CCDF:

$$P_{\geq}(x) \propto x^{-\gamma+1}$$

- ▶ Use when tail of *P* follows a power law.
- Increases exponent by one.
- Useful in cleaning up data.

Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







CCDF:

>

$$P_{\geq}(x) \propto x^{-\gamma+1}$$

- ▶ Use when tail of *P* follows a power law.
- Increases exponent by one.
- ▶ Useful in cleaning up data.

Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







CCDF:

>

$$P_{\geq}(x) \propto x^{-\gamma+1}$$

- ▶ Use when tail of *P* follows a power law.
- Increases exponent by one.
- ▶ Useful in cleaning up data.

Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





CCDF:

>

$$P_{\geq}(x) \propto x^{-\gamma+1}$$

- Use when tail of P follows a power law.
- Increases exponent by one.
- Useful in cleaning up data.

Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





Power Law Size Distributions

CCDF:


PDF:

 $\log_{10} N_q$

$$P_{\geq}(x) \propto x^{-\gamma+1}$$

- ▶ Use when tail of P follows a power law.
- Increases exponent by one.
- ► Useful in cleaning up data.

 $\log_{10} q$



efinition

Examples

Wild vs. Mild

CCDFs

Zipf's law $Zipf \Leftrightarrow CCDF$







 $P_{>}(k) = P(k' \geq k)$

Discrete variables:

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References

▶ Use integrals to approximate sums.





CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References

Discrete variables:

$$P_{\geq}(k) = P(k' \geq k)$$

$$=\sum_{k'=k}^{\infty}P(k)$$

Use integrals to approximate sums.





Discrete variables:

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF

References

 $P_{>}(k) = P(k' \geq k)$

 $r \geq (\kappa) - r (\kappa \geq \kappa)$

$$=\sum_{k'=k}^{\infty}P(k)$$

$$\propto k^{-\gamma+1}$$

▶ Use integrals to approximate sums.







Discrete variables:

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$

References

 $P_{>}(k) = P(k' \geq k)$

$$=\sum_{k'=k}^{\infty}P(k)$$

$$\propto k^{-\gamma+1}$$

Use integrals to approximate sums.





George Kingsley Zipf:

- Noted various rank distributions followed power laws, often with exponent -1 (word frequency, city sizes...)
- ► Zipf's 1949 Magnum Opus (⊞):
 "Human Behaviour and the Principle of Least-Effort" [12]
- ▶ We'll study Zipf's law in depth...





George Kingsley Zipf:

- Noted various rank distributions followed power laws, often with exponent -1 (word frequency, city sizes...)
- ➤ Zipf's 1949 Magnum Opus (⊞):
 "Human Behaviour and the Principle of Least-Effort" [12]
- ► We'll study Zipf's law in depth...





George Kingsley Zipf:

- Noted various rank distributions followed power laws, often with exponent -1 (word frequency, city sizes...)
- ➤ Zipf's 1949 Magnum Opus (⊞):
 "Human Behaviour and the Principle of Least-Effort" [12]
- ▶ We'll study Zipf's law in depth...





Examples

Wild vs. Mild CCDFs

Zipf's law

Zipf ⇔ CCDF

Zipf's way:

- Given a collection of entities, rank them by size, largest to smallest.
- x_r = the size of the rth ranked entity.
- ightharpoonup r = 1 corresponds to the largest size.
- \triangleright Example: x_1 could be the frequency of occurrence of
- Zipf's observation:







- ► Given a collection of entities, rank them by size, largest to smallest.
- x_r = the size of the *r*th ranked entity.
- ightharpoonup r = 1 corresponds to the largest size.
- \triangleright Example: x_1 could be the frequency of occurrence of the most common word in a text.
- ► Zipf's observation:

 $x_r \propto r^{-\alpha}$

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $Zipf \Leftrightarrow CCDF$







- Given a collection of entities, rank them by size, largest to smallest.
- \rightarrow x_r = the size of the rth ranked entity.
- ightharpoonup r = 1 corresponds to the largest size.
- ► Example: *x*₁ could be the frequency of occurrence of the most common word in a text.
- ► Zipf's observation:

 $X_r \propto r^{-\alpha}$

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







- Given a collection of entities, rank them by size, largest to smallest.
- \rightarrow x_r = the size of the rth ranked entity.
- ightharpoonup r = 1 corresponds to the largest size.
- ► Example: *x*₁ could be the frequency of occurrence of the most common word in a text.
- ➤ Zipf's observation:

$$x_r \propto r^{-\alpha}$$

Definition

Examples

Wild vs. Mild

CCDFs


Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$





Brown Corpus (1,015,945 words):



- ► The, of, and, to, a, ... = 'objects'
- ► 'Size' = word frequency

Definition

Examples

Wild vs. Mild

CCDFs


Zipf's law

Zipf ⇔ CCDF





Brown Corpus (1,015,945 words):



- ► The, of, and, to, a, ... = 'objects'
- ► 'Size' = word frequency
- Beep: CCDF and Zipf plots are related...

Definition

Examples

Wild vs. Mild

Zipf's law


Zipf ⇔ CCDF








Brown Corpus (1,015,945 words):



Zipf (axes flipped):



- ► The, of, and, to, a, ... = 'objects'
- 'Size' = word frequency
- Beep: CCDF and Zipf plots are related...

Definition

Examples

Wild vs. Mild

Zipf's law

Zipf ⇔ CCDF







- ▶ $NP_{\geq}(x)$ = the number of objects with size at least x where N = total number of objects.
- ▶ If an object has size x_r , then $NP_{\geq}(x_r)$ is its rank r.
- ► So

$$x_r \propto r^{-\alpha} = (NP_{\geq}(x_r))^{-\alpha}$$

$$\propto x_r^{(-\gamma+1)(-\alpha)}$$
 since $P_{\geq}(x) \sim x^{-\gamma+1}$.

We therefore have $1 = (-\gamma + 1)(-\alpha)$ or:

$$\alpha = \frac{1}{\gamma - 1}$$

A rank distribution exponent of $\alpha = 1$ corresponds to a size distribution exponent $\gamma = 2$.

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







- NP_≥(x) = the number of objects with size at least x where N = total number of objects.
- ▶ If an object has size x_r , then $NP_{\geq}(x_r)$ is its rank r.
- ► So

$$x_r \propto r^{-\alpha} = (NP_{\geq}(x_r))^{-\alpha}$$

$$\propto x_r^{(-\gamma+1)(-\alpha)}$$
 since $P_{\geq}(x) \sim x^{-\gamma+1}$.

We therefore have $1 = (-\gamma + 1)(-\alpha)$ or:

$$\alpha = \frac{1}{\gamma - 1}$$

A rank distribution exponent of $\alpha = 1$ corresponds to a size distribution exponent $\gamma = 2$.

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







- NP_≥(x) = the number of objects with size at least x where N = total number of objects.
- ▶ If an object has size x_r , then $NP_{\geq}(x_r)$ is its rank r.
- ▶ So

$$x_r \propto r^{-\alpha} = (NP_{\geq}(x_r))^{-\alpha}$$

$$\propto x_r^{(-\gamma+1)(-\alpha)}$$
 since $P_{\geq}(x) \sim x^{-\gamma+1}$.

We therefore have $1 = (-\gamma + 1)(-\alpha)$ or:

$$\alpha = \frac{1}{\gamma - 1}$$

A rank distribution exponent of $\alpha = 1$ corresponds to a size distribution exponent $\gamma = 2$.

Definition

xamples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







- NP_≥(x) = the number of objects with size at least x where N = total number of objects.
- ▶ If an object has size x_r , then $NP_{\geq}(x_r)$ is its rank r.
- ► So

$$x_r \propto r^{-\alpha} = (NP_{\geq}(x_r))^{-\alpha}$$

$$\propto x_r^{(-\gamma+1)(-\alpha)}$$
 since $P_{\geq}(x) \sim x^{-\gamma+1}$.

We therefore have $1 = (-\gamma + 1)(-\alpha)$ or:

$$\alpha = \frac{1}{\gamma - 1}$$

A rank distribution exponent of $\alpha = 1$ corresponds to a size distribution exponent $\gamma = 2$.

Definition

xamples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







- NP_≥(x) = the number of objects with size at least x where N = total number of objects.
- ▶ If an object has size x_r , then $NP_{\geq}(x_r)$ is its rank r.
- ► So

$$x_r \propto r^{-\alpha} = (NP_{\geq}(x_r))^{-\alpha}$$

$$\propto x_r^{(-\gamma+1)(-\alpha)}$$
 since $P_{\geq}(x) \sim x^{-\gamma+1}$.

We therefore have $1 = (-\gamma + 1)(-\alpha)$ or:

$$\alpha = \frac{1}{\gamma - 1}$$

A rank distribution exponent of $\alpha = 1$ corresponds to a size distribution exponent $\gamma = 2$.

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







- NP_≥(x) = the number of objects with size at least x where N = total number of objects.
- ▶ If an object has size x_r , then $NP_{\geq}(x_r)$ is its rank r.
- ► So

$$x_r \propto r^{-\alpha} = (NP_{\geq}(x_r))^{-\alpha}$$

$$\propto x_r^{(-\gamma+1)(-\alpha)}$$
 since $P_{\geq}(x) \sim x^{-\gamma+1}$.

We therefore have $1 = (-\gamma + 1)(-\alpha)$ or:

$$\alpha = \frac{1}{\gamma - 1}$$

A rank distribution exponent of $\alpha = 1$ corresponds to a size distribution exponent $\gamma = 2$.

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law


Zipf ⇔ CCDF





The Don. (⊞)

Extreme deviations in test cricket:



Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$








The Don. (⊞)

Extreme deviations in test cricket:



Don Bradman's batting average (⊞) = 166% next best. Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

 $\mathsf{Zipf} \Leftrightarrow \mathsf{CCDF}$







[2] A.-L. Barabási and R. Albert. Emergence of scaling in random networks. Science, 286:509–511, 1999. pdf (⊞)

[3] K. Christensen, L. Danon, T. Scanlon, and P. Bak. Unified scaling law for earthquakes.

Proc. Natl. Acad. Sci., 99:2509–2513, 2002. pdf (⊞)

[4] P. Grassberger.
Critical behaviour of the Drossel-Schwabl forest fire model.

New Journal of Physics, 4:17.1–17.15, 2002. pdf (⊞)

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF





Ann. Phys., 58:577-, 1917.

[6] M. E. J. Newman.

Power laws, pareto distributions and zipf's law.

Contemporary Physics, pages 323–351, 2005.

pdf (⊞)

[7] D. J. d. S. Price. Networks of scientific papers. Science, 149:510–515, 1965. pdf (⊞)

[8] D. J. d. S. Price.

A general theory of bibliometric and other cumulative advantage processes.

J. Amer. Soc. Inform. Sci., 27:292-306, 1976.

Definition

Examples

Wild vs. Mild

CCDFs

Zipf's law

Zipf ⇔ CCDF







References III

[9] L. F. Richardson.Variation of the frequency of fatal quarrels with magnitude.

J. Amer. Stat. Assoc., 43:523–546, 1949. pdf (⊞)

[10] H. A. Simon.
On a class of skew distribution functions.
Biometrika, 42:425–440, 1955. pdf (⊞)

[11] N. N. Taleb.

The Black Swan.

Random House, New York, 2007.

[12] G. K. Zipf.

Human Behaviour and the Principle of Least-Effort.

Addison-Wesley, Cambridge, MA, 1949.

Power Law Size Distributions

Definition

Examples

Wild vs. Mild

CCDFs
Zipf's law

Zipf ⇔ CCDF



