

**Complex Networks, CSYS/MATH 303**  
**University of Vermont, Spring 2014**  
**Assignment 3 • code name: Skipperdee** ↗

**Dispersed:** Thursday, February 6, 2014.

**Due:** By start of lecture, 2:30 pm, Thursday, February 13, 2014.

*Some useful reminders:*

**Instructor:** Peter Dodds

**Office:** Farrell Hall, second floor, Trinity Campus

**E-mail:** peter.dodds@uvm.edu

**Office hours:** 3:45 pm to 4:15 pm, Tuesday, and 12:45 pm to 2:15 pm, Wednesday

**Course website:** <http://www.uvm.edu/~pdodds/teaching/courses/2014-01UVM-303>

---

All parts are worth 3 points unless marked otherwise. Please show all your working clearly and list the names of others with whom you collaborated.

Graduate students are requested to use  $\LaTeX$  (or related  $\TeX$  variant).

---

**Supply networks and allometry:**

1. From lectures on Supply Networks:

Show that for large  $V$  and  $0 < \epsilon < 1/2$

$$\min V_{\text{net}} \propto \int_{\Omega_{d,D}(V)} \rho \|\vec{x}\|^{1-2\epsilon} d\vec{x} \sim \rho V^{1+\gamma_{\max}(1-2\epsilon)}$$

Reminders: we defined  $L_i = c_i^{-1} V^{\gamma_i}$  where  $\gamma_1 + \gamma_2 + \dots + \gamma_d = 1$ ,  $\gamma_1 = \gamma_{\max} \geq \gamma_2 \geq \dots \geq \gamma_d$ , and  $c = \prod_i c_i \leq 1$  is a shape factor.

Hints: assume the first  $k$  lengths scale in the same way with

$\gamma_1 = \dots = \gamma_k = \gamma_{\max}$ , and write  $\|\vec{x}\| = (x_1^2 + x_2^2 + \dots + x_d^2)^{1/2}$ .

2. Consider a set of rectangular areas with side lengths  $L_1$  and  $L_2$  such that  $L_1 \propto A^{\gamma_1}$  and  $L_2 \propto A^{\gamma_2}$  where  $A$  is area and  $\gamma_1 + \gamma_2 = 1$ . Assume  $\gamma_1 > \gamma_2$  and that  $\epsilon = 0$ .


Now imagine that material has to be distributed from a central source in each of these areas to sinks distributed with density  $\rho(A)$ , and that these sinks draw the same amount of material per unit time independent of  $L_1$  and  $L_2$ .

Find an exact form for how the volume of the most efficient distribution network scales with overall area  $A = L_1 L_2$ . (Hint: you will have to set up a double integration over the rectangle.)

If network volume must remain a constant fraction of overall area, determine the maximal scaling of sink density  $\rho$  with  $A$ .

Extra hints:

- Integrate over triangles as follows.
- You need to only perform calculations for one triangle.



3. (a) For a family of  $d$ -dimensional regions, with scaling as per Question 1, determine, to leading order, the scaling of hyper-surface area  $S$  with volume  $V$ . In other words, find the exponent  $\beta$  in  $S \propto V^\beta$  as  $V \rightarrow \infty$ . Assume that nothing peculiar happens with the shapes (as we have always implicitly done), in that there is no fractal roughening. Hint: figure out how the circumference for the rectangles in the previous question scales with area  $A$ . For  $d$  dimensions, think about how the hyper-surface area of a hyperrectangle (or orthotope) would scale.
- (b) For general  $d$ , what is the minimum and maximum possible values of  $\beta$  and for what values of the  $\gamma_i$  does these extrema occur?